

PDF issue: 2025-04-28

Bombardier beetles repel invasive bullfrogs

Sugiura, Shinji Date, Tomoki

(Citation) PeerJ,10:e13805

(Issue Date) 2022-09-15

(Resource Type) journal article

(Version) Version of Record

(Rights)

© 2022 Sugiura and Date This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, reproduction and adaptation in any medium and for any purpose provided that it is properly attributed...

(URL)

https://hdl.handle.net/20.500.14094/0100476928



Peer

Submitted 7 April 2022 Accepted 7 July 2022 Published 15 September 2022

Corresponding author Shinji Sugiura, ssugiura@people.kobe-u.ac.jp, sugiura.shinji@gmail.com

Academic editor Ann Hedrick

Additional Information and Declarations can be found on page 12

DOI 10.7717/peerj.13805

Copyright 2022 Sugiura and Date

Distributed under Creative Commons CC-BY 4.0

OPEN ACCESS

Bombardier beetles repel invasive bullfrogs

Shinji Sugiura and Tomoki Date

Graduate School of Agricultural Science, Kobe University, Kobe, Hyogo, Japan

ABSTRACT

Invasive non-native predators negatively affect native species; however, some native species can survive the predation pressures of invasive species by using pre-existing antipredator strategies or evolving defenses against invasive predators. The American bullfrog Lithobates catesbeianus (Anura: Ranidae) has been intentionally introduced to many countries and regions, and has impacted native animals through direct predation. Bombardier beetles (Coleoptera: Carabidae: Brachininae: Brachinini) discharge chemicals at a temperature of approximately 100 °C from the tip of the abdomen when they are attacked by predators. This "bombing" can successfully repel predators. However, adults of a native bombardier beetle Pheropsophus (Stenaptinus) occipitalis jessoensis have been reportedly found in the gut contents of the introduced bullfrog L. catesbeianus in Japan. These records suggest that the invasive bullfrog L. catesbeianus attacks the native bombardier beetle P. occipitalis jessoensis under field conditions in Japan; however, the effectiveness of the bombing defense against invasive bullfrogs is unclear. To test the effectiveness of the bombing defense against bullfrogs, we investigated the behavioral responses of *L. catesbeianus* juveniles to *P.* occipitalis jessoensis adults under laboratory conditions. Contrary to previous gut content results, almost all the bullfrogs (96.3%) rejected bombardier beetles before swallowing them; 88.9% rejected the beetles after being bombed, and 7.4% stopped attacking the beetles before being bombed. Only 3.7% successfully swallowed and digested the beetle. All of the beetles collected from non-bullfrog-invaded sites could deter bullfrogs, suggesting that the pre-existing defenses of bombardier beetles played an essential role in repelling bullfrogs. When treated beetles that were unable to discharge hot chemicals were provided, 77.8% of bullfrogs successfully swallowed and digested the treated beetles. These results indicate that bombing is important for the successful defense of *P. occipitalis jessoensis* against invasive bullfrogs. Although invasive bullfrogs have reportedly impacted native insect species, P. occipitalis jessoensis has an existing defense mechanism strong enough to repel the invasive predators.

Subjects Animal Behavior, Ecology, Entomology, Evolutionary Studies, ZoologyKeywords Bombardier beetles, Brachinini, Carabidae, Chemical defences, Introduced predators, Invasive alien species

INTRODUCTION

Invasive non-native species negatively impact native biota (*Doherty et al., 2016; Sugiura, 2016; David et al., 2017*). In particular, invasive predators affect native communities and ecosystems through cascading effects (*Goldschmidt, Witte & Wanink, 1993; O'Dowd, Green & Lake, 2003; Kenis et al., 2009; David et al., 2017; Rogers et al., 2017; McGruddy*

et al., 2021). Because native prey species do not share a history with invasive predators (*Fritts & Rodda, 1998; Strauss, Lau & Carroll, 2006; Carthey & Banks, 2014*), many native species suffer predation by invasive species (*Goldschmidt, Witte & Wanink, 1993; Doherty et al., 2016; Sugiura, 2016*). However, some native species have survived the predation pressures of invasive species by using pre-existing antipredator strategies (*Davis, Epp & Gabor, 2012; Carthey & Banks, 2014*) or evolving defenses against invasive predators (*Vermeij, 1982; Strauss, Lau & Carroll, 2006*). Pre-existing antipredator defenses can play an important role in repelling invasive predators that have similar ecological traits to native predators (*Carthey & Banks, 2014; Melotto et al., 2021*). However, pre-existing defenses have received less attention than the evolution of anti-predator defenses in terms of native species' tolerance to invasive predators (*Strauss, Lau & Carroll, 2006*). Investigating the effectiveness of the pre-existing defenses of native species against invasive predators would enable a better understanding of how to mitigate the impacts of invasive predators on native species.

The American bullfrog Lithobates catesbeianus (Shaw) (formerly called Rana catesbeiana Shaw; Lowe et al., 2000) (Anura: Ranidae) has been intentionally introduced for various purposes to many countries and regions (western North America, South America, East and Southeast Asia, and Western Europe) from eastern North America (Ficetola et al., 2007; Ficetola, Thuiller & Miaud, 2007; Giovanelli, Haddad & Alexandrino, 2008; Bissattini & Vignoli, 2017; Groffen et al., 2019; Johovic et al., 2020). Eggs are laid in still water such as ponds (Govindarajulu, Price & Anholt, 2006). The larvae feed on algae, diatoms, cyanobacteria, protists, and tiny invertebrates in water (Kupferberg, 1997; Pryor, 2003; Ruibal & Laufer, 2012). Postmetamorphic juveniles and adults prey on various animals (including aquatic and terrestrial species) in and near water (*Hirai*, 2004; Govindarajulu, Price & Anholt, 2006; Dontchev & Matsui, 2016; Flynn, Kreofsky & Sepulveda, 2017; Laufer et al., 2021; Sarashina & Yoshida, 2021). Because bullfrog adults commonly reach a size (snout-vent length) of 180-200 mm (Werner, Wellborn & McPeek, 1995), they are able to swallow small vertebrates (e.g., fish, mammals, reptiles, and frogs) as well as invertebrates (Raney & Ingram, 1941; Stewart & Sandison, 1972; Bruneau & Magnin, 1980, Clarkson & DeVos, 1986, Govindarajulu, Price & Anholt, 2006, Flynn, Kreofsky & Sepulveda, 2017; Oda et al., 2019). Consequently, invasive bullfrogs have impacted native communities in invaded habitats (Kats & Ferrer, 2003; Li et al., 2011; Adriaens, Devisscher & Louette, 2013; Gobel, Laufer & Cortizas, 2019). Therefore, L. catesbeianus has been listed as one of the 100 "world's worst invaders" (Lowe et al., 2000). Many studies have investigated the gut or stomach contents of adult and juvenile bullfrogs in native (Raney & Ingram, 1941; Korschgen & Moyle, 1955; Fulk & Whitaker, 1968; Stewart & Sandison, 1972; Bruneau & Magnin, 1980; Werner, Wellborn & McPeek, 1995) and invaded (Clarkson & DeVos, 1986; Balfour & Morey, 1999; Krupa, 2002; Hirai, 2004, 2005; Wu et al., 2005; Hirai & Inatani, 2008; Mori, 2008; Silva et al., 2009; Barrasso et al., 2009; Leivas, Leivas & Moura, 2012; Boelter et al., 2012; Jancowski & Orchard, 2013; Ortíz-Serrato, Ruiz-Campos & Valdez-Villavicencio, 2014; Quiroga et al., 2015; Liu et al., 2015; Dontchev & Matsui, 2016; Flynn, Kreofsky & Sepulveda, 2017; Vrcibradic et al., 2017; Park et al., 2018; Bissattini, Buono & Vignoli, 2018, 2019; Oda et al., 2019; Matsumoto,

Suwabe & Karube, 2020; Laufer et al., 2021; Nakamura & Tominaga, 2021) ranges, with the results indicating that introduced bullfrogs frequently attack native animal species in invaded areas. However, few studies have directly observed how invasive bullfrogs can attack and swallow native prey. Investigating the attack behavior of bullfrogs would help to assess which native species suffer from bullfrog predation.

Carabidae is one of the most diverse families in Coleoptera. Carabid beetles have frequently been used as bioindicators (Rainio & Niemelä, 2003) and biocontrol agents (Kromp, 1999). Carabid adults also exhibit morphological, physiological, chemical, and behavioral defenses against predators (Giglio et al., 2021). For example, adult bombardier beetles (Coleoptera: Carabidae: Brachininae: Brachinini) discharge toxic chemicals (e.g., 1,4-benzoquinone and 2-methyl-1,4-benzoquinone) and water (vapor) at a temperature of approximately 100 °C (*i.e.*, bombing) from the tip of abdomen when they are attacked by predators (Video S1; Aneshansley et al., 1969; Kanehisa & Murase, 1977; Dean, 1979; Kanehisa, 1996; Eisner, Eisner & Siegler, 2005; Arndt et al., 2015; Sugiura, 2018, 2021). The hot chemicals can effectively protect the beetles from predators such as arthropods (Eisner, 1958; Eisner & Meinwald, 1966; Eisner & Dean, 1976; Eisner et al., 2006; Sugiura, 2021), amphibians (Eisner & Meinwald, 1966; Dean, 1980; Sugiura & Sato, 2018; Sugiura, 2018), reptiles (Bonacci et al., 2008), and birds (Kojima & Yamamoto, 2020). The bombardier beetle *Pheropsophus* (Stenaptinus) occipitalis jessoensis Morawitz (formerly called Pheropsophus jessoensis Morawitz; Sugiura & Sato, 2018; Sugiura, 2018, 2021), which is commonly found in grassland, farmland, and forest edge environments in Japan, Korea, China, and Vietnam (Habu & Sadanaga, 1965; Yahiro et al., 1992; Ishitani & Yano, 1994; Fujisawa, Lee & Ishii, 2012; Ohwaki, Kaneko & Ikeda, 2015; Fedorenko, 2021), has been frequently studied to investigate the effectiveness of bombing as an anti-predator defense (Sugiura, 2018; Sugiura & Sato, 2018; Kojima & Yamamoto, 2020; Sugiura, 2021). Pheropsophus occipitalis jessoensis can successfully deter birds (Kojima & Yamamoto, 2020), frogs (Sugiura, 2018), and praying mantises (Sugiura, 2021). However, adults of the bombardier beetle Ph. occipitalis jessoensis have been reportedly found in the stomach contents of field-collected bullfrogs in central Japan; for example, two adult beetles were found in a dead bullfrog (Mori, 2008) and an adult beetle was found in a juvenile bullfrog (Matsumoto, Suwabe & Karube, 2020). These records suggest that the invasive bullfrog L. catesbeianus attacks the native bombardier beetle Ph. occipitalis jessoensis under field conditions in Japan, but the bombing defense of Ph. occipitalis jessoensis against invasive bullfrogs remains unexplored. To test the effectiveness of the bombing defense against bullfrogs, we investigated how L. catesbeianus juveniles respond to Ph. occipitalis jessoensis adults under laboratory conditions. In addition, the responses of bullfrogs to Ph. occipitalis jessoensis collected from bullfrog-invaded sites were compared with those of beetles collected from non-invaded sites to investigate whether native bombardier beetles that coexist with invasive bullfrogs exhibit a stronger defense than beetles that do not coexist with bullfrogs.

MATERIALS AND METHODS

Study species

To investigate how bullfrogs respond to bombardier beetles under laboratory conditions, we used juveniles of the invasive bullfrog *L. catesbeianus* and adults of the bombardier beetle *Ph. occipitalis jessoensis*. We observed *Ph. occipitalis jessoensis* adults, *L. catesbeianus* juveniles, and native pond frogs *Pelophylax nigromaculatus* (Hallowell) (Anura: Ranidae) in the same grassland around a pond in Hyogo, central Japan, on the same date (Fig. 1). Therefore, bullfrog juveniles may frequently encounter adults of *Ph. occipitalis jessoensis* in grassland around ponds, lakes, and paddy fields in Japan.

Fifty-four juvenile bullfrogs (snout-vent length: 42.2-59.6 mm) were collected from grassland around a pond in Hyogo, Japan, between August and October 2021. The bombardier beetle *Ph. occipitalis jessoensis* was frequently found in this sampling site. The snout-vent length and body weight of each bullfrog were measured to the nearest 0.01 mm and 0.1 mg, using an electronic slide caliper (CD-S15C; Mitutoyo, Kanagawa, Japan) and an electronic balance (CPA64; Sartorius Japan K.K., Tokyo, Japan), respectively. Juvenile bullfrogs were maintained separately in small plastic cages (120×85 \times 130 mm, length \times width \times height) in the laboratory at 25 °C (cf. Sugiura, 2018, 2020b). Live mealworms (larvae of *Tenebrio molitor* Linnaeus (Coleoptera: Tenebrionidae)) were provided as food (cf. Sugiura, 2018, 2020b). Bullfrogs were starved for 24 h before the experiments to standardize their hunger level (cf. Sugiura, 2018, 2020b). Individual bullfrogs were not used repeatedly (cf. Sugiura, 2018, 2020b). Introduced bullfrogs have been designated as an "invasive alien species" in Japan. Therefore, transportation, laboratory keeping, and behavioral experiments of bullfrogs were performed with permission from the Kinki Regional Environmental Office of the Ministry of the Environment, Government of Japan (Number: 20000085).

Fifty-four adults of the bombardier beetle *Ph. occipitalis jessoensis* (body length: 15.2–20.2 mm) were collected from grasslands and farmlands in Hyogo (three sites), Shiga (one site), Kyoto (one site), and Shimane (one site), central Japan (Fig. S1), in July–September 2020 and May–October 2021; 39 and 15 beetles were collected from bullfrog-invaded sites (three sites in Hyogo and one in Shimane) and non-invaded sites (one site in Kyoto and one in Shiga), respectively (Fig. S1). All adult beetles displayed bombing when manually caught by our researchers under field conditions. Body length and body weight of each beetle were also measured. Bombardier beetles were maintained separately in small plastic cases (diameter: 85 mm; height: 25 mm) in the laboratory at 25 °C (cf. *Sugiura, 2018; Sugiura & Sato, 2018; Sugiura, 2021*). Dead larvae of *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) were provided as food (cf. *Sugiura, 2018; Sugiura, 2021*). Individual beetles were not used repeatedly (cf. *Sugiura, 2018; Sugiura, 2018; Sugiura, 2021*).

Experiments

Following the method of *Sugiura (2018, 2020b*), we investigated how bullfrogs can attack bombardier beetles in our laboratory (25 °C) between September and November 2021.



Figure 1 A bombardier beetle, an invasive bullfrog, and a native frog. (A) An adult bombardier beetle *Pheropsophus occipitalis jessoensis.* (B) A juvenile bullfrog *Lithobates catesbeianus.* (C) An adult pond frog *Pelophylax nigromaculatus.* These photographs were taken at the same site and microhabitat on the same date. Photo credit: Shinji Sugiura. Full-size DOI: 10.7717/peerj.13805/fig-1

Bullfrogs that ate mealworms >1 day before experiments were used. We were unable to sex either bullfrogs or beetles due to their age and morphology, respectively. First, we placed a bullfrog in a plastic cage ($120 \times 85 \times 130$ mm, length \times width \times height). Then, we placed an

adult beetle in the cage. We recorded the behavior of the bullfrog and the beetle using digital cameras (iPhone 12 Pro Max; Apple Inc., Cupertino, CA, USA; Handycam HDR-CX630V; Sony, Tokyo, Japan). When a bullfrog rejected a beetle, we played back the footage of the recorded behavior to investigate whether the rejection was due to bombing. When bombing sounds were heard or ejected vapor was seen, we considered that bombing forced the bullfrog to reject the beetle. When a bullfrog swallowed a beetle, we investigated whether it vomited the beetle (cf. *Sugiura, 2018; Sugiura & Sato, 2018*). Bullfrogs that did not vomit were considered to have digested the beetle. When a bullfrog several minutes after beetle, we provided a mealworm as a palatable prey to the bullfrog several minutes after beetle rejection to determine whether this rejection was due to satiation (cf. *Sugiura, 2018, 2020b*). In total, 27 bullfrogs and 27 beetles were used in this experiment.

To test the role of hot chemical ejection by bombardier beetles in deterring bullfrogs, we provided bullfrogs with treated *Ph. occipitalis jessoensis* that were unable to eject hot chemicals (thereafter, treated beetles; cf. *Sugiura & Sato*, 2018; *Sugiura*, 2021). Following the method of *Sugiura & Sato* (2018) and *Sugiura* (2021), we used forceps to repeatedly stimulate an adult *Ph. occipitalis jessoensis*. This treatment caused the beetle to release all the chemicals. Each beetle repeatedly bombed before exhausting its chemicals. We then used the same procedure as for the control beetles to observe whether a bullfrog successfully attacked the treated beetle in a transparent plastic case (length × width × height, $120 \times 85 \times 130$ mm). In total, 27 bullfrogs and 27 beetles were used in the experiments. The sample size was determined based on the previous study (*Sugiura, 2018*).

All experiments were undertaken in accordance with the Kobe University Animal Experimentation Regulations (Kobe University's Animal Care and Use Committee, 30–01). No bullfrogs were injured during the feeding experiments. Because the release of bullfrogs into the wild is banned in Japan, the bullfrogs used in this study were euthanized by CO² asphyxiation after all experiments were completed.

Data analysis

Fisher's exact test was used to compare the rejection rate of control beetles with that of treated beetles by bullfrogs; the rejection rate of bombardier beetles collected from invaded sites with that of beetles from non-invaded sites; and the rejection rate of control beetles by bullfrogs with that by native pond frogs. Odds ratios and 95% confidence intervals (CIs) were also calculated. Data from *Sugiura (2018)* were also used as the rejection rate by the native pond frog *Pe. nigromaculatus*. Welch's *t*-test was used to compare the body size of bullfrogs and bombardier beetles between control and treated experiments. A generalized linear model (GLM) with a binomial error distribution and logit link was used to determine the factors contributing to the rejection of bombardier beetles by bullfrogs was used as the response variable. Beetle weight, bullfrog weight, the beetle weight × bullfrog weight interaction, and beetle treatment (control or treated) were included as fixed factors. A quasi-binomial error distribution was used rather than a binomial error distribution, which is necessary if the residual degrees of freedom (cf. *Sugiura, 2018*). All the tests were

performed at the 0.05 significance level. All analyses were performed using R ver. 3.5.2 (*R Core Team, 2018*).

RESULTS

All bullfrogs (n = 27) opened their mouths to attack bombardier beetles (control beetles); however, 26 bullfrogs (96.3%) rejected bombardier beetles (Table 1). The rejection was not due to satiation because 25 (96.2%) of the bullfrogs that rejected beetles ate mealworms immediately after the rejection. Only one bullfrog (3.7%) successfully swallowed and digested the beetle (Table 1). The swallowed beetle did not bomb when attacked. This beetle was relatively old (its sampling date was the earliest among all beetles). Two bullfrogs (7.4%) rejected the beetles before being bombed; one bullfrog (3.7%)stopped attacking the beetle immediately after its tongue touched it, and one bullfrog (3.7%) spat out the beetle <1 s after taking it into its mouth (Table 1). Two bullfrogs (7.4%) were bombed before taking the beetles into their mouths and immediately stopped the attack (Table 1). Twenty-two bullfrogs (81.5%) were bombed within 5 s of taking the beetles into their mouths and then spat them out within 2 s after being bombed (Video S2; Fig. 2; Table 1). The collection sites of bombardier beetles did not influence the rejection rates by bullfrogs (Fisher's exact test, P = 1.0, odds ratio [95% CI] = ∞ [0.01283594– ∞]); 94.4% of the beetles (n = 18) collected from bullfrog-invaded sites and 100% of the beetles (n = 9) from non-invaded sites were rejected by bullfrogs. The behavioral responses of bullfrogs to bombardier beetles were compared with those of the native pond frog species Pe. nigromaculatus (Fig. 3). The rate of swallowing and rejection of beetles did not significantly differ between the two species (Fisher's exact test, P = 1.0, odds ratio [95% CI] = 0.506127 [0.008172-10.284953]), but the rate of rejection before bombing significantly differed between the two species (Fig. 3; P = 0.000005, odds ratio [95% CI] = 0.042546 [0.004042-0.229305]).

When treated beetles that were unable to bomb were provided, all bullfrogs (n = 27) attacked the beetles. Twenty-one bullfrogs (77.8%) successfully swallowed and digested treated beetles, while six bullfrogs (22.2%) spat out treated beetles within 7 s of taking them into their mouths (Table 1). All of the bullfrogs that rejected treated beetles (n = 6) ate mealworms after the rejection. The rejection rate of treated beetles by bullfrogs (22.2%) differed significantly from that of control beetles (96.3%; Table 1; Fisher's exact test, P = 0.00000002, odds ratio [95% CI] = 0.01246560 [0.00026351-0.10342453]).

The body lengths and weights of treated beetles were not significantly different from those of control beetles (Table 2; t-test, P = 0.16-0.86). The snout–vent lengths and weights of bullfrogs that attacked control beetles were not significantly different from those of bullfrogs that attacked treated beetles (Table 2; t-test, P = 0.50-0.66). The GLM results indicated that the rejection rate of bombardier beetles by bullfrogs was influenced by beetle treatment, but not by the body size of either bombardier beetles or bullfrogs (Table 3).

DISCUSSION

The American bullfrog *L. catesbeianus* can eat any animals smaller than itself (*Adriaens*, *Devisscher & Louette*, 2013). Consequently, introduced bullfrogs have negatively affected



Figure 2 Temporal sequence of the bullfrog *Lithobates catesbeianus* rejecting a control adult *Pheropsophus occipitalis jessoensis.* (A) 0 ms. (B) 375 ms. (C) 900 ms. (D) 2,200 ms. (E) 2,575 ms. (F) 2,625 ms. (G) 2,650 ms. (H) 3,475 ms. The bullfrog spat out the beetle after taking it into its mouth. Bombing by the beetle was audible and the ejected vapor (E) was observed just before the bullfrog spat out the beetle (see Video S2). Credit: Shinji Sugiura and Tomoki Date.

Full-size 🖾 DOI: 10.7717/peerj.13805/fig-2

native arthropods and amphibians through direct predation in invaded areas (*Kats & Ferrer*, 2003; *Li et al.*, 2011; *Adriaens, Devisscher & Louette*, 2013; *Gobel, Laufer & Cortizas, 2019*; *Groffen et al.*, 2019; *Nakamura & Tominaga*, 2021). Although the native bombardier beetle *Ph. occipitalis jessoensis* has reportedly been identified in the stomach contents of



Figure 3 Behavioral responses of the invasive bullfrog *Lithobates catesbeianus* and the native pond frog *Pelophylax nigromaculatus* to adults of the bombardier beetle *Pheropsophus occipitalis jessoensis*. Swallow: bullfrogs or frogs successfully swallowed control beetles. Reject before bombed: bullfrogs or frogs stopped attacking control beetles before being bombed. Reject after bombed: bullfrogs or frogs rejected control beetles after being bombed. The graph showing data for *Pe. nigromaculatus* was taken from *Sugiura (2018)*. Photo credit: Shinji Sugiura. Full-size DOI: 10.7717/peerj.13805/fig-3

Frog response ^a	Frog behavior ^b	Control beetles ^c % (<i>n</i>)	Treated beetles ^c % (<i>n</i>)
Eat	Swallow	3.7 (1)	77.8 (21)
Reject (subtotal)		96.3 (26)	22.2 (6)
Reject before bombed	Stop attack	3.7 (1)	0.0 (0)
	Spit out	3.7 (1)	22.2 (6)
Reject after bombed	Stop attack	7.4 (2)	_
	Spit out	81.5 (22)	_
Total		100.0 (27)	100.0 (27)

 Table 1 Responses of the bullfrog Lithobates catesbeianus to control and treated adults of the bombardier beetle Pheropsophus occipitalis jessoensis.

Notes:

^a Eat: bullfrogs successfully ate beetles. Reject before bombed: bullfrogs rejected beetles before or without being bombed. Reject after bombed: bullfrogs rejected beetles after being bombed.

^b Swallow: bullfrogs successfully swallowed beetles. Stop attack: bullfrogs stopped attacking beetles before taking them into their mouths. Spit out: bullfrogs spat out beetles after taking them into their mouths.

^c Control beetles and treated beetles are the *Pheropsophus occipitalis jessoensis* that were able and unable to discharge hot chemicals, respectively.

introduced bullfrogs in Japan (*Mori, 2008; Matsumoto, Suwabe & Karube, 2020*), our laboratory experiments showed that almost all bullfrogs rejected *Ph. occipitalis jessoensis* before swallowing them. Therefore, *Ph. occipitalis jessoensis* can successfully repel invasive bullfrogs using a chemical weapon. To our knowledge, this is the first study to demonstrate the successful defense of a native insect species against invasive bullfrogs. However, this study may reflect limited aspects of prey-predator interactions between native bombardier beetles and invasive bullfrogs, as it was not designed to assess the potential effects of

	1 1	1)	0		'
Species	Boy size	Treatment		Statistical comparison	
		Control beetles $n = 27$	Treated beetle $n = 27$	t value	P value
Bombardier beetle	Body length (mm) ^a	17.6 ± 0.2	17.6 ± 0.2	0.25	0.80
		(15.2–20.2)	(15.5–19.6)		
	Boy weight (mg) ^a	265.8 ± 12.4	241.7 ± 11.4	1.43	0.16
		(149.1-411.3)	(146.5-376.2)		
			268.9 ± 12.4	$(-0.18)^{c}$	(0.86) ^c
			$(164.4 - 409.9)^{\rm b}$		
Bullfrog	Snout-vent length (mm) ^a	48.2 ± 0.8	47.8 ± 0.7	0.44	0.66
		(43.5–59.6)	(42.2–57.3)		
	Body weight (mg) ^a	9206.6 ± 554.7	8720.9 ± 458.1	0.68	0.50
		(6136.9-18257.1)	(5575.6-16763.8)		

Table 2 Sizes of the bombardier beetle Pheropsophus occipitalis jessoensis and the bullfrog Lithobates catesbeianus used in this study.

Notes:

^a Values are the mean ± standard error (range: minimum-maximum).

^b Body weight of bombardier beetles before treatment.

^c Statistical result of a comparison between treated beetles (before treatment) and control beetles.

Table 3 Results of a generalized linear model (GLM) identifying factors affecting whether the bullfrog *Lithobates catesbeianus* rejected the bombardier beetle *Pheropsophus occipitalis jessoensis*.

Response variable	Explanatory variable (fixed effect)	Coefficient estimate	SE	t value	P value
Rejection ^a	Intercept	11.68	7.466	1.564	0.12418
	Beetle treatment ^b	-5.389	1.673	-3.222	0.00226
	Beetle size (weight)	-0.02395	0.0308	-0.777	0.44061
	Frog size (weight)	-0.0007511	0.000699	-1.075	0.28783
	Beetle size \times frog size	0.000002211	0.000003243	0.682	0.49872

Notes:

^a A quasi-binomial error distribution (rather than a binomial error distribution) was used because the residual deviance was smaller than the residual degrees of freedom. ^b Control beetles were used as a reference.

bullfrog size and learning on successful defenses of *Ph. occipitalis jessoensis*. Considering this limitation, we discuss the importance of bombing behavior as a pre-existing defense of *Ph. occipitalis jessoensis* against invasive bullfrogs, and the potential impact of invasive bullfrogs on native *Ph. occipitalis jessoensis*.

Some native species can evolve a tolerance to or defense against invasive predators (*Strauss, Lau & Carroll, 2006*). However, all adults of *Ph. occipitalis jessoensis* collected from non-bullfrog-invaded sites could successfully defend against bullfrogs, suggesting that the pre-existing defense of *Ph. occipitalis jessoensis* was strong enough to repel bullfrogs. Like invasive bullfrogs, the native pond frog *Pe. nigromaculatus* has been shown to frequently reject *Ph. occipitalis jessoensis* under laboratory conditions (*Sugiura, 2018*). Because both the native frog and the introduced bullfrog are frequently found in the same habitats in Japan (*Kambayashi et al., 2016; Sato, 2016; Tawa & Sagawa, 2017*), the defenses of *Ph. occipitalis jessoensis* that originally functioned against native frogs could play an important role in repelling invasive bullfrogs.

Sugiura (2018) showed that 67.9% of the native frog *Pe. nigromaculatus* rejected *Ph. occipitalis jessoensis* before being bombed. When dead adults of *Ph. occipitalis jessoensis* were provided, 71.4% of *Pe. nigromaculatus* rejected them (*Sugiura, 2018*). The native frog species stopped attacking live and dead *Ph. occipitalis jessoensis* immediately after their tongues contacted them, indicating that this frog species may avoid being bombed by detecting chemicals on the surface of the beetle (*Sugiura, 2018*). The present study showed that only 7.4% of bullfrogs rejected *Ph. occipitalis jessoensis* before being bombed. Therefore, bombing by *Ph. occipitalis jessoensis* is much more important for a successful defense against invasive bullfrogs than against native frogs. Unlike native frogs, bullfrogs may not use their tongue to detect a deterrent chemical or the physical characteristics of *Ph. occipitalis jessoensis*.

Adults of Ph. occipitalis jessoensis were found in the stomach contents of introduced bullfrogs in Japan (Mori, 2008; Matsumoto, Suwabe & Karube, 2020), although our results showed that almost all bullfrogs failed to eat adult Ph. occipitalis jessoensis. At least three factors may help to explain this inconsistency: high encounter rates between adult Ph. occipitalis jessoensis and bullfrogs; deficiency of defensive chemicals in old adult Ph. occipitalis jessoensis; and different body sizes of the bullfrogs used in the present and previous studies. First, because Ph. occipitalis jessoensis is commonly found in grassland and farmland around ponds, lakes, and paddy fields invaded by bullfrogs, adults of Ph. occipitalis jessoensis frequently encounter bullfrogs in Japan. The high encounter rate between Ph. occipitalis jessoensis and bullfrogs could result in successful predation events by bullfrogs even when the overall success rate of predation on *Ph. occipitalis jessoensis* by bullfrogs is low. To the second point, old adults of *Ph. occipitalis jessoensis* that are unable to produce enough defensive chemicals can easily be eaten by bullfrogs. In our experiment, the swallowed adult Ph. occipitalis jessoensis was relatively older than the other beetles and did not bomb when attacked. Lastly, Matsumoto, Suwabe & Karube (2020) found an adult *Ph. occipitalis jessoensis* in the stomach content of a juvenile bullfrog (snout-vent length: 83 mm) that was larger than the juveniles used in our experiments (snout-vent length: 43.4-59.6 mm). This suggests that Ph. occipitalis jessoensis may fail to defend itself against bullfrog adults and large juveniles. The importance of predator size for the successful defense of Ph. occipitalis jessoensis was suggested by Sugiura & Sato (2018) who showed that adult and large juvenile toads could more frequently eat adult Ph. occipitalis jessoensis than the small juveniles. However, juvenile bullfrogs of the size used in this study are much more abundant than the adults and large juveniles in invaded areas in Japan (Sato & Nishihara, 2017; Matsumoto, Suwabe & Karube, 2020). Therefore, unlike other native insect species, the native bombardier beetle Ph. occipitalis jessoensis may not suffer predation by invasive bullfrogs. However, no studies have quantitatively compared the abundance of Ph. occipitalis jessoensis between bullfrog-invaded and non-invaded areas. Further studies are needed to demonstrate the impacts of invasive bullfrogs on Ph. occipitalis jessoensis.

CONCLUSIONS

Some native animal species can tolerate invasive predators by evolving defenses against the predators (*Vermeij, 1982; Strauss, Lau & Carroll, 2006*) or using pre-existing defensive strategies (*Davis, Epp & Gabor, 2012; Carthey & Banks, 2014*). Although bombardier beetles possess chemical weapons to deter various types of predators (*Eisner, Eisner & Siegler, 2005; Sugiura, 2020a*), how they defend against invasive predators has been unclear. Our laboratory experiments demonstrated that the native bombardier beetle *Ph. occipitalis jessoensis* was able to repel invasive bullfrogs by bombing. Because *Ph. occipitalis jessoensis* can defend itself against the native pond frog *Pe. nigromaculatus* (*Sugiura, 2018*) and other native predators (*Sugiura & Sato, 2018; Kojima & Yamamoto, 2020; Sugiura, 2021*) in Japan, *Ph. occipitalis jessoensis* uses its pre-existing defense to defend against invasive bullfrogs, which occupy a similar niche to that of native pond frogs.

ACKNOWLEDGEMENTS

We are grateful to M. Hayashi for helping to collect bombardier beetles.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The following grant information was disclosed by the authors: Grant-in-Aid for Scientific Research (JSPS KAKENHI): 19K06073. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors: Grant-in-Aid for Scientific Research (JSPS KAKENHI): 19K06073.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Shinji Sugiura conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Tomoki Date performed the experiments, prepared figures and/or tables, and approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (*i.e.*, approving body and any reference numbers):

The authors followed Kobe University's Animal Experimentation Regulations and obtained approval from Kobe University's Animal Care and Use Committee (30–01).

Data Availability

The following information was supplied regarding data availability:

The raw data are available at figshare: Sugiura, Shinji; Date, Tomoki (2022): Data from: Bombardier beetles repel invasive bullfrogs. figshare. Dataset. https://doi.org/10.6084/m9. figshare.19439453.

Supplemental Information

Supplemental information for this article can be found online at http://dx.doi.org/10.7717/ peerj.13805#supplemental-information.

REFERENCES

- Adriaens T, Devisscher S, Louette G. 2013. Risk analysis report of non-native organisms in Belgium: risk analysis of American bullfrog Lithobates catesbeianus (Shaw). Brussel: Instituut voor Natuur- en Bosonderzoek.
- Aneshansley DT, Eisner T, Widom JM, Widom B. 1969. Biochemistry at 100 °C: explosive secretory discharge of bombardier beetles (*Brachinus*). *Science* 165(3888):61–63 DOI 10.1126/science.165.3888.61.
- Arndt EM, Moore W, Lee WK, Ortiz C. 2015. Mechanistic origins of bombardier beetle (Brachinini) explosion-induced defensive spray pulsation. *Science* 348(6234):563–567 DOI 10.1126/science.1261166.
- **Balfour PS, Morey SR. 1999.** Prey selection by juvenile bullfrogs in a constructed vernal pool complex. *Transactions of the Western Section of the Wildlife Society* **35**:34–40.
- Barrasso DA, Cajade R, Nenda S, Baloriani G, Herrera R. 2009. Introduction of the American bullfrog *Lithobates catesbeianus* (Anura: Ranidae) in natural and modified environments: an increasing conservation problem in Argentina. *South American Journal of Herpetology* 4(1):69–75 DOI 10.2994/057.004.0109.
- **Bissattini AM, Buono V, Vignoli L. 2018.** Field data and worldwide literature review reveal that alien crayfish mitigate the predation impact of the American bullfrog on native amphibians. *Aquatic Conservation: Marine and Freshwater Ecosystems* **28(6)**:1465–1475 DOI 10.1002/aqc.2978.
- Bissattini AM, Buono V, Vignoli L. 2019. Disentangling the trophic interactions between American bullfrogs and native anurans: complications resulting from post-metamorphic ontogenetic niche shifts. *Aquatic Conservation: Marine and Freshwater Ecosystems* 29(2):270–281 DOI 10.1002/aqc.3023.
- Bissattini AM, Vignoli L. 2017. Let's eat out, there's crayfish for dinner: American bullfrog niche shifts inside and outside native ranges and the effect of introduced crayfish. *Biological Invasions* 19(9):2633–2646 DOI 10.1007/s10530-017-1473-6.
- Boelter RA, Kaefer IL, Both C, Cechin S. 2012. Invasive bullfrogs as predators in a Neotropical assemblage: what frog species do they eat? *Animal Biology* 62(4):397–408 DOI 10.1163/157075612X634111.
- Bonacci T, Aloise G, Brandmayr P, Brandmayr TZ, Capula M. 2008. Testing the predatory behaviour of *Podarcis sicula* (Reptilia: Lacertidae) towards aposematic and non-aposematic preys. *Amphibia-Reptilia* 29(3):449–453 DOI 10.1163/156853808785111986.
- Bruneau M, Magnin E. 1980. Croissance, nutrition et reproduction des souaouarons Rana catesbeiana Shaw (Amphibia Anura) des Laurentides au nord de Montréal. Canadian Journal of Zoology 58(2):175–183 DOI 10.1139/z80-019.

- Carthey AJR, Banks PB. 2014. Naïveté in novel ecological interactions: lessons from theory and experimental evidence. *Biological Reviews* 89(4):932–949 DOI 10.1111/brv.12087.
- Clarkson RW, DeVos JCJ. 1986. The bullfrog, *Rana catesbeiana* Shaw, in the lower Colorado River, Arizona-California. *Journal of Herpetology* 20(1):42–49 DOI 10.2307/1564123.
- David P, Thébault E, Anneville O, Duyck PF, Chapuis E. 2017. Impacts of invasive species on food webs: a review of empirical data. *Advances in Ecological Research* 56(2):1–60 DOI 10.1016/bs.aecr.2016.10.001.
- Davis DR, Epp KJ, Gabor CR. 2012. Predator generalization decreases the effect of introduced predators in the San Marcos salamander, *Eurycea nana*. *Ethology* 118(12):1191–1197 DOI 10.1111/eth.12025.
- **Dean J. 1979.** Defensive reaction time of bombardier beetles: an investigation of the speed of a chemical defense. *Journal of Chemical Ecology* **5(5)**:691–701 DOI 10.1007/BF00986554.
- Dean J. 1980. Encounters between bombardier beetles and two species of toads (*Bufo americanus*, B. *marinus*): speed of prey-capture does not determine success. *Journal of Comparative Physiology* 135(1):41–50 DOI 10.1007/BF00660180.
- Doherty TS, Glen AS, Nimmo DG, Ritchie EG, Dickman CR. 2016. Invasive predators and global biodiversity loss. Proceedings of the National Academy of Sciences of the United States of America 113(40):11261–11265 DOI 10.1073/pnas.1602480113.
- **Dontchev K, Matsui M. 2016.** Food habits of the American bullfrog *Lithobates catesbeianus* in the city of Kyoto, central Japan. *Current Herpetology* **35(2)**:93–100 DOI 10.5358/hsj.35.93.
- Eisner T. 1958. The protective role of the spray mechanism of the bombardier beetle, *Brachynus* ballistarius Lec. Journal of Insect Physiology 2(3):215–220 DOI 10.1016/0022-1910(58)90006-4.
- Eisner T, Aneshansley DJ, del Campo ML, Eisner M, Frank JH, Deyrup M. 2006. Effect of bombardier beetle spray on a wolf spider: repellency and leg autotomy. *Chemoecology* 16(4):185–189 DOI 10.1007/s00049-006-0346-8.
- **Eisner T, Dean J. 1976.** Ploy and counterploy in predator-prey interactions: orb-weaving spiders versus bombardier beetles. *Proceedings of the National Academy of Sciences of the United States of America* **73(4)**:1365–1367 DOI 10.1073/pnas.96.17.9705.
- Eisner T, Eisner M, Siegler M. 2005. Secret weapons: defenses of insects, spiders, scorpions, and other many-legged creatures. Cambridge: The Belknap Press of the Harvard University Press.
- Eisner T, Meinwald J. 1966. Defensive secretions of arthropods. *Science* 153(3742):1341–1350 DOI 10.1126/science.153.3742.1341.
- Fedorenko DN. 2021. *Stenaptinus* (Coleoptera: Carabidae: Brachininae) of Vietnam. Note 3. *Russian Entomology Journal* 30(3):252–263 DOI 10.15298/rusentj.30.3.02.
- Ficetola GF, Coïc C, Detaint M, Berroneau M, Lorvelec O, Miaud C. 2007. Pattern of distribution of the American bullfrog *Rana catesbeiana* in Europe. *Biological Invasions* 9(7):767–772 DOI 10.1007/s10530-006-9080-y.
- Ficetola GF, Thuiller W, Miaud C. 2007. Prediction and validation of the potential global distribution of a problematic alien invasive species—the American bullfrog. *Diversity and Distributions* 13(4):476–485 DOI 10.1111/j.1472-4642.2007.00377.x.
- Flynn LM, Kreofsky TM, Sepulveda AJ. 2017. Introduced American bullfrog distribution and diets in Grand Teton National Park. Northwest Science 91(3):244–256 DOI 10.3955/046.091.0305.
- Fritts TH, Rodda GH. 1998. The role of introduced species in the degradation of island ecosystems: a case history of Guam. *Annual Review of Ecology and Systematics* 29:113–140 DOI 10.1146/ANNUREV.ECOLSYS.29.1.113.

- Fujisawa T, Lee CM, Ishii M. 2012. Species diversity of ground beetle assemblages in the distinctive landscapes of the Yodo River flowing through northern Osaka Prefecture, central Japan. Japanese Journal of Environmental Entomology and Zoology 23(2):89–100 DOI 10.11257/jjeez.23.89.
- **Fulk FD, Whitaker JO Jr. 1968.** The food of *Rana catebeiana* in three habitats in Owen Country, Indiana. *Proceedings of the Indiana Academy of Science* **78**:491–496.
- Giglio A, Vommaro ML, Brandmayr P, Talarico F. 2021. Pygidial glands in Carabidae, an overview of morphology and chemical secretion. *Life* 11(6):562 DOI 10.3390/life11060562.
- Giovanelli JGR, Haddad CFB, Alexandrino J. 2008. Predicting the potential distribution of the alien invasive American bullfrog (*Lithobates catesbeianus*) in Brazil. *Biological Invasions* 10(5):585–590 DOI 10.1007/s10530-007-9154-5.
- **Gobel N, Laufer G, Cortizas S. 2019.** Changes in aquatic communities recently invaded by a top predator: evidence of American bullfrogs in Aceguá, Uruguay. *Aquatic Sciences* **81(1)**:8 DOI 10.1007/s00027-018-0604-1.
- Goldschmidt T, Witte F, Wanink J. 1993. Cascading effects of the introduced Nile perch on the detritivorous/phytoplanktivorous species in the sublittoral areas of Lake Victoria. *Conservation Biology* 7(3):686–700 DOI 10.1046/j.1523-1739.1993.07030686.x.
- **Govindarajulu P, Price WS, Anholt BR. 2006.** Introduced bullfrogs (*Rana catesbeiana*) in western Canada: has their ecology diverged? *Journal of Herpetology* **40(2)**:249–260 DOI 10.1670/68-05A.1.
- Groffen J, Kong S, Jang Y, Borzée A. 2019. The invasive American bullfrog (*Lithobates catesbeianus*) in the Republic of Korea: history and recommendations for population control. Management of Biological Invasions 10(3):517–535 DOI 10.3391/mbi.2019.10.3.08.
- Habu A, Sadanaga K. 1965. Illustrations for identification of larvae of the Carabidae found in cultivated fields and paddy-fields (III). *Bulletin of the National Institute of Agricultural Sciences, Series C: Plant Pathology and Entomology* 19:81–216 [in Japanese with English summary].
- Hirai T. 2004. Diet composition of introduced bullfrog, *Rana catesbeiana*, in the Mizorogaike Pond of Kyoto, Japan. *Ecological Research* 19(4):375–380 DOI 10.1111/j.1440-1703.2004.00647.x.
- Hirai T. 2005. On the giant water bug, *Lethocerus deyrolli*, found in stomach contents of a bullfrog, *Rana catesbeiana. Bulletin of Kansai Organization for Nature Conservation* 27(1):57–58 (in Japanese with English summary).
- **Hirai T, Inatani Y. 2008.** Predation by *Rana catesbeiana* on an adult male of *R. porosa brevipoda*. *Bulletin of the Herpetological Society of Japan* **2008(1)**:6–7 [in Japanese].
- **Ishitani M, Yano K. 1994.** Species composition and seasonal activities of ground beetles (*Coleoptera*) in a fig orchard. *Japanese Journal of Entomology* **62(1)**:201–210.
- Jancowski K, Orchard SA. 2013. Stomach contents from invasive American bullfrogs *Rana* catesbeiana (= Lithobates catesbeianus) on southern Vancouver Island, British Columbia, Canada. *NeoBiota* 16(2):17–37 DOI 10.3897/neobiota.16.3806.
- Johovic I, Gama M, Banha F, Tricarico E, Anastácio PM. 2020. A potential threat to amphibians in the European Natura 2000 network: forecasting the distribution of the American bullfrog *Lithobates catesbeianus. Biological Conservation* 245(6):108551 DOI 10.1016/j.biocon.2020.108551.
- Kambayashi C, Uto T, Shioji T, Kurabayashi A, Shimizu N. 2016. Amphibian fauna in the Higashi-Hiroshima Campus, Hiroshima University. *Bulletin of the Hiroshima University Museum* 8:17–29 (in Japanese with English Abstract).

- Kanehisa K. 1996. Secretion of defensive substance by Carabidae and Brachinidae. *Bulletin of the Research Institute for Bioresources, Okayama University* 4(1):9–23 [in Japanese with English summary].
- Kanehisa K, Murase M. 1977. Comparative study of the pygidial defensive systems of carabid beetles. *Applied Entomology and Zoology* 12(3):225–235 DOI 10.1303/aez.12.225.
- Kats LB, Ferrer RP. 2003. Alien predators and amphibian declines: review of two decades of science and the transition to conservation. *Diversity and Distributions* 9(2):99–110 DOI 10.1046/j.1472-4642.2003.00013.x.
- Kenis M, Auger-Rozenberg MA, Roques A, Timms L, Péré C, Cock MJW, Settele J, Augustin S, Lopez-Vaamonde C. 2009. Ecological effects of invasive alien insects. *Biological Invasions* 11(1):21–45 DOI 10.1007/s10530-008-9318-y.
- Kojima W, Yamamoto R. 2020. Defense of bombardier beetles against avian predators. *The Science of Nature* 107(4):36 DOI 10.1007/s00114-020-01692-z.
- Korschgen LJ, Moyle DL. 1955. Food habits of the bullfrog in central Missouri farm ponds. *The American Midland Naturalist* 54(2):332–341 DOI 10.2307/2422571.
- Kromp B. 1999. Carabid beetles in sustainable agriculture: a review on pest control efficacy, cultivation impacts and enhancement. *Agriculture, Ecosystems and Environment* 74(1-3):187–228 DOI 10.1016/S0167-8809(99)00037-7.
- Krupa JJ. 2002. Temporal shift in diet in a population of American bullfrog (*Rana catesbeiana*) in Carlsbad Caverns National Park. *The Southwestern Naturalist* 47(3):461–467 DOI 10.2307/3672506.
- Kupferberg SJ. 1997. The role of larval diet in anuran metamorphosis. *American Zoologist* 37(2):146–159 DOI 10.1093/icb/37.2.146.
- Laufer G, Gobel N, Berazategui M, Zarucki M, Cortizas S, Soutullo A, Martinez-Debat C, de Sá RO. 2021. American bullfrog (*Lithobates catesbeianus*) diet in Uruguay compared with other invasive populations in Southern South America. North-Western Journal of Zoology 17(1):e211502.
- Leivas PT, Leivas FWT, Moura MO. 2012. Diet and trophic niche of *Lithobates catesbeianus* (Amphibia: Anura). *Zoologia* 29(5):405–412 DOI 10.1590/S1984-46702012000500003.
- Li Y, Ke Z, Wang Y, Blackburn TM. 2011. Frog community responses to recent American bullfrog invasions. *Current Zoology* 57(1):83–92 DOI 10.1093/czoolo/57.1.83.
- Liu X, Luo Y, Chen J, Guo Y, Bai C, Li Y. 2015. Diet and prey selection of the invasive American bullfrog (*Lithobates catesbeianus*) in southwestern China. *Asian Herpetological Research* 6(1):34–44 DOI 10.16373/j.cnki.ahr.140044.
- Lowe S, Browne M, Boudjelas S, De Poorter M. 2000. 100 of the world's worst invasive alien species: a selection from the global invasive species database. Auckland: IUCN/SSC Invasive Species Specialist Group (ISSG).
- Matsumoto R, Suwabe S, Karube H. 2020. Diet of *Xenopus laevis* and *Lithobates catesbeianus* trapped in Nakaogino Area, Atsugi, Kanagawa Prefecture, Japan. *Bulletin of the Kanagawa Prefectural Museum (Natural Science)* **49**:85–99 [in Japanese with English abstract].
- McGruddy RA, Howse MWF, Haywood J, Ward CJI, Staufer TB, Hayek-Williams M, Toft RJ, Lester PJ. 2021. Invasive paper wasps have strong cascading effects on the host plant of monarch butterflies. *Ecological Entomology* 46(2):459–469 DOI 10.1111/een.12992.
- Melotto A, Ficetola GF, Alari E, Romagnoli S, Manenti R. 2021. Visual recognition and coevolutionary history drive responses of amphibians to an invasive predator. *Behavioral Ecology* 32(6):1352–1362 DOI 10.1093/beheco/arab101.

- Mori I. 2008. Predation by introduced bullfrog *Rana catesbeiana* on a breeding male of *Rhacophorus schlegelii* and the other animals. *Bulletin of the Okayama Prefecture Nature Conservation Center* 16:61–62 [in Japanese].
- Nakamura Y, Tominaga A. 2021. Diet of the American bullfrog *Lithobates catesbeianus* naturalized on Okinawajima, Ryukyu Archipelago, Japan. *Current Herpetology* 40(1):40–53 DOI 10.5358/hsj.40.40.
- Oda FH, Guerra V, Grou E, de Lima LD, Proença HC, Gambale PG, Takemoto RM, Teixeira CP, Campião KM, Ortega JCG. 2019. Native anuran species as prey of invasive American bullfrog *Lithobates catesbeianus* in Brazil: a review with new predation records. *Amphibian & Reptile Conservation* 13(2):217–226.
- O'Dowd DJ, Green PT, Lake PS. 2003. Invasional 'meltdown' on an oceanic island. *Ecology Letters* 6(9):812–817 DOI 10.1046/j.1461-0248.2003.00512.x.
- **Ohwaki A, Kaneko Y, Ikeda H. 2015.** Seasonal variability in the response of ground beetles (Coleoptera: Carabidae) to a forest edge in a heterogeneous agricultural landscape in Japan. *European Journal of Entomology* **112(1)**:135–144 DOI 10.14411/eje.2015.022.
- Ortíz-Serrato L, Ruiz-Campos G, Valdez-Villavicencio JH. 2014. Diet of the exotic American bullfrog, *Lithobates catesbeianus* in a stream of northwestern Baja California, Mexico. *Western North American Naturalist* 74(1):116–122 DOI 10.3398/064.074.0112.
- Park CD, Lee CW, Lim JC, Yang BG, Lee JH. 2018. A study on the diet items of American bullfrog (*Lithobates catesbeianus*) in Ga-hang wetland. *Korea Korean Journal of Environment and Ecology* 32(1):55–65 [in Korean with English abstract] DOI 10.13047/KJEE.2018.32.1.55.
- Pryor GS. 2003. Growth rates and digestive abilities of bullfrog tadpoles (*Rana catesbeiana*) fed algal diets. *Journal of Herpetology* 37(3):560–566 DOI 10.1670/153-02N.
- Quiroga LB, Moreno MD, Cataldo AA, Aragón-Traverso JH, Pantano MV, Olivares JPS, Sanabria EA. 2015. Diet composition of an invasive population of *Lithobates catesbeianus* (American Bullfrog) from Argentina. *Journal of Natural History* **49**(27–28):1703–1716 DOI 10.1080/00222933.2015.1005711.
- **R Core Team. 2018.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. *Available at http://www.R-project.org/*.
- Rainio J, Niemelä J. 2003. Ground beetles (Coleoptera: Carabidae) as bioindicators. *Biodiversity* and Conservation 12(3):487–506 DOI 10.1023/A:1022412617568.
- Raney EC, Ingram WM. 1941. Growth of tagged frogs (*Rana catesbeiana* Shaw and *Rana clamitans* Daudin) under natural conditions. *The American Midland Naturalist* 26(1):201–206 DOI 10.2307/2420767.
- Rogers HS, Buhle ER, HilleRisLambers J, Fricke EC, Miller RH, Tewksbury JJ. 2017. Effects of an invasive predator cascade to plants via mutualism disruption. *Nature Communications* 8(1):14557 DOI 10.1038/ncomms14557.
- Ruibal M, Laufer G. 2012. Bullfrog *Lithobates catesbeianus* (Amphibia: Ranidae) tadpole diet: description and analysis for three invasive populations in Uruguay. *Amphibia-Reptilia* 33(3-4):355-363 DOI 10.1163/15685381-00002838.
- Sarashina M, Yoshida T. 2021. Diet composition of the invasive American bullfrog (*Lithobates catesbeianus*) in Onuma Quasi-National Park, Hokkaido, Japan. *Current Herpetology* 40(1):77–82 DOI 10.5358/hsj.40.77.
- Sato R, Nishihara S. 2017. Impacts of invasive bullfrogs and the control strategy. In: Takahashi K, ed. *Recovering fish*. Tokyo: Kouseisha, 68–80 [in Japanese].

- **Sato T. 2016.** Reptiles and amphibians distributed on the periphery of an army cemetery in Nara City. *Annual Bulletin of Oyasato Institute for the Study of Religion, Tenri University* **22**:49–74 [in Japanese with English summary].
- Silva ET, Reis EP, Feio RN, Filho OPR. 2009. Diet of the invasive frog Lithobates catesbeianus (Shaw, 1802) (Anura: Ranidae) in Viçosa, Minas Gerais state, Brazil. South American Journal of Herpetology 4(3):286–294 DOI 10.2994/057.004.0312.
- Stewart MM, Sandison P. 1972. Comparative food habits of sympatric mink frogs, bullfrogs, and green frogs. *Journal of Herpetology* 3(4)):241–244 DOI 10.2307/1562781.
- Strauss SY, Lau JA, Carroll SP. 2006. Evolutionary responses of natives to introduced species: what do introductions tell us about natural communities? *Ecology Letters* **9(3)**:357–374 DOI 10.1111/j.1461-0248.2005.00874.x.
- Sugiura S. 2016. Impacts of introduced species on the biota of an oceanic archipelago: the relative importance of competitive and trophic interactions. *Ecological Research* 31(2):155–164 DOI 10.1007/s11284-016-1336-0.
- **Sugiura S. 2018.** Anti-predator defences of a bombardier beetle: is bombing essential for successful escape from frogs? *PeerJ* 6:e5942 DOI 10.7717/peerj.5942.
- Sugiura S. 2020a. Predators as drivers of insect defenses. *Entomological Science* 23(3):316–337 DOI 10.1111/ens.12423.
- Sugiura S. 2020b. Active escape of prey from predator vent via the digestive tract. *Current Biology* 30(15):R867–R868 DOI 10.1016/j.cub.2020.06.026.
- Sugiura S. 2021. Beetle bombing always deters praying mantises. *PeerJ* 9(3888):e11657 DOI 10.7717/peerj.11657.
- Sugiura S, Sato T. 2018. Successful escape of bombardier beetles from predator digestive systems. *Biology Letters* 14(2):20170647 DOI 10.1098/rsbl.2017.0647.
- Tawa K, Sagawa S. 2017. Breeding habitats of frogs in paddy field and fallow field biotopes in Shounji (Toyooka City, Hyogo Prefecture). *Yaseihukki* 5:29–38 [in Japanese with English Abstract].
- Vermeij GJ. 1982. Phenotypic evolution in a poorly dispersing snail after arrival of a predator. *Nature* 299(5881):349–350 DOI 10.1038/299349a0.
- Vrcibradic D, Diaz A, Cosendey BN, Nascimento BB, Borges-Júnior VNT. 2017. Trichodactylus dentatus (Crustacea, Decapoda, Trichodactylidae) and other prey of a large adult of the exotic American bullfrog, Lithobates catesbeianus (Ranidae), caught in a disturbed habitat in southeastern Brazil. Herpetology Notes 10:375–378.
- Werner EE, Wellborn GA, McPeek MA. 1995. Diet composition in postmetamorphic bullfrogs and green frogs: implications for interspecific predation and competition. *Journal of Herpetology* 29(4):600–607 DOI 10.2307/1564744.
- Wu Z, Li Y, Wang Y, Adams MJ. 2005. Diet of introduced bullfrogs (*Rana catesbeiana*): predation on and diet overlap with native frogs on Daishan Island, China. *Journal of Herpetology* 39(4):668–674 DOI 10.1670/78-05N.1.
- Yahiro K, Fujimoto T, Tokuda M, Yano K. 1992. Species composition and seasonal abundance of ground beetles (Coleoptera) in paddy fields. *Japanese Journal of Entomology* 60(4):805–813.