



# Factors Related to Mental Health of Foreign Care Workers in Long-Term Care Facilities in Japan during the COVID-19 Pandemic—A Comparative Study

Wu, Qian

Yamaguchi, Yuko

Greiner, Chieko

---

## (Citation)

International Journal of Environmental Research and Public Health, 19(24):16491

## (Issue Date)

2022-12

## (Resource Type)

journal article

## (Version)

Version of Record

## (Rights)

© 2022 by the authors. Licensee MDPI, Basel, Switzerland.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## (URL)

<https://hdl.handle.net/20.500.14094/0100477982>





Article

# Factors Related to Mental Health of Foreign Care Workers in Long-Term Care Facilities in Japan during the COVID-19 Pandemic—A Comparative Study

Qian Wu, Yuko Yamaguchi  and Chieko Greiner \*

Graduate School of Health Sciences, Kobe University, 7-10-2 Tomogaoka, Suma, Kobe 6540142, Hyogo, Japan

\* Correspondence: greiner@harbor.kobe-u.ac.jp

**Abstract:** This study aimed to evaluate the factors related to the mental health of foreign care workers in Japan's long-term care (LTC) facilities and compare their results with those of native care workers. We conducted a cross-sectional survey covering 80 LTC facilities across Japan between August and November 2021. The survey mainly included mental health, workload, reward, sense of coherence, loneliness, COVID-19-specific factors and sociodemographic variables. The results show that workload was a distinct feature associated with the mental health of foreign care workers ( $n = 172$ ) when compared with those of native care workers ( $n = 154$ ). In addition, we found that the relationship between cultural adaptation and mental health in a sample of foreign care workers was mediated by loneliness and sense of coherence (SOC). Finally, we found that reward, loneliness, SOC, and COVID-19-specific factors had significant impacts on the mental health of both foreign and native care workers. These findings highlight the importance of support measures from the workplace for foreign care workers. Workplace interventions that focus on workload, reward, and sense of coherence strategies are required to address mental health improvement and may still be of value in dealing with the continuing COVID-19 pandemic.

**Keywords:** mental health; foreign care workers; workload; sense of coherence



**Citation:** Wu, Q.; Yamaguchi, Y.; Greiner, C. Factors Related to Mental Health of Foreign Care Workers in Long-Term Care Facilities in Japan during the COVID-19 Pandemic—A Comparative Study. *Int. J. Environ. Res. Public Health* **2022**, *19*, 16491. <https://doi.org/10.3390/ijerph192416491>

Academic Editor: Stefano Porru

Received: 8 November 2022

Accepted: 7 December 2022

Published: 8 December 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

By 2050, the number of people in OECD countries who are aged 80 years or older will rise from about 5%, as it currently stands, to almost 10% on average [1]. However, the growth of long-term care (LTC) services has been slower than the growth of the elderly population. To address this issue, many countries regard attracting and employing immigrants as one viable option. According to statistical data in OECD countries, foreign care workers make up almost 20% of the population who worked in LTC facilities [2]. Japan is no exception. As a country with one of the fastest-aging populations in the world, about 8% of the population was aged 80 years or older in 2015, and that number is expected to increase substantially by 2050 [3]. To compensate for the labor deficit in long-term care institutions, the Japanese government has recruited foreign care workers under economic partnership agreements (EPA) with Asian countries, starting with Indonesia and followed by the Philippines (in 2009) and Vietnam (in 2014) [4]. In 2017, the policy-related care work visa was issued to stimulate an increase in foreign participation in the LTC industry [5]. According to government statistics, 43,446 foreign workers were employed in long-term care facilities in 2021, an increase of 18.0 percent from the previous year [6]. As the number of foreign care workers is growing annually, they have already assumed a significant role in caring for the elderly.

However, the complexity of LTC environments and intensive workloads have a negative impact on the mental health of foreign care workers, manifesting as anxiety and depression, for example [7]. Studies on foreign care workers reported that they have poor

working conditions such as low wages, higher working hours, low job security and workplace abuse in host countries [8,9]. In addition, visa restrictions have been placed on foreign care workers. The majority of foreign care workers in Japan possess a visa that restricts their employment to the healthcare profession and limits their length of stay, hence increasing their stress levels [10,11]. In addition, governmental restrictions, such as the requirement for 2–5 years of experience working in Japan's long-term care institutions, require them to exert greater effort than locals in order to get the national nursing qualification [4]. In a previous study, responses from 22.5% of foreign care workers suggested that they were at risk of developing mental health problems [12]. Mental health problems were associated with the high rate of turnover and work performance [13,14]. To improve qualities of care and maintain well-being in the workplace, interventions regarding care workers' mental health should be implemented [15]. However, research thus far has primarily focused on the qualitative findings [7]; there has been little research on the mental health and related factors of foreign care workers serving the public.

The relevant factors of foreign care workers' mental health are complex and diverse, and mainly close to their migration environment and work conditions [15]. In most previous studies, the relationship between mental health and coping skills has been investigated as well [16–18]. Therefore, the following hypotheses were established.

### *1.1. Work Conditions and Mental Health*

There are several studies indicating the mental health of foreign workers associated with work conditions. Work-related stressors such as long working hours, high workload, and no free time were related to the mental health status among migrant workers in India [19]. Work overload stress of foreign care workers was the most common cause of mental health disorders [20]. According to a qualitative study, foreign care workers' dissatisfactions regarding the workplace were due to the workload, constraints with time, and poor peer relations [21]. These factors might become work-related stressors of migrant workers, which result in mental stress [22]. Moreover, greater work-related stress results from perceived low rewards, which was identified as a risk factor of mental health among LTC workers [23]. In a previous study, Benjamin et al. found a correlation between the mental health of workers and the degree to which they had autonomy in their jobs, as well as interactions with coworkers and supervisors [24]. However, the specific factors of work conditions that are connected with mental health among foreign care workers have not been studied in depth in comparison to native care workers.

**Hypothesis 1.** *Heavy workload is associated with negative mental health.*

**Hypothesis 2.** *There is a relationship between reward and positive mental health.*

### *1.2. Sense of Coherence and Mental Health*

There is empirical evidence of a strong relationship between mental health and sense of coherence (SOC), such that individuals with a strong SOC report a better overall health status [25]. It is a concept that reflects the ability to cope with stress and is at the core of the autogenesis theory [26]. In light of the current pandemic crisis, stress and challenges have been evident. In order to cope with them, SOC is a crucial aspect for mental health in care professionals who are exposed to stressful settings [27]. A solid SOC emerged as the strongest predictor for less severe symptoms of anxiety and depression among healthcare workers in Germany [28]. In Japan, it was identified that poor mental health was related to weak SOC among healthcare workers amid the COVID-19 pandemic [29]. It is extremely important that an attempt is made to understand the connection between the mental health and SOC of foreign care workers because so little research has been carried out on this topic.

**Hypothesis 3.** *Sense of coherence is associated with positive mental health.*

### 1.3. Loneliness and Mental Health

Immigrants' social factors have been identified as important contributors to anxiety and depression symptoms [30]. Importantly, with a migrant background, most foreign care workers experienced a sense of loneliness [31]. Previous research also indicates that loneliness negatively affects the health and well-being of immigrants [32]. Additionally, the COVID-19 pandemic introduced global experiences of social isolation. Because of the great distance between Japan and their home countries, it is especially possible for foreign care workers to experience negative effects on their mental health while working in Japan.

**Hypothesis 4.** *Loneliness was related to negative mental health.*

### 1.4. COVID-19-Specific Factors and Mental Health

On March 2020, the World Health Organization (WHO) declared the rapid worldwide spread of coronavirus disease 2019 (COVID-19) to be a pandemic. Since the onset of the ongoing pandemic, numerous cluster outbreaks of COVID-19 have been reported in long-term care (LTC) facilities worldwide, affecting both the residents and the care staff [33]. In the Japan Geriatrics Society, LTC facilities are particularly vulnerable places with the majority of residents at high risk of complications. To date, it was estimated that 1,600 LTC facilities reported cluster infection cases due to the emergence of COVID-19 [34]. Caring for the elderly and for persons with physical and mental disabilities or pre-existing medical conditions, LTC workers are the first-line defense in protecting those most vulnerable to infections and disease. Their health and safety are directly related to the benefits of residents and facilities [35]. During the COVID-19 pandemic, the immigrant population in Norway reported high levels of depression and anxiety [36]. The COVID-19 outbreak could be considered an uncontrollable stressful life event that can contribute to the development of or increase in mental health problems in foreign care workers. There is little evidence about how COVID-19 has impacted foreign care workers in LTC facilities.

**Hypothesis 5.** *COVID-19 specific factors are associated with mental health.*

### 1.5. Cultural Adaptation and Mental Health

Cultural adaptation is the process by which immigrants decide which parts of their culture from their home country to keep and which ones to change to fit their new surroundings [37–39]. These decisions might cause acculturative stress among immigrants. Studies in the U.S. have concluded that the mental health of immigrants was highly influenced by acculturative stress [40]. Research on foreign care workers also identified that cultural adaptation was an important predictor of mental health status [41,42]. Notably, according to the results of a survey regarding the problems foreign care workers face in Japan, they are experiencing difficulty with administrative procedures, real estate contracts, and automobile sales, as well as other procedures [43]. In addition, changes in temperature, the cost of living, culture, and language may make adaptation difficult for foreign workers who live abroad [44].

**Hypothesis 6.** *Cultural adaptation is related to the mental health.*

At present, there is a worldwide increase in interest in foreign care workers' difficulties and mental health. Based on the literature review, limited studies are available on comprehensive mental health support for foreign care workers. Additionally, little is known about the distinct characteristics of foreign care workers in long-term care facilities in comparison with native workers. To fill this gap, the goal of this study was to examine the factors that affect the mental health of foreign care workers in Japan's LTC facilities and compare their results with those of native care workers.

## 2. Materials and Methods

### 2.1. Study Design

This quantitative study used a cross-sectional design. Data were collected using online questionnaires (Microsoft Forms). We followed the reporting guidelines of the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement for observational studies (Supplementary File S1).

### 2.2. Setting and Data Collection

Data collection took place between August and November 2021. The target population included foreign care workers and native care workers in long-term care facilities in Japan. We randomly selected 80 care facilities to distribute the study instructions as well as the questionnaire invitations. Care facilities which accepted EPA program were chosen from the latest open data on the Ministry of Health, Labour and Welfare website. The study was performed using an online survey addressed to care workers. The link to the questionnaire survey along with the introductory letter was sent to these facilities.

The participant selection criteria included: (1) working at the current facility for more than one month, (2) providing direct nursing care, except in cases where the participant was the facility head and (3) ability to read Japanese or English.

We developed separate questionnaires for foreign care workers and Japanese care workers, including demographic characteristics, work-related information (reward and workload), mental health and other variables. English and Japanese versions were available for foreign care workers and added extra items including cultural adaptation and migration-related characteristics.

### 2.3. Sample Size

The sample size was determined by using the G\*Power software (v3.1.9.2, Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany) [45,46]. We used 2-sided testing, odds ratio = 2,  $\Pr(Y = 1 | X = 1) / \Pr(Y = 1 | X = 0) = 2$ ,  $H_0 = 0.5$ ,  $\alpha$  err prob = 0.05, power ( $1 - \beta$  err prob) = 0.85,  $R^2$  other  $X = 0.6$ . The minimum sample sizes for foreign and native care workers were 167 and 120, respectively.

### 2.4. Measurements

#### 2.4.1. Mental Health

Mental health was assessed using the Kessler 6 scale. This scale measures non-specific psychological distress and is utilized as a screening tool for serious mental illness in community-based samples [47]. The original version of the K6 was developed in English and then translated into Japanese [47]. The reliability and validity of the K6 have been systematically confirmed [47,48]. The Kessler 6 scale uses a 5-point Likert scale (ranging from 0 = none of the time to 4 = all of the time) to assess how often the respondent felt (a) nervous, (b) hopeless, (c) restless or fidgety, (d) so depressed that nothing could cheer them up, (e) that everything was a great effort, and (f) worthless over the past 30 days. Scale items were summed to achieve a score of up to 24, with higher scores on the Kessler 6 scale indicating a worse state of mental health [47,48].

#### 2.4.2. Workload

Workload was assessed using the short version of the effort–reward imbalance questionnaire. Workload was assessed by the effort sub-scale, which determines time pressure, interrupted work, and increased workload, with a total range from 3 to 12 [49,50]. Every item was scored on a 4-point Likert scale ranging from 1 (strongly disagree) to 4 (strongly agree). The validity and reliability of the Japanese version and English version were confirmed, with Cronbach's  $\alpha$  coefficient 0.74–0.89 [50,51].

#### 2.4.3. Reward

Reward was assessed using the short version of the effort–reward imbalance questionnaire, which includes a seven-item reward subscale [48,49]. Items were scored using a four-point Likert scale ranging from 1 (strongly disagree) to 4 (strongly agree), with higher scores indicating a positive perception of reward, including mutual respect and promotion. The validity and reliability of the Japanese version and English version were confirmed, with Cronbach's  $\alpha$  coefficient 0.79–0.89 [50,51]. Accordingly, this sample had a Cronbach's  $\alpha$  coefficient of 0.79. Subscale scores were determined by summing the seven items.

#### 2.4.4. Sense of Coherence (SOC)

Sense of coherence was measured by the Sense of Coherence (SOC-3) scale [52]. It measures SOC based on three items. It consists of three questions concerning manageability, meaningfulness and comprehensibility—factors representing one's internal resources when coping with difficulties in life [52,53]. The questions were: “Do you usually see a solution to problems and difficulties that other people find hopeless?” (manageability), “Do you usually feel that your daily life is a source of personal satisfaction?” (meaningfulness), “Do you usually feel that the things that happen to you in your daily life are hard to understand?” (comprehensibility) [52,53]. The sum score of the SOC-3 ranged from 1 to 7, and higher values indicated a higher sense of coherence [53]. The Japanese version and English version of the SOC scale have been shown to have adequate reliability and validity [53,54]. In this study, the Cronbach's  $\alpha$  coefficient was 0.73.

#### 2.4.5. Loneliness

Loneliness was assessed using the UCLA Loneliness Scale (UCLA) [55]. The UCLA Loneliness Scale has shown satisfactory reliability (Cronbach's  $\alpha$  = 0.84) [56] and both concurrent and discriminant validity [57]. This scale contains 10 items that assess how frequently a participant has felt certain emotions, with values ranging from 1 = never, 2 = rarely, 3 = sometimes, to 4 = often [58]. The scale consists of 5 positively and 5 negatively scored items with a total score of 10–40 points. Higher scores indicate greater degrees of loneliness and there is no identified cut-off score that defines loneliness [55,58]. In Japan, the reliability and validity of the original version have been evaluated [58]. In this study, the Cronbach's  $\alpha$  coefficient was 0.88.

#### 2.4.6. COVID-19-Specific Factors

COVID-19-specific factors included income changes during the pandemic, COVID-19 clusters in the workplace, periodic PCR tests, whether they had been infected with COVID-19, and whether they had cared for elderly individuals with COVID-19.

#### 2.4.7. Cultural Adaptation

The cultural adaptation items were created by us, based on the Sociocultural Adaptation Scale [59]. Based on a survey on foreign care workers, from the 40-item scale, we chose 5 general-culture-relevant items [59]. These items were making friends, becoming used to Japanese food, disaster preparedness, house hunting, and valuing freedom. Participants were asked to indicate the extent to which each statement was true for them on a 5-point Likert scale (from 1 “No difficulty” to 5 “Extreme difficulty”).

### 2.5. Statistical Analysis

This study included both descriptive and inferential statistics. A Kolmogorov–Smirnov test was used to examine normality. The continuous variables with normal distribution were represented by the mean and standard deviation (SD), and the continuous variables with non-normal distribution were represented by the median. The distribution of categorical variables was expressed as a frequency and percentage. The distribution difference in the demographic characteristics, as well as other variables, between foreign care workers and native care workers were evaluated by a t-test, a Mann–Whitney U test, a chi-squared test



or Fisher's exact test. Correlations, alpha reliabilities ( $\alpha$ ) for each scale, and descriptive statistics were performed with R software.

Data were further analyzed using Partial Least Squares Structural Equation Modeling (PLS-SEM). SmartPLS v3.0 software was used, selecting 5000 samples for the bootstrapping procedure [60]. PLS-SEM analyses follow a two-step approach given by Hair et al. [61]. PLS-SEM is a non-parametric technique that takes advantage of the variance that can be explained in latent unobserved dimensions. Smart PLS-SEM requires less information about residual distributions, measurement scales, and sample sizes compared with the covariance-based SEM [62]. In PLS-SEM, relations between latent and manifest variables, as well as between latent variables (structural/inner model), are defined in the form of path models. The directional interpreted paths are represented by connecting arrows.

### 2.6. Ethical Considerations

This study was conducted upon receiving the approval of the Ethics Review Committee of the Kobe University department of Health Sciences, Japan (No.1014). After obtaining approval from all the ethics committees involved, we contacted the institutions which provided us with formal authorization. All the participants who met the inclusion criteria were invited to participate in the study. A complete description of this survey and informed consent forms were sent to participants prior to the questionnaires. After the participants selected "Yes," data collection was able to continue. The participants responded to the questionnaires anonymously.

## 3. Results

In total, we recruited 335 care workers, including certified care workers and care assistants. Overall, 5 of those participating through web-based entries declined to participate. After data cleaning, we found that 4 entries were missing substantial data. We excluded these entries from further analysis, leaving 326 valid responses (foreign care workers = 172; native care workers = 154). Regarding the nationality of foreign care workers, most of them were Vietnamese ( $n = 54$ , 31.40%), followed by Indonesian ( $n = 50$ , 29.07%), and the remaining were of other nationalities ( $n = 68$ , 39.53%) (Supplementary File S2). Table 1 shows the comparisons of the demographic characteristics between the foreign care workers and native care workers. The total participants' average age was  $35.05 \pm 9.35$  years old. However, there was an age difference between foreign and native care workers. The mean age of foreign care workers was  $29.06 \pm 5.25$  years old. Additionally, the mean age of native care workers was  $41.73 \pm 8.35$  years old. Moreover, the number of years of care work experience in native care workers was  $10.65 \pm 5.72$ , which was much longer than foreign care workers. There were also significant differences in professional status between the two groups. Overall, 62.21% foreign care workers did not yet have the necessary national qualifications. On the contrary, 66.23% native care workers had obtained their national qualifications.

As shown in Table 2, no significant differences were observed between the foreign care workers and the native care workers regarding mental health. Foreign care workers had a higher level of SOC than native care workers. Regarding workload and reward, the workload of native care workers was greater than that of foreign care workers, as was the reward level. A drop in income due to COVID-19 was evident among native care workers. Other COVID-19-specific factors show no difference between the two groups.

The results of hypothesis testing among foreign care workers and native care workers are summarized in Table 3. After controlling for demographic factors (age and professional status), we found that the impact of COVID-19 was the strongest factor that positively affected mental health among both foreign care workers ( $\beta = 0.325$ ,  $p < 0.001$ ) and native care workers ( $\beta = 0.316$ ,  $p < 0.001$ ). It was also shown that loneliness has a significant effect on the mental health of both foreign care workers ( $\beta = 0.246$ ,  $p < 0.001$ ) and native care workers ( $\beta = 0.220$ ,  $p = 0.001$ ). Meanwhile, mental health was significantly associated with reward (FCW:  $\beta = -0.112$ ; NCW:  $\beta = -0.195$ ) and SOC (FCW:  $\beta = -0.232$ ; NCW:

$\beta = -0.293$ ). Workload only had a significant impact on the mental health of foreign care workers ( $\beta = 0.198$ ,  $p < 0.001$ ).

**Table 1.** Demographic characteristics of care workers (n = 326).

Characteristics	All	Foreign Care Workers	Native Care Workers	p-Value
	n = 326	n = 172	n = 154	
	%/M $\pm$ SD	%/M $\pm$ SD	%/M $\pm$ SD	
Age	35.05 $\pm$ 9.35	29.06 $\pm$ 5.25	41.73 $\pm$ 8.35	<0.001 <sup>a</sup>
Year of Care work experience	6.92 $\pm$ 5.61	3.58 $\pm$ 2.61	10.65 $\pm$ 5.72	<0.001 <sup>a</sup>
Gender				0.648 <sup>b</sup>
Female	220 (67.48)	118 (68.60)	102 (66.23)	
Male	106 (32.51)	54 (31.40)	52 (33.77)	
Marriage				<0.001 <sup>b</sup>
Unmarried	181 (55.52)	115 (66.86)	66 (42.86)	
Married	145 (44.48)	57 (33.14)	88 (57.14)	
Professional status				<0.001 <sup>b</sup>
Certified care worker	167 (51.23)	65 (37.79)	102 (66.23)	
Care assistant	159 (48.77)	107 (62.21)	52 (33.77)	
Workplace				0.498 <sup>b</sup>
Special elderly nursing home	165 (50.61)	84 (48.84)	81 (52.60)	
Long-term care health facility	161 (49.39)	88 (51.16)	73 (47.40)	
Educational background				<0.001 <sup>c</sup>
High school or technical school	36 (11.04)	2 (1.16)	34 (22.07)	
Junior college	173 (47.79)	94 (54.65)	79 (51.30)	
Bachelor degree	114 (34.97)	73 (42.44)	41 (26.62)	
Master degree	3 (0.92)	3 (1.74)	0 (00.00)	
Night shift				0.487 <sup>b</sup>
Yes	216 (66.26)	111 (64.53)	105 (68.18)	
No	110 (33.74)	61 (35.47)	49 (31.82)	

Notes: M = mean, SD = standard deviation; <sup>a</sup>: Mann–Whitney U test; <sup>b</sup>: Chi-square test; <sup>c</sup>: Fisher’s exact test.

**Table 2.** Average scores of variables in foreign care workers and native care workers.

Variable	Foreign Care Workers	Native Care Workers	p-Value
	n = 172	n = 154	
	M $\pm$ SD	M $\pm$ SD	
Mental health	8.18 $\pm$ 3.87	8.70 $\pm$ 4.22	0.245 <sup>d</sup>
SOC	15.03 $\pm$ 2.82	13.91 $\pm$ 3.37	0.001 <sup>a</sup>
Workload	8.35 $\pm$ 1.93	8.90 $\pm$ 2.16	0.014 <sup>a</sup>
Reward	19.20 $\pm$ 3.88	20.76 $\pm$ 3.67	<0.001 <sup>d</sup>
Loneliness	22.45 $\pm$ 4.89	22.95 $\pm$ 4.86	0.346 <sup>a</sup>
COVID-19-specific factors			0.464 <sup>b</sup>
Periodic PCR test in workplace			
Yes	148 (86.05)	128 (83.12)	
No	24 (13.95)	26 (16.88)	
COVID-19 outbreak in workplace			0.793 <sup>b</sup>
Yes	35 (20.35)	37 (24.02)	
No	137 (79.65)	117 (75.98)	
Income changes during COVID-19			<0.001 <sup>c</sup>
Increased a lot	1 (0.58)	1 (0.65)	
Increased	4 (2.32)	7 (4.55)	
Increased a little	36 (20.93)	15 (9.74)	
No change	113 (65.70)	92 (59.74)	
Decreased a little	11 (6.40)	33 (21.43)	
Decreased	4 (2.33)	6 (3.90)	
Decreased a lot	3 (1.74)	0 (0.00)	
Infected with COVID-19			0.225 <sup>c</sup>



Table 2. Cont.

Variable	Foreign Care Workers	Native Care Workers	<i>p</i> -Value
	<i>n</i> = 172	<i>n</i> = 154	
	<i>M</i> ± <i>SD</i>	<i>M</i> ± <i>SD</i>	
Yes	9 (5.23)	4 (2.60)	0.057 <sup>b</sup>
No	155 (90.12)	142 (92.21)	
Unsure	8 (4.65)	8 (5.19)	
Cared for elderly with COVID-19			
Yes	25 (14.53)	35 (22.73)	
No	147 (85.47)	119 (77.27)	

Notes: *M* = mean, *SD* = standard deviation; SOC = sense of coherence; <sup>a</sup>: Mann–Whitney U test; <sup>b</sup>: Chi-square test; <sup>c</sup>: Fisher’s exact test; <sup>d</sup>: Student’s *t*-test.

Table 3. Factors associated with mental health among foreign care workers and native care workers.

Variable	Foreign Care Workers			Native Care Workers		
	n = 172			n = 154		
	Beta(β)	T-Value	<i>p</i> Values	Beta(β)	T-Value	<i>p</i> Values
SOC	−0.232	3.613	<0.001	−0.293	4.009	<0.001
Workload	0.198	3.739	<0.001	0.050	0.751	0.453
Reward	−0.112	2.120	0.035	−0.195	2.322	0.021
Loneliness	0.246	3.846	<0.001	0.220	3.297	0.001
COVID-19	0.325	6.263	<0.001	0.316	5.669	<0.001
Adjusted R <sup>2</sup> (Mental health) = 0.617			Adjusted R <sup>2</sup> (Mental health) = 0.575			

Notes: SOC = sense of coherence; COVID-19 = COVID-19-specific factors.

The PLS analysis for foreign care workers is shown in Table 4. The relationship between workload and mental health ( $\beta = 0.247$ ,  $p < 0.001$ ) is statistically significant, in line with the results shown in Table 3. Additionally, reward was found to have a negative effect on mental health ( $\beta = -0.114$ ,  $p = 0.023$ ). Conversely, cultural adaptation had no direct effect on the mental health of foreign care workers ( $\beta = 0.038$ ,  $p = 0.460$ ). Finally, by looking at the internal variance inflation factor (VIF) index to determine the multicollinearity between latent variables, we can see that all values were lower than the standard value of 5.00 (inner VIF: 1.200–1.640). Therefore, multicollinearity was not confirmed [63]. As a result, Hypothesis 6 was rejected, and Hypotheses 1, 2, 3, 4, and 5 were accepted.

Table 4. Hypothesis testing of mental health model in foreign care workers (*n* = 172).

Hypotheses	Beta(β)	T-Value	95%CI	f <sup>2</sup>	<i>p</i> Values	VIF
H1 Workload → Mental health	0.247	4.683	[0.147,0.347]	0.132	<0.001	1.170
H2 Reward → Mental health	−0.114	2.139	[−0.228,0.001]	0.023	0.030	1.410
H3 SOC → Mental health	−0.239	3.828	[−0.366, −0.111]	0.097	<0.001	1.490
H4 Loneliness → Mental health	0.207	3.194	[0.081,0.334]	0.066	<0.001	1.640
H5 COVID-19 → Mental health	0.327	6.822	[0.235,0.419]	0.211	<0.001	1.280
H6 CA → Mental health	0.038	0.783	[−0.057,0.134]	0.003	0.460	1.200

Notes: SOC = sense of coherence; COVID-19 = COVID-19-specific factors; CA: Cultural Adaptation.

In this study, there were two indirect effects, which are given in Table 5. The first indirect effect concerns the mediating role of loneliness in the relationship between mental health and cultural adaptation. This hypothesis is supported, according to the following results:  $t = 2.890$ ,  $p = 0.004$ . The second indirect effect concerns the mediating role of SOC in the relationship between mental health and cultural adaptation. This hypothesis is supported, according to the following results:  $t = 2.545$  and  $p = 0.011$ .

**Table 5.** Direct and indirect effects predicting mental health of foreign care workers (n = 172).

Path	Beta	T-Value	95%CI	p Values
Workload → Mental health	0.241	4.788	[0.131, 0.330]	<0.001
Reward → Mental health	−0.117	2.187	[−0.245, −0.035]	0.029
SOC → Mental health	−0.246	3.987	[−0.368, −0.119]	<0.001
Loneliness → Mental health	0.213	3.491	[0.076, 0.323]	0.001
COVID-19 → Mental health	0.329	6.538	[0.228, 0.424]	<0.001
Cultural Adaptation → Loneliness	0.362	6.094	[0.269, 0.493]	<0.001
Cultural Adaptation → SOC	−0.286	4.332	[−0.436, −0.178]	<0.001
Cultural Adaptation → Loneliness → Mental Health	0.077	2.890	[0.024, 0.125]	0.004
Cultural Adaptation → SOC → Mental Health	0.071	2.545	[0.023, 0.127]	0.011
Adjusted R <sup>2</sup> (Mental health) = 0.591				

Notes: SOC = sense of coherence; COVID-19 = COVID-19-specific factors.

#### 4. Discussion

The present study investigated mental health and related factors in foreign care workers and compared these factors with those of native care workers in Japan. To the best of our knowledge, this is the first study to compare the mental health outcomes of foreign care workers and native care workers in long-term care (LTC) facilities. This study produced three key findings. First, compared with native care workers, we determined that workload was a distinct feature connected with the mental health of foreign care workers. Second, we found that the relationship between cultural adaptation and mental health in a sample of foreign care workers is mediated by loneliness and SOC. Thirdly, COVID-19-specific factors had a strong impact on the mental health of both foreign and native care workers. In addition, we have found that reward, loneliness, and SOC are significant factors associated with the mental health of both foreign and native care workers. In terms of cultural adaptation, there is no direct evidence linking the mental health of foreign care workers.

Regarding workload, we found that foreign care workers were prone to having a lower external workload than native care workers in LTC facilities in Japan. This finding may be explained by the difference in professional status between the two groups. A majority of foreign care workers are care assistants without a national qualification related to care work, whereas most native care workers are certified care workers. In Japan, most certified care workers have more than 3 years of care work experience and can undertake more specific age-related care, such as dementia care [64]. Care assistants, on the other hand, can only carry out daily life care for older individuals; for instance, they can help with bathing [65]. The age difference between local and foreign care workers may also play a role in the noticeable variation in workload. Consistent with prior research [66,67], there was a substantial age disparity between foreign care workers and local care workers in our study. Regarding physical workload, such as transferring a patient, younger care workers may feel less burdened [68]. In addition, owing to the status of foreign care workers, some leadership responsibilities, such as training new care employees, may not be assigned to them [7].

In the hypothesis testing, our findings demonstrated that only foreign care workers with a heavier workload are more likely to experience mental health issues. This was consistent with a study on the mental health of immigrant workers, which indicated that bad working conditions, particularly heavy workloads, were strongly related to mental illness [23]. Contrary to the findings of the care staff study [69], we did not find a relationship between the workload and mental health of native care workers. There is a rational explanation for this, in that immigrant workers' mental health declines as a result of the stress brought about by growing workloads and low levels of reward in host countries [70]. Work-related stress has long been recognized as a key risk factor for migrant workers' mental health [71]. An interesting finding from a study of interviews with foreign care workers in Japan is that completing paper care records and other documents in Japanese

is also seen as a heavy workload that can lead to burnout [7]. However, the focus of this study was not the types of workloads in the LTC sectors, but rather on the relationships between mental health and workload.

Another finding is that both foreign and native care workers with a positive perception of reward tend to possess better mental health. This finding ties well with previous studies [21,72]. This is because increasing the appropriateness of rewards can make people feel positive emotions that are good for their mental health and well-being [24]. To be more specific, occupational rewards can be broken down into three categories: satisfaction with earnings, esteem, and job security [50]. There is a tendency toward declining mental well-being after migration among foreign care workers living with lower economic conditions [10]. This is because most foreign workers emigrated to developed countries for a better economic life [12,73]. In comparison with the high cost of living in Japan, foreign care workers are typically dissatisfied with their low pay, which may have a negative impact on their mental health [7]. A new study has also identified a correlation between poor self-esteem at work and mental health issues [74]. Many care workers have an esteem need for clear acknowledgement of their sacrifices and extraordinary efforts [75]. Meanwhile, lack of job security has been highlighted as a significant mental health risk factor for care workers during the COVID-19 pandemic [76]. Fear of becoming infected may make care workers feel unsafe at work, which could worsen their mental stress [77].

Although we assumed that cultural adaptation has a direct effect on mental health based on previous reports showing that such cultural-related stress facilitates poor mental health among migrant care workers [41,42], our results did not support this assumption. On the other hand, we found that cultural adaptation could have an indirect effect on mental health through the mediating effects of loneliness and sense of coherence (SOC).

Regarding loneliness, it has been defined as an unpleasant experience caused by a person's assessment that their network of social relations is insufficient [78]. Our findings supported the psychological pathway of the conceptual model proposed by Berkman et al. [42], who argued that social relationships may influence mental health outcomes via multiple mechanisms. As one of the most significant post-migration stressors [30], loneliness was strongly associated with cultural adaptation among foreign care workers, and was further related to their mental health.

It was also found that the SOC completely mediated the effect of cultural adaptation and mental health. This was consistent with previous studies on immigrants [79]. Our result also identified a strong relationship between SOC and mental health status which is directly in line with previous findings [28,29]. By identifying the fact that SOC mediates the relation between cultural adaptation and mental health, more emphasis can be placed on utilizing interventions that strengthen SOC for the migration population with growing mental health disorders [26]. This may be particularly beneficial in terms of mental health policy, since it may suggest that governments and institutions offer multiple means of supports rather than relying solely on language and training in caregiving skills [80].

COVID-19-specific factors were found to be strongly associated with both foreign and native care workers' mental health, even after controlling for demographic features. Research on the mental health status of care workers has largely focused on the associations between workplace safety [81] and work-related changes by COVID-19 [82]. Many scholars found that the worst mental health among care workers was associated with caring for COVID-19-infected patients [83,84] and the fear of infection [85]. A Japanese longitudinal study also revealed that the effects of COVID-19 may have substantially reduced their income, thereby threatening their mental health [86]. Importantly, many long-term care facilities have experienced COVID-19 outbreaks [87,88], and were associated with some significant drop-in care workers, owing to elevated absences and departures [89]. Therefore, the mental health of care workers may be negatively impacted by the stress caused by insufficient manpower and increased workloads [90].

This study possesses several limitations. First, due to the nature of cross-sectional study data collection, causal relationships between factors and mental health cannot be

inferred. Second, the data collected were only from the early phase of the pandemic, which may not represent other periods of the pandemic in which circumstances might have changed. Thirdly, this study tested the mediating effects of loneliness and SOC on the relationship between cultural adaptation and mental health. However, several other factors, such as gender and social support, may act as moderators that may affect the tested relationship. Our survey was only available in Japanese and English. This limitation affected our participants because it was difficult for foreign care workers with less than N2 Japanese proficiency to participate in the Japanese version. In addition, there are a large number of Chinese and Vietnamese care workers in Japan. Since English is not their first language, they are ineligible to participate in our studies if they cannot comprehend English well. Therefore, it is strongly suggested that the scope of this study be broadened in the future by including multiple languages among the questionnaires.

## 5. Conclusions

These findings highlight the importance of support measures from the workplace for foreign care workers. Workplace interventions that focus on workload, reward, and sense of coherence strategies are required in order to improve workers' mental health, and may still be of value in dealing with the continuing COVID-19 pandemic.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijerph192416491/s1>, Supplementary File S1 and Supplementary File S2.

**Author Contributions:** Conceptualization, Q.W.; methodology, Q.W.; software, Q.W.; validation, Q.W., Y.Y. and C.G.; formal analysis, Q.W.; investigation, Q.W.; resources, Q.W.; data curation, Q.W.; writing—original draft preparation, Q.W.; writing—review and editing, Q.W. and C.G.; supervision, C.G.; project administration, C.G.; funding acquisition, Q.W. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by Kobe University Premium Program (189k012k).

**Institutional Review Board Statement:** Kobe University department of Health Sciences, Japan (No.1014).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** We thank all of the participating LTC facilities and the questionnaire respondents.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. OECD. *Health at a Glance 2021: OECD Indicators*; OECD Publishing: Paris, France, 2021. [CrossRef]
2. OECD. *Who Cares? Attracting and Retaining Care Workers for the Elderly, OECD Health Policy Studies*; OECD Publishing: Paris, France, 2020. [CrossRef]
3. Ministry of Health, Labour and Welfare, 2021. Status Report on Long-Term Care Insurance. Available online: <https://www.mhlw.go.jp/topics/kaigo/osirase/jigyos/20/index.html> (accessed on 12 August 2022).
4. Ministry of Health, Labour and Welfare: Committee for the Acceptance of Foreign Care Workers. Available online: [https://www.mhlw.go.jp/stf/shingi/other-syakai\\_225506.html](https://www.mhlw.go.jp/stf/shingi/other-syakai_225506.html) (accessed on 12 August 2022).
5. Immigration Services Agency of Japan, Publications. 2016. Available online: [https://www.moj.go.jp/isa/laws/h28\\_kaisei.html](https://www.moj.go.jp/isa/laws/h28_kaisei.html) (accessed on 12 August 2022).
6. Portal Site of Official Statistics of Japan, Browse Statistics, System of Social and Demographic Statistics. Available online: <https://www.e-stat.go.jp/en/stat-search?page=1&toukei=00200502> (accessed on 12 August 2022).
7. Asis, E.; Carandang, R.R. The plight of migrant care workers in Japan: A qualitative study of their stressors on caregiving. *J. Migr. Health* **2020**, 1–2, 100001. [CrossRef] [PubMed]
8. Hurtado, D.A.; Sabbath, E.L.; Ertel, K.A.; Buxton, O.; Berkman, L.F. Racial disparities in job strain among American and immigrant long-term care workers. *Int. Nurs. Rev.* **2011**, 59, 237–244. [CrossRef] [PubMed]
9. Ha, N.H.L.; Chong, M.S.; Choo, R.W.M.; Tam, W.J.; Yap, P.L.K. Caregiving burden in foreign domestic workers caring for frail older adults in Singapore. *Int. Psychogeriatr.* **2018**, 30, 1139–1147. [CrossRef] [PubMed]
10. Nugraha, S.; Ohara-Hirano, Y. The Mental Health Predictors of Indonesian EPA Nurses and Certified Care Worker Candidates in Japan. *Jpn. Psychol. Res.* **2015**, 58, 85–96. [CrossRef]

11. Hirano, Y.O.; Nugraha, S.; Shiozu, H.; Higashijima, M.; Rahardjo, T.B.W. Measuring attentiveness toward oral care needs: A comparative study of Indonesian care workers in Japan and Indonesia. *Hum. Resour. Health* **2021**, *19*, 71. [CrossRef]
12. Sato, F.; Hayakawa, K.; Kamide, K. Investigation of mental health in Indonesian health workers immigrating to Japan under the Economic Partnership Agreement. *Nurs. Health Sci.* **2016**, *18*, 342–349. [CrossRef]
13. Ki, J.; Choi-Kwon, S. Health problems, turnover intention, and actual turnover among shift work female nurses: Analyzing data from a prospective longitudinal study. *PLoS ONE* **2022**, *17*, e0270958. [CrossRef]
14. Dhaini, S.R.; Zúñiga, F.; Ausserhofer, D.; Simon, M.; Kunz, R.; De Geest, S.; Schwendimann, R. Are nursing home care workers' health and presenteeism associated with implicit rationing of care? A cross-sectional multi-site study. *Geriatr. Nurs.* **2016**, *38*, 33–38. [CrossRef]
15. Moll, S.E.; Patten, S.; Stuart, H.; MacDermid, J.C.; Kirsh, B. Beyond Silence: A Randomized, Parallel-Group Trial Exploring the Impact of Workplace Mental Health Literacy Training with Healthcare Employees. *Can. J. Psychiatry* **2018**, *63*, 826–833. [CrossRef]
16. Lee, M.-C.; Chen, C.-H.; Hsieh, P.-H.; Ling, C.-H.; Yang, C.-C.; Chang, Y.-C.; Yeh, L.-Y.; Hung, H.-C.; Yeh, T.-F. Psychological impact, coping behaviors, and traumatic stress among healthcare workers during COVID-19 in Taiwan: An early stage experience. *PLoS ONE* **2022**, *17*, e0276535. [CrossRef]
17. López-Martínez, C.; Serrano-Ortega, N.; Moreno-Cámara, S.; Del-Pino-Casado, R. Association between Sense of Coherence and Mental Health in Caregivers of Older Adults. *Int. J. Environ. Res. Public Health* **2019**, *16*, 3800. [CrossRef]
18. Ho, K.H.; Chiang, V. A meta-ethnography of the acculturation and socialization experiences of migrant care workers. *J. Adv. Nurs.* **2014**, *71*, 237–254. [CrossRef]
19. Regmi, P.R.; van Teijlingen, E.; Mahato, P.; Aryal, N.; Jadhav, N.; Simkhada, P.; Zahiruddin, Q.S.; Gaidhane, A. The Health of Nepali Migrants in India: A Qualitative Study of Lifestyles and Risks. *Int. J. Environ. Res. Public Health* **2019**, *16*, 3655. [CrossRef]
20. De Diego-Cordero, R.; Vega-Escañó, J.; Tarriño-Concejero, L.; García-Carpintero-Muñoz, M. The Occupational Health of Female Immigrant Caregivers: A Qualitative Approach. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7807. [CrossRef]
21. Goel, K.; Penman, J. Employment experiences of immigrant workers in aged care in regional South Australia. *Rural Remote Health* **2015**, *15*, 15–28. [CrossRef]
22. Capasso, R.; Zurlo, M.C.; Smith, A.P. Ethnicity, work-related stress and subjective reports of health by migrant workers: A multi-dimensional model. *Ethn. Health* **2016**, *23*, 174–193. [CrossRef]
23. Siegrist, J.; Lunau, T.; Wahrendorf, M.; Dragano, N. Depressive symptoms and psychosocial stress at work among older employees in three continents. *Glob. Health* **2012**, *8*, 27. [CrossRef]
24. Schilgen, B.; Nienhaus, A.; Mösko, M. The Extent of Psychosocial Distress among Immigrant and Non-Immigrant Homecare Nurses—A Comparative cross Sectional Survey. *Int. J. Environ. Res. Public Health* **2020**, *17*, 1635. [CrossRef]
25. Erim, Y.; Morawa, E.; Atay, H.; Aygün, S.; Gökalp, P.; Senf, W. Sense of coherence and depression in the framework of immigration: Turkish patients in Germany and in Turkey. *Int. Rev. Psychiatry* **2011**, *23*, 542–549. [CrossRef]
26. Antonovsky, A. The structure and properties of the sense of coherence scale. *Soc. Sci. Med.* **1993**, *36*, 725–733. [CrossRef]
27. Schäfer, S.K.; Sopp, M.R.; Schanz, C.G.; Staginnus, M.; Göritz, A.S.; Michael, T. Impact of COVID-19 on Public Mental Health and the Buffering Effect of a Sense of Coherence. *Psychother. Psychosom.* **2020**, *89*, 386–392. [CrossRef] [PubMed]
28. Schmuck, J.; Hiebel, N.; Rabe, M.; Schneider, J.; Erim, Y.; Morawa, E.; Jerg-Bretzke, L.; Beschoner, P.; Albus, C.; Hannemann, J.; et al. Sense of coherence, social support and religiosity as resources for medical personnel during the COVID-19 pandemic: A web-based survey among 4324 health care workers within the German Network University Medicine. *PLoS ONE* **2021**, *16*, e0255211. [CrossRef] [PubMed]
29. Tanaka, K.; Tahara, M.; Mashizume, Y.; Takahashi, K. Effects of Lifestyle Changes on the Mental Health of Healthcare Workers with Different Sense of Coherence Levels in the Era of COVID-19 Pandemic. *Int. J. Environ. Res. Public Health* **2021**, *18*, 2801. [CrossRef] [PubMed]
30. Chen, W.; Wu, S.; Ling, L.; Renzaho, A.M. Impacts of social integration and loneliness on mental health of humanitarian migrants in Australia: Evidence from a longitudinal study. *Aust. N. Z. J. Public Health* **2018**, *43*, 46–55. [CrossRef] [PubMed]
31. Hall, B.J.; Garabiles, M.R.; Latkin, C.A. Work life, relationship, and policy determinants of health and well-being among Filipino Domestic Workers in China: A qualitative study. *BMC Public Health* **2019**, *19*, 229. [CrossRef]
32. Toselli, S.; Gualdi-Russo, E.; Marzouk, D.; Sundquist, J.; Sundquist, K. Psychosocial health among immigrants in central and southern Europe. *Eur. J. Public Health* **2014**, *24*, 26–30. [CrossRef]
33. Dosa, D.; Jump, R.L.; LaPlante, K.; Gravenstein, S. Long-Term Care Facilities and the Coronavirus Epidemic: Practical Guidelines for a Population at Highest Risk. *J. Am. Med. Dir. Assoc.* **2020**, *21*, 569–571. [CrossRef]
34. Ministry of Health, Labour and Welfare. Available online: [https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/kenkou\\_iryuu/dengue\\_fever\\_qa\\_00001.html](https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/kenkou_iryuu/dengue_fever_qa_00001.html) (accessed on 9 June 2021).
35. UNI Global Union, The Impact of COVID-19 on Long-Term Care Workers in the US, UK, Canada, Ireland, and Australia. Available online: [https://www.uniglobalunion.org/sites/default/files/imce/the\\_impact\\_of\\_covid19\\_fin.pdf](https://www.uniglobalunion.org/sites/default/files/imce/the_impact_of_covid19_fin.pdf) (accessed on 9 June 2021).
36. Harris, S.M.; Sandal, G.M. COVID-19 and psychological distress in Norway: The role of trust in the healthcare system. *Scand. J. Public Health* **2020**, *49*, 96–103. [CrossRef]
37. Berry, J.W. Acculturation and adaptation: Health consequences of culture contact among circumpolar peoples. *Arct. Med Res.* **1990**, *49*, 142–150.



38. Caplan, S. Latinos, Acculturation, and Acculturative Stress: A Dimensional Concept Analysis. *Policy Politi-Nurs. Pr.* **2007**, *8*, 93–106. [\[CrossRef\]](#)
39. Schwartz, S.J.; Unger, J.B.; Zamboanga, B.L.; Szapocznik, J. Rethinking the concept of acculturation: Implications for theory and research. *Am. Psychol.* **2010**, *65*, 237–251. [\[CrossRef\]](#)
40. Hagen, D.; Goldmann, E.; Parikh, N.S.; Goodman, M.; Boden-Albala, B. Acculturation and Depressive Symptoms Among Dominicans in New York City. *J. Immigr. Minor. Health* **2021**, *24*, 1186–1195. [\[CrossRef\]](#)
41. Adebayo, B.; Nichols, P.; Albrecht, M.A.; Brijnath, B.; Heslop, K. Investigating the Impacts of Acculturation Stress on Migrant Care Workers in Australian Residential Aged Care Facilities. *J. Transcult. Nurs.* **2020**, *32*, 389–398. [\[CrossRef\]](#)
42. Bekteshi, V.; Kang, S.-W. Contextualizing acculturative stress among Latino immigrants in the United States: A systematic review. *Ethn. Health* **2018**, *25*, 897–914. [\[CrossRef\]](#)
43. The Japan Association of Training Institutions for Certified Care Workers. Foreign Certified Care Candidates' Questionnaire Survey Report. Available online: [http://kaiyokyo.net/news/04\\_report\\_01.pdf](http://kaiyokyo.net/news/04_report_01.pdf) (accessed on 11 August 2022).
44. Efendi, F.; Chen, C.-M.; Nursalam, N.; Indarwati, R.; Ulfiana, E. Lived experience of Indonesian nurses in Japan: A phenomenological study. *Jpn. J. Nurs. Sci.* **2016**, *13*, 284–293. [\[CrossRef\]](#)
45. Kang, H. Sample size determination and power analysis using the G\*Power software. *J. Educ. Evaluation Health Prof.* **2021**, *18*, 17. [\[CrossRef\]](#)
46. Levenson, R.L. Statistical Power Analysis: Implications for Researchers, Planners, and Practitioners in Gerontology. *Gerontologist* **1980**, *20*, 494–498. [\[CrossRef\]](#)
47. Kessler, R.C.; Andrews, G.; Colpe, L.J.; Hiripi, E.; Mroczek, D.K.; Normand, S.-L.; Walters, E.E.; Zaslavsky, A.M. Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychol. Med.* **2002**, *32*, 959–976. [\[CrossRef\]](#)
48. Furukawa, T.; Kessler, R.C.; Slade, T.; Andrews, G. The performance of the K6 and K10 screening scales for psychological distress in the Australian National Survey of Mental Health and Well-Being. *Psychol. Med.* **2003**, *33*, 357–362. [\[CrossRef\]](#)
49. Siegrist, J. Adverse health effects of high-effort/low-reward conditions. *J. Occup. Health Psychol.* **1996**, *1*, 27–41. [\[CrossRef\]](#)
50. Siegrist, J.; Wege, N.; Pühlhofer, F.; Wahrendorf, M. A short generic measure of work stress in the era of globalization: Effort–reward imbalance. *Int. Arch. Occup. Environ. Health* **2008**, *82*, 1005–1013. [\[CrossRef\]](#) [\[PubMed\]](#)
51. Tsutsumi, A.; Ishitake, T.; Peter, R.; Siegrist, J.; Matoba, T. The Japanese version of the Effort-Reward Imbalance Questionnaire: A study in dental technicians. *Work. Stress* **2001**, *15*, 86–96. [\[CrossRef\]](#)
52. Schmalbach, B.; Tibubos, A.N.; Zenger, M.; Hinz, A.; Brähler, E. Psychometric Evaluation and Norm Values of an Ultra-Short Version of the Sense of Coherence Scale “SOC-3”. *Psychother. Psychosom. Med. Psychol.* **2019**, *70*, 86–93. [\[CrossRef\]](#) [\[PubMed\]](#)
53. Togari, T.; Yamazaki, Y.; Nakayama, K.; Shimizu, J. Development of a short version of the sense of coherence scale for population survey. *J. Epidemiol. Community Health* **2007**, *61*, 921–922. [\[CrossRef\]](#) [\[PubMed\]](#)
54. Lundberg, O.; Peck, M.N. A simplified way of measuring sense of coherence. *Eur. J. Public Health* **1995**, *5*, 56–59. [\[CrossRef\]](#)
55. Russell, D. UCLA Loneliness Scale (Version 3): Reliability, Validity, and Factor Structure. *J. Pers. Assess.* **1996**, *66*, 20–40. [\[CrossRef\]](#)
56. Rico-Uribe, L.A.; Caballero, F.F.; Olaya, B.; Tobiasz-Adamczyk, B.; Koskinen, S.; Leonardi, M.; Haro, J.M.; Chatterji, S.; Ayuso-Mateos, J.L.; Miret, M. Loneliness, Social Networks, and Health: A Cross-Sectional Study in Three Countries. *PLoS ONE* **2016**, *11*, e0145264. [\[CrossRef\]](#)
57. Hughes, M.E.; Waite, L.J.; Hawkey, L.C.; Cacioppo, J.T. A Short Scale for Measuring Loneliness in Large Surveys: Results from Two Population-Based Studies. *Res. Aging* **2004**, *26*, 655–672. [\[CrossRef\]](#)
58. Arimoto, A.; Tadaka, E. Reliability and validity of Japanese versions of the UCLA loneliness scale version 3 for use among mothers with infants and toddlers: A cross-sectional study. *BMC Women's Health* **2019**, *19*, 105. [\[CrossRef\]](#)
59. Searle, W.; Ward, C. The prediction of psychological and sociocultural adjustment during cross-cultural transitions. *Int. J. Intercult. Relations* **1990**, *14*, 449–464. [\[CrossRef\]](#)
60. Ramayah, T.J.F.H.; Cheah, J.; Chuah, F.; Ting, H.; Memon, M.A. *Partial Least Squares Structural Equation Modeling (PLS-SEM) Using smartPLS 3.0: An updated guide and practical guide to statistical analysis*; Pearson Malaysia Sdn Bhd: Kuala Lumpur, Malaysia, 2018.
61. Hair, J.F.; Hult, G.T.M.; Ringle, C.M.; Sarstedt, M. *A Primer on Partial Least Squares Structural Equation Modeling (PLS-SEM)*, 2nd ed.; Sage Publications: Thousand Oaks, CA, USA, 2017.
62. Hair, J.F.; Risher, J.J.; Sarstedt, M.; Ringle, C.M. When to use and how to report the results of PLS-SEM. *Eur. Bus. Rev.* **2019**, *31*, 2–24. [\[CrossRef\]](#)
63. Kock, N. Common method bias in PLS-SEM: A full collinearity assessment approach. *Int. J. e-Collab.* **2015**, *11*, 1–10. [\[CrossRef\]](#)
64. Yamada, Y.; Sekiya, M. Licensing and Training Requirements for Direct Care Workers in Japan: What can the United States and Japan learn from each other? *J. Aging Soc. Policy* **2003**, *15*, 113–129. [\[CrossRef\]](#)
65. Yamamoto-Mitani, N.; Saito, Y.; Takaoka, M.; Takai, Y.; Igarashi, A. Nurses' and Care Workers' Perception of Care Quality in Japanese Long-Term Care Wards: A Qualitative Descriptive Study. *Glob. Qual. Nurs. Res.* **2018**, *5*, 2333393618812189. [\[CrossRef\]](#)
66. Kameyama, J.; Hashizume, Y.; Takamura, Y.; Nomura, S.; Gomi, T.; Yanagi, H. Work engagement, well-being, and intent to continue working based on educational support among foreign care workers in Japan. *Environ. Health Prev. Med.* **2022**, *27*, 4. [\[CrossRef\]](#)



67. Kurazono, K.; Ikeoka, H.; Hiroi, S.; Iwasaki, K.; Takeshima, T.; Akazawa, M. Estimation of the cost of influenza antiviral medication guidance or support provided by healthcare professionals: A questionnaire survey in Japan. *J. Med Econ.* **2021**, *25*, 38–50. [\[CrossRef\]](#)
68. Suliman, M. Prevalence of low back pain and associated factors among nurses in Jordan. *Nurs. Forum* **2018**, *53*, 425–431. [\[CrossRef\]](#)
69. Zhang, Y.; Punnett, L.; Mawn, B.; Gore, R. Working Conditions and Mental Health of Nursing Staff in Nursing Homes. *Issues Ment. Health Nurs.* **2016**, *37*, 485–492. [\[CrossRef\]](#)
70. Li, W.; Moriyama, M.; Cui, Y.; Kazawa, K.; Nakaya, T.; Susanto, T. Presenteeism among Chinese workers in Japan and its relationship with mental health and health-promoting lifestyles. *Ind. Health* **2020**, *58*, 35–45. [\[CrossRef\]](#)
71. Chambers, C.N.L.; Frampton, C.M.A. Burnout, stress and intentions to leave work in New Zealand psychiatrists; a mixed methods cross sectional study. *BMC Psychiatry* **2022**, *22*, 380. [\[CrossRef\]](#)
72. Pelissier, C.; Fontana, L.; Fort, E.; Vohito, M.; Sellier, B.; Perrier, C.; Glerant, V.; Couprie, F.; Agard, J.P.; Charbotel, B. Impaired mental well-being and psychosocial risk: A cross-sectional study in female nursing home direct staff. *BMJ Open* **2015**, *5*, e007190. [\[CrossRef\]](#) [\[PubMed\]](#)
73. Hill, N.S.; Dorow, S.; Barnetson, B.; Martinez, J.F.; Matsunaga-Turnbull, J. Occupational Health and Safety for Migrant Domestic Workers in Canada: Dimensions of (Im)mobility. *NEW SOLUTIONS J. Environ. Occup. Health Policy* **2019**, *29*, 397–421. [\[CrossRef\]](#) [\[PubMed\]](#)
74. Hovey, J.D.; Magaña, C.G. Exploring the Mental Health of Mexican Migrant Farm Workers in the Midwest: Psychosocial Predictors of Psychological Distress and Suggestions for Prevention and Treatment. *J. Psychol.* **2002**, *136*, 493–513. [\[CrossRef\]](#) [\[PubMed\]](#)
75. Poh, L.W.; Teo, I.; Tewani, K.; Tan, H.K.; Yoon, S.; Sung, S.C. Understanding the needs of health care workers in Singapore during the COVID-19 outbreak: A qualitative analysis. *Am. J. Infect. Control* **2022**, *50*, 1133–1139. [\[CrossRef\]](#)
76. Aughterson, H.; McKinlay, A.R.; Fancourt, D.; Burton, A. Psychosocial impact on frontline health and social care professionals in the UK during the COVID-19 pandemic: A qualitative interview study. *BMJ Open* **2021**, *11*, e047353. [\[CrossRef\]](#)
77. Nielsen, M.B.D.; Ekholm, O.; Møller, S.P.; Ersbøll, A.K.; Santini, Z.I.; Grønbaek, M.K.; Thygesen, L.C. Mental wellbeing among Danish employees during the COVID-19 pandemic: Results from a longitudinal study on the role of industry and working environment. *Eur. J. Public Health* **2022**, *32*, 871–876. [\[CrossRef\]](#)
78. de Jong-Gierveld, J. Developing and testing a model of loneliness. *J. Pers. Soc. Psychol.* **1987**, *53*, 119–128. [\[CrossRef\]](#)
79. Abu-Kaf, S.; Khalaf, E. Acculturative Stress among Arab Students in Israel: The Roles of Sense of Coherence and Coping Strategies. *Int. J. Environ. Res. Public Health* **2020**, *17*, 5106. [\[CrossRef\]](#)
80. Kaneko, K. The Inflow of Southeast Asian Healthcare Worker Candidates in Japan: Japanese Reactions to the Possibility of Cultural and Ethnic Diversity. *Cph. J. Asian Stud.* **2016**, *33*, 78–96. [\[CrossRef\]](#)
81. Lai, J.; Ma, S.; Wang, Y.; Cai, Z.; Hu, J.; Wei, N.; Wu, J.; Du, H.; Chen, T.; Li, R.; et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw. Open* **2020**, *3*, e203976. [\[CrossRef\]](#)
82. Shiota, N.; Ishimaru, T.; Okawara, M.; Fujino, Y.; Tabuchi, T. Association between work-related changes caused by the COVID-19 pandemic and severe psychological distress among Japanese workers. *Ind. Health* **2021**, *60*, 216–223. [\[CrossRef\]](#)
83. Seys, D.; De Decker, E.; Waelkens, H.; Claes, S.; Panella, M.; Danckaerts, M.; Vanhaecht, K. A Comparative Study Measuring the Difference of Healthcare Workers Reactions Among Those Involved in a Patient Safety Incident and Healthcare Professionals While Working During COVID-19. *J. Patient Saf.* **2022**, *18*, 717–721. [\[CrossRef\]](#)
84. Maideen, A.A.; Idris, D.R.; Lupat, A.; Chung, Y.F.; Haji-Badarudin, H.; Suhai, H.; Abdullah, H.N.; Omar, H.; Kisut, R.; Rahman, H.A.; et al. Nurses' mental health and coping strategies throughout COVID -19 outbreak: A nationwide qualitative study. *Int. J. Ment. Health Nurs.* **2022**, *31*, 1213–1227. [\[CrossRef\]](#)
85. van Dijk, Y.; Janus, S.I.M.; de Boer, M.R.; Achterberg, W.P.; Roelen, C.A.M.; Zuidema, S.U. Job Demands, Work Functioning and Mental Health in Dutch Nursing Home Staff during the COVID-19 Outbreak: A Cross-Sectional Multilevel Study. *Int. J. Environ. Res. Public Health* **2022**, *19*, 4379. [\[CrossRef\]](#)
86. Kikuchi, H.; Machida, M.; Nakamura, I.; Saito, R.; Odagiri, Y.; Kojima, T.; Watanabe, H.; Fukui, K.; Inoue, S. Changes in Psychological Distress During the COVID-19 Pandemic in Japan: A Longitudinal Study. *J. Epidemiol.* **2020**, *30*, 522–528. [\[CrossRef\]](#)
87. Maeda, K. Outbreaks of COVID -19 infection in aged care facilities in Japan. *Geriatr. Gerontol. Int.* **2020**, *20*, 1241–1242. [\[CrossRef\]](#)
88. Gorges, R.J.; Konetzka, R.T. Factors Associated With Racial Differences in Deaths Among Nursing Home Residents With COVID-19 Infection in the US. *JAMA Netw. Open* **2021**, *4*, e2037431. [\[CrossRef\]](#)
89. Shen, K.; McGarry, B.E.; Grabowski, D.C.; Gruber, J.; Gandhi, A.D. Staffing Patterns in US Nursing Homes During COVID-19 Outbreaks. *JAMA Health Forum* **2022**, *3*, e222151. [\[CrossRef\]](#)
90. Bernotaite, L.; Malinauskiene, V. Workplace bullying and mental health among teachers in relation to psychosocial job characteristics and burnout. *Int. J. Occup. Med. Environ. Health* **2017**, *30*, 629–640. [\[CrossRef\]](#)