



Chlorogenic Acid and Caffeine in Coffee Restore Insulin Signaling in Pancreatic Beta Cells

Ihara, Yuka ; Asahara, Shunichirou ; Inoue, Hiroyuki ; Seike, Masako ; Ando, Misaki ; Kabutoya, Hiroki ; Kimura, Maki ; Kido, Yoshiaki

(Citation)

The Kobe journal of the medical sciences, 69(1):1-8

(Issue Date)

2023

(Resource Type)

departmental bulletin paper

(Version)

Version of Record

(JaLCD0I)

<https://doi.org/10.24546/0100480554>

(URL)

<https://hdl.handle.net/20.500.14094/0100480554>



Chlorogenic Acid and Caffeine in Coffee Restore Insulin Signaling in Pancreatic Beta Cells

YUKA IHARA¹, SHUN-ICHIRO ASAHARA^{2,*}, HIROYUKI INOUE¹,
MASAKO SEIKE², MISAKI ANDO¹, HIROKI KABUTOYA¹,
MAKI KIMURA-KOYANAGI², and YOSHIAKI KIDO¹

¹*Division of Medical Chemistry, Department of Metabolism and Diseases,
Kobe University Graduate School of Health Sciences, Kobe, Japan*

²*Division of Diabetes and Endocrinology, Department of Internal Medicine,
Kobe University Graduate School of Medicine, Kobe, Japan*

**Corresponding author*

Received 22 December 2022 / Accepted 6 January 2023

Keywords: Chlorogenic acid, Pancreatic β cell, Caffeine, Insulin signaling, Endoplasmic reticulum stress

The incidence of type 2 diabetes is reported to be lower in frequent coffee drinkers than in non-coffee drinkers. To elucidate the mechanism by which coffee prevents the onset of type 2 diabetes, we analyzed how caffeine and chlorogenic acid, which are components of coffee, alter insulin signaling in MIN6 cells, a mouse pancreatic β cell line. The results showed that caffeine improved insulin signaling under endoplasmic reticulum stress, and chlorogenic acid protected pancreatic β cells by enhancing the expression of insulin receptor substrate 2 via cAMP response element-binding protein and promoting insulin signaling downstream of insulin receptor substrate 2. In addition, chlorogenic acid was a potent antioxidant for the protection of pancreatic β cells. Furthermore, *in vivo* and *in vitro* analyses revealed that the pancreatic β cell-protective effect of chlorogenic acid was mediated by the alleviation of endoplasmic reticulum stress. The results suggest that these components of coffee have the potential to reduce the pathogenesis of type 2 diabetes and improve pancreatic β cell insufficiency.

In recent years, the number of patients with type 2 diabetes has been increasing worldwide (1). The main pathogenic mechanisms of type 2 diabetes can be divided into two categories: insulin resistance and insulin insufficiency. Furthermore, insulin insufficiency is thought to be caused by abnormalities in the secretory mechanism of pancreatic β cells and a decrease in the number of pancreatic β cells (2). Our laboratory has previously reported that decreased insulin signaling markedly reduces pancreatic β cell volume, resulting in impaired glucose tolerance (3,4). We have also shown that pancreatic β cell insufficiency is caused by autophagy disorders (5) and that the accumulation of cell cycle inhibitory factors reduces pancreatic β cell volume (6,7). If substances that alleviate these causative factors of pancreatic β cell insufficiency are identified, they are expected to make a significant contribution to the treatment and prevention of type 2 diabetes mellitus. Recently, novel diabetes medications such as incretin and sodium/glucose cotransporter 2 inhibitors have become widely used, but because drugs are associated with side effects, we are focusing on the therapeutic potential of nutrients contained in food. The advantage of nutrients is their established safety when administered over long periods of time.

Coffee is a beverage that has long been the subject of numerous health-related studies. It has been reported that long-term coffee consumption reduces the risk of ischemic heart disease (8) and that coffee intake has a protective effect on cerebral infarction (9). Several other studies have indicated that coffee prevents Alzheimer's disease, osteoporosis, and cancer (10,11,12).

Multiple studies have also reported an association between type 2 diabetes and coffee consumption (13,14,15). The mechanism by which these effects of coffee occur is still under investigation, but caffeine and chlorogenic acid, which are components of coffee, are thought to play an important role. Caffeine and chlorogenic acid are both nutrients whose effects on the body have been widely reported, and although their association with health status and disease has attracted attention, there are few reports on their effects on pancreatic β cells. In this study, we examined the effects of caffeine and chlorogenic acid on pancreatic β cells.

MATERIALS AND METHODS

Mice

Pancreatic β cell-specific C/EBP β TG mice on the C57BL/6J background were generated as previously described by inserting the C/EBP β gene under the control of the rat insulin promoter (16). The animals were

maintained in a 12-h light/dark cycle and fed normal chow from weaning at 3 weeks of age. All experiments were performed with male mice. After intraperitoneal injection of 10-week-old C/EBP β TG mice with 40 mg/kg/day chlorogenic acid or saline for 2 weeks, the pancreas was removed and islet morphology was observed. This study was approved by the Institutional Animal Care and Use Committee (Permission number: P190503) and carried out in accordance with the Kobe University Animal Experimentation Regulations.

Cell culture

MIN6 cells, a murine pancreatic β cell line, was routinely maintained in Dulbecco's modified Eagle's medium (Sigma-Aldrich) containing 15% heat-inactivated fetal calf serum at 37°C with 5% CO₂. The medium was replaced every 3–4 days. Chlorogenic acid loading of MIN6 cells was performed for 24 h at the concentrations indicated. Caffeine loading of MIN6 cells was performed for 48 h at the concentrations indicated.

Immunoblot analysis

Lysates of isolated islets and MIN6 cells were prepared as previously described (17) and probed with antibodies to β -actin (Sigma-Aldrich), phosphorylated (phospho)-Akt (Thr308), phospho-Akt (Ser473), Akt, phospho-p70 S6 kinase, p70 S6 kinase, phospho-S6, S6, phospho-glycogen synthase kinase 3 beta (GSK3 β), GSK3 β , insulin receptor substrate 2 (IRS2), phospho-cAMP response element-binding protein (CREB), and CREB (Cell Signaling Technology).

Real-time reverse transcription (RT)-PCR analysis

Total RNA was extracted from MIN6 cells using a RNeasy Kit (Qiagen). RNA was subjected to reverse transcription (RT), and real-time polymerase chain reaction (PCR) analysis was performed as previously described (18). The cDNA synthesized from the RNA was analyzed using a sequence detector (model 7500; Applied Biosystems) with specific primers and SYBR Green PCR Master Mix (Qiagen). The relative abundance of each mRNA was normalized to the mRNA of the housekeeping gene cyclophilin A. Primers (sense and antisense, respectively) were as follows: cyclophilin A, 5'-CAGACGCCACTGTCGCTTT-3' and 5'-TGTCTTTGGAACCTTTGTCTGCAA-3'; insulin receptor, 5'-TTTGTCATGGATGGAGGCTA-3' and 5'-CCTCATCTTGGGGTTGAACT-3'; *Irs2*, 5'-AGTCCCACATCGGGCTTGAAG-3' and 5'-GGTCTGCACGGATGACCTTAG-3'; and *PI3K-p85a*, 5'-CCTTGTCCGGGAGAGCAGTA-3' and 5'-TTGACTTCGCCGTCTACCACT-3'.

Statistical analysis

Quantitative data are presented as the mean $\pm$ standard error of the mean, and differences between means were assessed with Student's *t*-test (two-tailed). A P-value <0.05 was considered statistically significant.

RESULTS

First, we examined the effect of chlorogenic acid in a pancreatic β cell line. Western blotting showed that treatment of MIN6 cells with 0.1 mM chlorogenic acid promoted the phosphorylation of Akt, GSK3 β , and S6K compared with controls (Fig. 1A).

Next, we examined how chlorogenic acid affects pancreatic β cells in the pathogenesis of diabetes. We performed endoplasmic reticulum (ER) stress loading as a mimic of diabetic pathology. Previously, our laboratory has shown that the islets of the db/db diabetic mouse model are exposed to ER stress loading (15,19). Other groups have reported that ER stress is an important factor in pancreatic β cell failure (20). Therefore, we loaded MIN6 cells with 2 μ g/ml tunicamycin, a drug that induces ER stress, for 6 hours (16). We found that tunicamycin loading attenuated Akt phosphorylation in MIN6 cells, which was restored by the addition of 0.1 mM chlorogenic acid (Fig. 1B)(21). This suggests that chlorogenic acid may contribute to the improvement of pancreatic β cell insufficiency in the diabetic state in addition to the normal state.




Fig. 1. Effect of chlorogenic acid on insulin signaling in a pancreatic β cell line

(A) Immunoblot analysis of insulin signaling with or without 0.1 mM chlorogenic acid in MIN6 cells

(B) Immunoblot analysis of insulin signaling with or without 0.1 mM chlorogenic acid in 2 μ g/ml tunicamycin-treated MIN6 cells. CGA, chlorogenic acid; TM, tunicamycin.

In order to confirm the mechanism by which insulin signaling is upregulated, we examined the expression of insulin receptor, *PI3K*, and *Irs2* by real-time RT-PCR and found that only *Irs2* expression was significantly increased in the chlorogenic acid-treated group (Fig. 2A). To further investigate the mechanism underlying the increased expression of *Irs2*, the phosphorylation of CREB, a transcription factor of IRS2, was examined by western blotting. We found that chlorogenic acid promoted CREB phosphorylation in pancreatic β cells (Fig. 2B). These results suggest that chlorogenic acid enhances IRS2 expression and insulin signaling by promoting the phosphorylation of CREB, a transcription factor of IRS2.




Fig. 2. Chlorogenic acid treatment increases *Irs2* expression upstream of insulin signaling

(A) Quantitative RT-PCR of molecules upstream of insulin signaling with or without 0.1 mM chlorogenic acid (black and white bars, respectively) in MIN6 cells ($n = 6$, respectively)

(B) Immunoblot analysis of phosphorylated CREB with or without 0.1 mM chlorogenic acid in MIN6 cells. CGA, chlorogenic acid.

Quantitative data in (A) are the mean $\pm$ SEM ($n = 6$). * $P < 0.05$, ** $P < 0.01$ (2-tailed Student's t -test).

Next, to examine the effect of caffeine on insulin signaling, MIN6 cells were loaded with 0, 50, or 100 μ M caffeine and protein quantification was performed. Under normal conditions, caffeine had no effect on insulin signaling (Fig. 3A). In contrast, the administration of tunicamycin induced ER stress, which reduced insulin signaling, and further administration of caffeine improved insulin signaling (Fig. 3B). Therefore, caffeine may have the potential to restore insulin signaling in pathological conditions. We further tested whether chlorogenic acid loading on tunicamycin-treated MIN6 cells, in which insulin signaling was restored by caffeine, further enhanced insulin signaling. However, no additive or synergistic effects were observed when chlorogenic acid and caffeine were loaded simultaneously (Fig. 3C).




Fig. 3. Effect of caffeine on insulin signaling in MIN6 cells
 (A) Immunoblot analysis of insulin signaling with 0, 50, or 100 μ M caffeine in MIN6 cells
 (B) Immunoblot analysis of insulin signaling with 0, 50, or 100 μ M caffeine in tunicamycin-treated MIN6 cells
 (C) Immunoblot analysis of insulin signaling with or without 50 μ M caffeine and 0.1 mM chlorogenic acid in tunicamycin-treated MIN6 cells

Finally, to examine the effects of chlorogenic acid *in vivo*, we administered chlorogenic acid to C/EBP β transgenic (TG) mice, a murine model of diabetes that overexpresses the transcription factor C/EBP β specifically in pancreatic β cells. In a previous report, ER stress was shown to reduce pancreatic β cell volume and hyperglycemia (19). These mice were administered 40 mg/kg/day chlorogenic acid or saline by intraperitoneal injection for 2 weeks (22). As a result, a significant improvement in blood glucose levels was observed, although the body weight of the C/EBP β TG mice did not change (Fig. 4A, 4B). Furthermore, the volume of pancreatic β cells in C/EBP β TG mice was increased in the chlorogenic acid-treated group, and islet structure was also improved (Fig. 4C). Isolated islets from these mice showed elevated Akt phosphorylation and improved insulin signaling (Fig. 4D).




Fig. 4. Effects of chlorogenic acid on pancreatic β cells in the diabetic C/EBP β TG mouse model
 (A) Body weight ($n = 12$, respectively) at 10 and 12 weeks of age. White bars: wild-type mice as control; Black bars: C/EBP β TG mice.
 (B) Blood glucose levels ($n = 12$, respectively) at 10 and 12 weeks of age. White bars: wild-type mice as control; Black bars: C/EBP β TG mice.
 (C) Immunostaining of pancreatic islets in wild-type mice, C/EBP β TG mice, and 40 mg/kg/day chlorogenic acid-treated C/EBP β TG mice. Green: glucagon; Red: insulin.
 (D) Immunoblot analysis of insulin signaling in pancreatic islets from C/EBP β TG mice with or without 40 mg/kg/day chlorogenic acid.
 Quantitative data in (A) and (B) are the mean $\pm$ SEM ($n = 12$). * $P < 0.05$, ** $P < 0.01$ (2-tailed Student's t -test).

DISCUSSION

The effects of coffee on the human body have long been the focus of attention, and its effects on various organs have been reported. Coffee consumption is also reported to suppress the onset of psychiatric disorders such as depression (23) and cardiovascular disease (24); however, it is known to cause hypertension (25). A meta-analysis has also shown that coffee consumption reduces the incidence of cancer (26). Furthermore, coffee consumption reduces mortality from all causes (27), and there seems to be no doubt that it contributes to the health of coffee lovers.

A cohort study of 17,111 Dutch subjects was conducted in 2002 to investigate the relationship between coffee and the development of type 2 diabetes. The results showed that the incidence of type 2 diabetes was reduced in those who drank more than 7 cups of coffee per day compared with those who drank less than 2 cups per day (28). Since that report, experiments using db/db mice, a typical murine model of diabetes, and various epidemiological studies have been conducted (29,30). Based on the results of these studies, it seems safe to say that there is near-epidemiological agreement that the higher the level of coffee consumption, the lower the risk of developing type

2 diabetes. A meta-analysis reported that coffee and caffeine consumption are associated with a lower incidence of type 2 diabetes (31). Furthermore, another meta-analysis found that coffee consumption, both caffeinated and decaffeinated, was associated with a lower incidence of type 2 diabetes depending on the amount of coffee consumed (32). In addition, many reports suggest that the protective effect of coffee is due to some component other than caffeine, based on the comparison between caffeinated and decaffeinated coffee, and there are various theories on the actual component and its mechanism.

Chlorogenic acid is attracting attention as the component of coffee with the highest potential for preventing type 2 diabetes. In fact, chlorogenic acid has been shown to reduce blood glucose levels in animal and human studies (33,34). Therefore, we examined the effects of chlorogenic acid and caffeine on the survival and maintenance of pancreatic β cells. Our results showed that chlorogenic acid enhances insulin signaling in pancreatic β cells. Of particular interest was the restoration of insulin signaling in mouse islets and MIN6 cells, which was attenuated by ER stress. In a previous report, chlorogenic acid was shown to reduce ER stress in the testis, heart, and brain (35), and although the detailed molecular mechanism is unknown, the results were sufficiently reliable to indicate the reduction of ER stress in pancreatic β cells. Furthermore, chlorogenic acid is reported to enhance insulin secretion and glucose uptake in a KATP channel-independent manner in INS-1 cells, a rat pancreatic β cell line (36). In the present study, CREB phosphorylation, which regulates IRS2 expression, was enhanced by chlorogenic acid, which in turn enhanced insulin signaling. It has also been reported that chlorogenic acid loading of human neuroblastoma SH-SY5Y cells inhibits PARP-1, a molecule associated with Parkinson's disease, by enhancing CREB phosphorylation (37). This suggests that chlorogenic acid also contributes to cell proliferation via CREB in pancreatic β cells. Incidentally, although oral administration of chlorogenic acid to mice is more physiological, intraperitoneal administration was also used in the present study, since it has been adopted in many previous reports (22,35,36).

In contrast, caffeine is the most well-known component of coffee and has been shown to antagonize adenosine action through competitive inhibition (38), which has an excitatory effect on the central nervous system and also causes diuretic and heart rate elevation effects (39,40). Caffeine intake is known to be detrimental to diabetes mellitus, as it elevates blood glucose levels in association with decreased insulin sensitivity in skeletal muscle and liver (41,42), but in pancreatic β cells, caffeine increases insulin secretion and pancreatic β cell volume by enhancing insulin signaling (43). In the present study, caffeine ameliorated the ER stress-induced decrease in insulin signaling, suggesting that it may have a protective effect in pancreatic β cells.

In summary, coffee is a beverage that can be expected to have a protective effect on pancreatic β cells through the actions of chlorogenic acid and caffeine, but decaffeinated coffee may be preferable for stable blood glucose control. Further development of coffee research is expected to play a greater role in the prevention and treatment of diabetes, leading to the provision of safe and inexpensive medical care.

ACKNOWLEDGEMENTS

This work was supported by Grants-in-Aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science, and Technology (20K08860 to S.A., 20K08906 to M.K.K., and 21K08579 to Y.K.).

REFERENCES

1. International Diabetes Federation. IDF Diabetes Atlas. 9th ed. Brussels:International Diabetes Federation;2019.
2. Prentki M, Nolan CJ. Islet β cell failure in type 2 diabetes. *J Clin Invest.* 2006;116(7):1802-1812.
3. Hashimoto N, Kido Y, Uchida T, Asahara S, Shigeyama Y, Matsuda T, et al. Ablation of PDK1 in pancreatic beta cells induces diabetes as a result of loss of beta cell mass. *Nat Genet.* 2006;38(5):589-593.
4. Shigeyama Y, Kobayashi T, Kido Y, Hashimoto N, Asahara S, Matsuda T, et al. Biphasic response of pancreatic beta-cell mass to ablation of tuberous sclerosis complex 2 in mice. *Mol Cell Biol.* 2008;28(9):2971-2979.
5. Bartolomé A, Kimura-Koyanagi M, Asahara S, Guillén C, Inoue H, Teruyama K, et al. Pancreatic β -cell failure mediated by mTORC1 hyperactivity and autophagic impairment. *Diabetes.* 2014;63(9):2996-3008.
6. Uchida T, Nakamura T, Hashimoto N, Matsuda T, Kotani K, Sakaue H, et al. Deletion of Cdkn1b ameliorates hyperglycemia by maintaining compensatory hyperinsulinemia in diabetic mice. *Nat Med.* 2005;11(2):175-182.
7. Asahara S, Etoh H, Inoue H, Teruyama K, Shibutani Y, Ihara Y, et al. Paternal allelic mutation at the Kcnq1 locus reduces pancreatic β -cell mass by epigenetic modification of Cdkn1c. *Proc Natl Acad Sci U S A.* 2015;112(27):8332-8337.
8. Wu JN, Ho SC, Zhou C, Ling WH, Chen WQ, Wang CL, et al. Coffee consumption and risk of coronary heart diseases: a meta-analysis of 21 prospective cohort studies. *Int J Cardiol.* 2009;137(3):216-225.

9. Larsson SC, Virtamo J, Wolk A. Coffee consumption and risk of stroke in women. *Stroke*. 2011;42(4):908–912.
10. Arendash GW, Cao C. Caffeine and coffee as therapeutics against Alzheimer's disease. *J Alzheimers Dis*. 2010;20 Suppl 1:S117–126.
11. Choi E, Choi KH, Park SM, Shin D, Joh HK, Cho E. The Benefit of Bone Health by Drinking Coffee among Korean Postmenopausal Women: A Cross-Sectional Analysis of the Fourth & Fifth Korea National Health and Nutrition Examination Surveys. *PLoS One*. 2016;11(1):e0147762.
12. Nkondjock A. Coffee consumption and the risk of cancer: an overview. *Cancer lett*. 2009;277(2):121–125.
13. van Dam RM, Hu FB. Coffee consumption and risk of type 2 diabetes: a systematic review. *JAMA*. 2005;294(1):97–104.
14. Faerch K, Lau C, Tatens I, Pedersen OB, Jørgensen T, Borch-Johnsen K, et al. A statistical approach based on substitution of macronutrients provides additional information to models analyzing single dietary factors in relation to type 2 diabetes in danish adults: the Inter99 study. *J Nutr*. 2005;135(5):1177–1182.
15. Greenberg JA, Axen KV, Schnoll R, Boozer CN. Coffee, tea and diabetes: the role of weight loss and caffeine. *Int J Obes (Lond)*. 2005;29(9):1121–1129.
16. Matsuda T, Kido Y, Asahara S, Kaisho T, Tanaka T, Hashimoto N, et al. Ablation of C/EBPbeta alleviates ER stress and pancreatic beta cell failure through the GRP78 chaperone in mice. *J Clin Invest*. 2010;120(1):115–126.
17. Asahara S, Shibutani Y, Teruyama K, Inoue HY, Kawada Y, Etoh H, et al. Ras-related C3 botulinum toxin substrate 1 (RAC1) regulates glucose-stimulated insulin secretion via modulation of F-actin. *Diabetologia*. 2013;56(5):1088–1097.
18. Kanno A, Asahara S, Furubayashi A, Masuda K, Yoshitomi R, Suzuki E, et al. GCN2 regulates pancreatic β cell mass by sensing intracellular amino acid levels. *JCI Insight*. 2020;5(9):e128820.
19. Asahara S, Matsuda T, Kido Y, Kasuga M. Increased ribosomal biogenesis induces pancreatic beta cell failure in mice model of type 2 diabetes. *Biochem Biophys Res Commun*. 2009;381(3):367–371.
20. Oyadomari S, Koizumi A, Takeda K, Gotoh T, Akira S, Araki E, et al. Targeted disruption of the Chop gene delays endoplasmic reticulum stress-mediated diabetes. *J Clin Invest*. 2002;109(4):525–532.
21. Pavlica S, Gebhardt R. Protective effects of ellagic and chlorogenic acids against oxidative stress in PC12 cells. *Free Radic Res*. 2005;39(12):1377–1390.
22. Ohkawara T, Takeda H, Nishihira J. Protective effect of chlorogenic acid on the inflammatory damage of pancreas and lung in mice with l-arginine-induced pancreatitis. *Life Sci*. 2017;190:91–96.
23. Wang L, Shen X, Wu Y, Zhang D. Coffee and caffeine consumption and depression: A meta-analysis of observational studies. *Aust N Z J Psychiatry*. 2016;50(3):228–242.
24. Ding M, Bhupathiraju SN, Satija A, van Dam RM, Hu FB. Long-term coffee consumption and risk of cardiovascular disease: a systematic review and a dose-response meta-analysis of prospective cohort studies. *Circulation*. 2014;129(6):643–659.
25. Nurminen ML, Niittynen L, Korpela R, Vapaatalo H. Coffee, caffeine and blood pressure: a critical review. *Eur J Clin Nutr*. 1999;53(11):831–839.
26. Zhao LG, Li ZY, Feng GS, Ji XW, Tan YT, Li HL, et al. Coffee drinking and cancer risk: an umbrella review of meta-analyses of observational studies. *BMC Cancer*. 2020;20(1):101.
27. Gunter MJ, Murphy N, Cross AJ, Dossus L, Dartois L, Fagherazzi G, et al. Coffee Drinking and Mortality in 10 European Countries: A Multinational Cohort Study. *Ann Intern Med*. 2017;167(4):236–247.
28. van Dam RM, Feskens EJM. Coffee consumption and risk of type 2 diabetes mellitus. *Lancet*. 2002;360(9344):1477–1478.
29. Jung UJ, Lee MK, Park YB, Jeon SM, Choi MS. Antihyperglycemic and antioxidant properties of caffeic acid in db/db mice. *J Pharmacol Exp Ther*. 2006;318(2):476–483.
30. Kato M, Noda M, Inoue M, Kadowaki T, Tsugane S; JPHC Study Group. Psychological factors, coffee and risk of diabetes mellitus among middle-aged Japanese: a population-based prospective study in the JPHC study cohort. *Endocr J*. 2009;56(3):459–468.
31. Jiang X, Zhang D, Jiang W. Coffee and caffeine intake and incidence of type 2 diabetes mellitus: a meta-analysis of prospective studies. *Eur J Nutr*. 2014;53(1):25–38.
32. Ding M, Bhupathiraju SN, Chen M, van Dam RM, Hu FB. Caffeinated and decaffeinated coffee consumption and risk of type 2 diabetes: a systematic review and a dose-response meta-analysis. *Diabetes Care*. 2014;37(2):569–586.
33. Mansour A, Mohajeri-Tehrani MR, Samadi M, Qorbani M, Merat S, Adibi H, et al. Effects of supplementation with main coffee components including caffeine and/or chlorogenic acid on hepatic, metabolic, and inflammatory indices in patients with non-alcoholic fatty liver disease and type 2 diabetes: a randomized, double-blind, placebo-controlled, clinical trial. *Nutr J*. 2021;20(1):35.

34. Yin XL, Xu BQ, Zhang YQ. *Gynura divaricata* rich in 3, 5-/4, 5-dicaffeoylquinic acid and chlorogenic acid reduces islet cell apoptosis and improves pancreatic function in type 2 diabetic mice. *Nutr Metab (Lond)*. 2018;15:73.
35. Kazaz IO, Demir S, Kerimoglu G, Colak F, Turkmen Alemdar N, Yilmaz Dogan S, et al. Chlorogenic acid ameliorates torsion/detorsion-induced testicular injury via decreasing endoplasmic reticulum stress. *J Pediatr Urol*. 2022;18(3):289.e1–289.e7.
36. Tousch D, Lajoix AD, Hosy E, Azay-Milhau J, Ferrare K, Jahannault C, et al. Chicoric acid, a new compound able to enhance insulin release and glucose uptake. *Biochem Biophys Res Commun*. 2008;377(1):131–135.
37. Kim H, Park J, Kang H, Yun SP, Lee YS, Lee YI, et al. Activation of the Akt1-CREB pathway promotes RNF146 expression to inhibit PARP1-mediated neuronal death. *Sci Signal*. 2020;13(663):eaax7119.
38. Schwarzschild MA, Chen JF, Ascherio A. Caffeinated clues and the promise of adenosine A(2A) antagonists in PD. *Neurology*. 2002;58(8):1154–1160.
39. Buscemi S, Mattina A, Tranchina MR, Verga S. Acute effects of coffee on QT interval in healthy subjects. *Nutrition J*. 2011;10:15.
40. Rieg T, Steiglele H, Schnermann J, Richter K, Osswald H, Vallon V. Requirement of intact adenosine A1 receptors for the diuretic and natriuretic action of the methylxanthines theophylline and caffeine. *J Pharmacol Exp Ther*. 2005;313(1):403–409.
41. Lee S, Min JY, Min KB. Caffeine and Caffeine Metabolites in Relation to Insulin Resistance and Beta Cell Function in U.S. Adults. *Nutrients*. 2020;12(6):1783.
42. Guarino MP, Ribeiro MJ, Sacramento JF, Conde SV. Chronic caffeine intake reverses age-induced insulin resistance in the rat: effect on skeletal muscle Glut4 transporters and AMPK activity. *Age (Dordr)*. 2013;35(5):1755–1765.
43. Park S, Jang JS, Hong SM. Long-term consumption of caffeine improves glucose homeostasis by enhancing insulinotropic action through islet insulin/insulin-like growth factor 1 signaling in diabetic rats. *Metabolism*. 2007;56(5):599–607.