



# Plant Flavonoid Production in Bacteria and Yeasts

Isogai, Shota  
Tominaga, Masahiro  
Kondo, Akihiko  
Ishii, Jun

---

**(Citation)**

Frontiers in Chemical Engineering, 4:880694

**(Issue Date)**

2022-07-08

**(Resource Type)**

journal article

**(Version)**

Version of Record

**(Rights)**

© 2022 Isogai, Tominaga, Kondo and Ishii.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited a...

**(URL)**

<https://hdl.handle.net/20.500.14094/0100482166>





# Plant Flavonoid Production in Bacteria and Yeasts

Shota Isogai<sup>1,2†</sup>, Masahiro Tominaga<sup>1,3</sup>, Akihiko Kondo<sup>1,2,3,4,5</sup> and Jun Ishii<sup>1,2,3\*</sup>

<sup>1</sup>Graduate School of Science, Technology and Innovation, Kobe University, Kobe, Japan, <sup>2</sup>Technology Research Association of Highly Efficient Gene Design (TRAHED), Kobe, Japan, <sup>3</sup>Engineering Biology Research Center, Kobe University, Kobe, Japan, <sup>4</sup>Department of Chemical Science and Engineering, Graduate School of Engineering, Kobe University, Kobe, Japan, <sup>5</sup>RIKEN Center for Sustainable Resource Science, Yokohama, Japan

## OPEN ACCESS

### Edited by:

Shang-Tian Yang,  
The Ohio State University,  
United States

### Reviewed by:

Jingwen Zhou,  
Jiangnan University, China  
Yu Wang,  
Tianjin Institute of Industrial  
Biotechnology (CAS), China

### \*Correspondence:

Jun Ishii  
junjun@port.kobe-u.ac.jp

### †Present address:

Shota Isogai,  
Division of Biological Science,  
Graduate School of Science and  
Technology, Nara Institute of Science  
and Technology, Ikoma, Japan

### Specialty section:

This article was submitted to  
Biochemical Engineering,  
a section of the journal  
Frontiers in Chemical Engineering

Received: 21 February 2022

Accepted: 09 June 2022

Published: 08 July 2022

### Citation:

Isogai S, Tominaga M, Kondo A and  
Ishii J (2022) Plant Flavonoid  
Production in Bacteria and Yeasts.  
Front. Chem. Eng. 4:880694.  
doi: 10.3389/fceng.2022.880694


Flavonoids, a major group of secondary metabolites in plants, are promising for use as pharmaceuticals and food supplements due to their health-promoting biological activities. Industrial flavonoid production primarily depends on isolation from plants or organic synthesis, but neither is a cost-effective or sustainable process. In contrast, recombinant microorganisms have significant potential for the cost-effective, sustainable, environmentally friendly, and selective industrial production of flavonoids, making this an attractive alternative to plant-based production or chemical synthesis. Structurally and functionally diverse flavonoids are derived from flavanones such as naringenin, pinocembrin and eriodictyol, the major basic skeletons for flavonoids, by various modifications. The establishment of flavanone-producing microorganisms can therefore be used as a platform for producing various flavonoids. This review summarizes metabolic engineering and synthetic biology strategies for the microbial production of flavanones. In addition, we describe directed evolution strategies based on recently-developed high-throughput screening technologies for the further improvement of flavanone production. We also describe recent progress in the microbial production of structurally and functionally complicated flavonoids *via* the flavanone modifications. Strategies based on synthetic biology will aid more sophisticated and controlled microbial production of various flavonoids.

**Keywords:** microbial production, flavonoid, naringenin, metabolic engineering, synthetic biology, directed evolution, biosensor

## 1 INTRODUCTION

Flavonoids are a major group of secondary metabolites in plants that shares a diphenylpropane backbone (C<sub>6</sub>-C<sub>3</sub>-C<sub>6</sub>) in which three methylene units link two aromatic rings. Flavonoids are divided into several subclasses based on modifications to the backbone: flavanone, flavone, dihydroflavonol, flavonol, flavane-3,4-diol, anthocyanidin, flavane-3-ol, isoflavone and the derivatives of these aglycones such as flavonoid glycoside and prenylated flavonoid (Figure 1). To date, approximately ten thousand types of flavonoids have been identified from plants (Dixon and Pasinetti, 2010) and are promising as pharmaceuticals and food supplements due to their health-promoting biological activities (Harborne and Williams, 2000; Fraga et al., 2019).


Flavonoids are primarily isolated from plants or generated by organic synthesis; however, neither plant-based isolation nor chemical synthesis are cost-effective and sustainable because 1) plants produce small quantities of flavonoids and large expanses of land and longtime are



required to grow these plants, making the isolation of flavonoids, including harvest, extraction and purification, arduous and expensive, and 2) organic synthesis is often challenging, given the structural complexity of flavonoids. As an alternative, flavonoids can be produced using recombinant microbes in a cost-effective and sustainable process due to their selective production of valuable compounds from inexpensive substrates (Liu et al., 2017b).

Most botanical flavonoids are biosynthesized from flavanone by various enzymatic modifications, such as hydroxylation, dehydration, methylation, glycosylation, and prenylation (Figure 1). The establishment of flavanone-producing microorganisms can therefore provide a platform for producing structurally complicated flavonoids. Flavonoid biosynthesis occurs in two steps: flavanone biosynthesis, and its modification(s) (Winkel, 2006; Tohge et al., 2017). Naringenin, the most common flavanone, is synthesized through the aromatic amino acids phenylalanine and

tyrosine by sequential enzymatic reactions in plants (Figure 2). First, phenylalanine ammonia lyase (PAL) catalyzes the deamination of phenylalanine to form *trans*-cinnamic acid, then *trans*-cinnamic acid is hydroxylated by cinnamate-4-hydroxylase (C4H, cytochrome P450) coupled with cytochrome P450 reductase (CPR) to yield *p*-coumaric acid. Alternatively, *p*-coumaric acid is synthesized by direct deamination of tyrosine by the enzymatic reaction of tyrosine ammonia lyase (TAL). Then, 4-coumaroyl-CoA ligase (4CL) catalyzes the ligation of CoA to *p*-coumaric acid to form *p*-coumaroyl-CoA. Chalcone synthase (CHS) subsequently catalyzes the Claisen-type condensation of *p*-coumaroyl-CoA with three molecules of malonyl-CoA to yield naringenin chalcone, then chalcone isomerase (CHI) converts naringenin chalcone to naringenin by formation of the C-ring. PAL has been identified in plants, whereas TAL has not. Alternatively, plant uses a bifunctional phenylalanine/tyrosine ammonia-lyase (PTAL) for *p*-coumaric acid synthesis from tyrosine, which catalyzes the deamination of both



phenylalanine and tyrosine (Barros and Dixon, 2020). The biosynthesis of flavanones uses a common route with the naringenin biosynthesis except for the difference in phenylpropanoyl-CoA precursors (i.e., *p*-coumaroyl-CoA for naringenin) for chalcone formation. In brief, pinocembrin and eriodictyol are produced from *trans*-cinnamic acid and caffeic acid via the phenylpropanoyl-CoA (*trans*-cinnamoyl-CoA and caffeoyl-CoA, respectively) by the enzymatic reactions of 4CL, CHS and CHI (Figure 2). Since flavanones share thus the same precursors, amino acids (phenylalanine and tyrosine) and malonyl-CoA, metabolic engineering strategies and synthetic biology approaches have been employed to increase the intracellular availability of these precursors in microbial production of flavonoids especially *de novo* synthesis of them.

The modification enzymes attach various functional groups to the flavanone core structure to form diverse flavonoids. The

modification steps are categorized into two types: variation in the heterocyclic C-ring by hydroxylation and dehydration, and attachment of diverse side chains such as glycosylation and prenylation (Figure 1). For example, flavone synthase (FNS) catalyzes the dehydration of the C-ring in flavanone to flavone, whereas flavanone-3-hydroxylase (F3H) attaches a hydroxyl group onto C3 in the C-ring of flavanone to yield dihydroflavonol, which is then converted to flavonol via dehydration by flavanol synthase (FLS). Dihydroflavonol is also converted to anthocyanidin or flavane-3-ol by sequential modifications by dihydroflavonol 4-reductase (DFR) along with anthocyanidin synthase (ANS) or leucoanthocyanidin reductase (LAR). Several carbon atoms in the B-ring of naringenin are hydroxylated to yield structurally different compounds during these modification steps such as hydroxylation of C3' or both C3' and C5' in the B-ring of flavonoids by flavanone-3'-hydroxylase (F3'H) and flavanone-

**TABLE 1 |** Flavanones production using different microbes.

Enzymes for flavanone biosynthesis <sup>a</sup>	Carbon source (+precursor supply) <sup>b</sup>	Pathway engineering for increasing availabilities of aromatic amino acids and malonyl-CoA	Product <sup>c</sup>	Yield (mg/l)	References
<i>Escherichia coli</i>					
RrPAL, Sc4CL, GeCHS	Glu	—	NAR	0.0833	Hwang et al., (2003)
	Glu (+Phe)	—	PIN	0.157	
	Glu (+Tyr)	—	NAR	0.0096	
			PIN	0.752	
			NAR	0.453	
RsTAL, At4CL, AtCHS	Glu/Gly	—	PIN	0.0389	Watts et al., (2004)
	Gly	—	NAR	7.53	
Pc4CL, PhCHS, MsCHI	Glu (+pCA)	Overexpression of ACC and ACS	NAR	20.8	Leonard et al., (2007)
	Glu (+tCA)		PIN	119	
	Glu (+CA)		EDT	429	Leonard et al., (2008)
	Glu (+pCA)	Overexpression of ACC and acetate kinase-phosphate acetyltransferase	NAR	52	
	Glu (+tCA)		PIN	104	
	Glu (+CA)		EDT	405	
	Glu (+pCA)	Bypass pathway from malonate by MatB and MatC; Downregulation of transacylases FabB and FabF by cerulenin	NAR	45	
	Glu (+pCA)	Overexpression of ACS; Knockout of genes <sup>d</sup> in TCA cycle and ethanol synthesis	NAR	186	
	Glu (+CA)	Overexpression of ACC and PDH; Downregulation of TCA cycle <sup>e</sup> ; Upregulation of glycolysis <sup>f</sup>	EDT	270	
At4CL, PhCHS		Downregulation of transacylase FabD by antisenseRNA	NAR	150	Fowler et al., (2009)
At4CL, PhCHS, CmCHI		—	—	91.3	Yang et al., (2015)
				>200	Xiong et al., (2017)
RgTAL, Pc4CL, PhCHS, MsCHI	Glu (+Tyr)	Bypass pathway from malonate by MatB and MatC; Downregulation of FabB and FabF by antisenseRNA	NAR	272	Hwang et al., (2021)
		Downregulation of malonyl-CoA consuming pathways by CRISPRi <sup>g</sup>		391	Wu et al., (2014a)
		Overexpression of ACC; Downregulation of fatty acid synthesis by CRISPRi <sup>h</sup>		4,216	Wu et al., (2015)
RgPAL, Pc4CL, PhCHS, MsCHI, GhF3'H, CrCPR		Overexpression of ACC and ACS; Deletion of <i>ackA</i>	EDT	1,073	Wu et al., (2021)
FJTAL, Gm4CL, AtCHS, AtCHI	Gly	Deletion of <i>tyrR</i> and <i>pheLA</i> ; Expression of PpsA, AroG <sup>D146N</sup> and TyrA <sup>M53I/A354V</sup>	NAR	107	Zhu et al., (2014)
AtPAL, Gm4CL, AtCHS, AtCHI	Gly (+Phe)	Downregulation of transacylases by cerulenin	PIN	484	Dunstan et al., (2020)
Gm4CL, AtCHS, AtCHI	Gly (+CA)		EDT	198	
FJTAL, Gm4CL, AtCHS, AtCHI, SeC3H, Pk3OMT	Gly		HEDT	88	
RgTAL, Pc4CL, PhCHS, MsCHI	Glu	Expression of AroG <sup>D146N</sup> and TyrA <sup>M53I/A354V</sup> ; Deletion of <i>tyrR</i> and <i>pheA</i> ; Bypass pathway from malonate by MatB and MatC	NAR	17	
		Expression of AroG <sup>D146N</sup> and TyrA <sup>M53I/A354V</sup> ; Downregulation of FabB and FabF by cerulenin		46	Santos et al., (2011)
		Expression of AroG <sup>D146N</sup> and TyrA <sup>M53I/A354V</sup> ; Bypass pathway from malonate by MatB and MatC		84	
		Downregulation of malonyl-CoA; Consuming pathways by CRISPRi <sup>i</sup>		100.6	Wu et al., (2014b)
AtPAL, Os4CL, PeCHS	Glu	Overexpression of ACC	PIN	97	Kim et al., (2014)
RgPAL, Pc4CL, PhCHS, MsCHI		Expression of AroF and PheA <sup>T326P</sup> ; Bypass pathway from malonate by MatB and MatC		40	Wu et al., (2013)
		Downregulation of malonyl-CoA; Consuming pathways by CRISPRi <sup>j</sup>		432	Wu et al., (2016a)
		Overexpression of ACC, ACS and FabF		526	Wu et al., (2016b)
BoPAL, Pc4CL, GuCHS, MsCHI		Overexpression of FabF; Supplementation of cerulenin		67.8	Cao et al., (2016a)
RmPAL, Sc4CL, GuCHS, MsCHI				29.9	Cao et al., (2016b)
SeTAL, At4CL, PhCHS, AtCHI	Gly	Expression of AroG <sup>D146N</sup> ; Knockout of <i>tyrR</i> ; Knocked down of fatty acid metabolism regulator FadR	NAR	103.8	Yang et al., (2018)
TcTAL, Pc4CL, PhCHS, MsCHI	Glu	Bypass pathway from malonate by MatB and MatC	NAR	191.9	Zhou et al., (2019)
		Optimization in expression levels of genes in malonyl CoA and fatty acid synthesis <sup>l</sup>		524	Zhou et al., (2021)
		Expression of AroG <sup>D146N</sup> and TyrA <sup>M53I/A354V</sup> ; Knockout of <i>tyrR</i> and <i>pheA</i> ; Optimization in expression levels of genes in malonyl CoA and fatty acid synthesis <sup>k</sup>		588	Zhou et al., (2020)

(Continued on following page)

**TABLE 1 |** (Continued) Flavanones production using different microbes.

Enzymes for flavanone biosynthesis <sup>a</sup>	Carbon source (+precursor supply) <sup>b</sup>	Pathway engineering for increasing availabilities of aromatic amino acids and malonyl-CoA	Product <sup>c</sup>	Yield (mg/l)	References
<i>Corynebacterium glutamicum</i>					
<u>FJTAL</u> , <u>Pc4CL</u> , <u>PhCHS</u> , <u>PhCHI</u>	Glu (+pCA)	Downregulation of transacylases by cerulenin	NAR	35	Kallscheuer et al., (2016b)
	Glu	Expression of <i>E. coli</i> AroH in addition to the above		32	Milke et al., (2019)
		Expression of <i>E. coli</i> AroH; Overexpression of ACC; Deletion of <i>fasR</i> and <i>sdhCAB</i> and downregulation of <i>gltA</i> <sup>f</sup>		24	
<i>Streptomyces albus</i>					
RcTAL, Sc4CL, GmCHS, GmCHI.	Glu	—		—	Marin et al., (2018)
<i>Saccharomyces cerevisiae</i>					
AtC4H, At4CL, PhCHS, PhCH	Glu (+tCA)	—	NAR	0.2	Yan et al., (2005b)
	Glu (+pCA)	—		28.3	
RtPAL, At4CL, HaCHS	Glu	—	NAR	7	Jiang et al., (2005)
			PIN	0.8	
Ptr-PdePAL, GmC4H, Ptr-PdePAL, Gm4CL, GmCHS, GmCHI	Glu (+Phe)	—	NAR	8.9	Trantas et al., (2009)
	Glu (+pCA)			15.6	
Pc4CL, PhCHS, PhCHI	Glu (+pCA)	Co-expression of ACS with Cab1 and Ald6	NAR	12.5	Liu et al., (2017a)
AtPAL, AtC4H, AtCPR, At4CL, AtCHS, AtCHI, RcTAL	Glu	Deletion of <i>ARO3</i> , <i>ARO10</i> , <i>PDC5</i> , and <i>PDC6</i> ; Expression of Aro4 <sup>G226S</sup>	NAR	113	Koopman et al., (2012)
<u>FJTAL</u> , <u>At4CL</u> , <u>HaCHS</u> , <u>MsCHI</u>	Suc/Gly	Deletion of <i>PHA2</i> , <i>ARO10</i> and <i>PDC5</i> ; Expression of Aro4 <sup>G226S</sup>	NAR	84	Lyu et al., (2017)
FJTAL, <u>Pc4CL</u> , <u>PhCHS</u> , <u>MsCHI</u>	Glu	Deletion of <i>ARO10</i> and <i>PDC5</i> ; Expression of Aro4 <sup>G226S</sup> and Aro7 <sup>G141S</sup>	NAR	1.55	Rodriguez et al., (2017)
FJTAL, At4CL, AtCHS, AtCHI	Glu	Deletion of <i>ARO10</i> and <i>PDC5</i> ; Expression of Aro4 <sup>G226S</sup> and Aro7 <sup>G141S</sup> ; Overexpression of ACC with <i>E. coli</i> PDH	NAR	144	Du et al., (2020)
At4CL, HaCHS, PhCHI	Glu (+pCA)	—	NAR	52	Wang et al., (2019)
Pc4CL, PhCHS, MsCHI	Glu (+pCA)	—	NAR	1,210	Gao et al., (2020)
FJTAL, Pc4CL, PhCHS, MsCHI	Glu	Deletion of <i>ARO10</i> ; Expression of Aro4 <sup>G226S</sup> and Aro7 <sup>G141S</sup> ; Overexpression of Pex11 <sup>m</sup>	NAR	1,129	Zhang et al., (2021)
<i>Yarrowia lipolytica</i>					
RtTAL, Pc4CL, PhCHS, MsCHI	Glu	Overexpression of Aro1 <sup>n</sup> ; Overexpression of ACC and ACS	NAR	254	Lv et al., (2019)
		Overexpression of Aro1; Overexpression of ACC and ACS; Downregulation of malonyl-CoA consuming pathways by CRISPR <sup>i</sup>		158	Lv et al., (2020)
		—		100–125	Bowman et al., (2021)
SeTAL, Nt4CL, HsCHS		Expression of tyrosine-insensitive Aro4 (Aro4 <sup>K221L</sup> )		898	Palmer et al., (2020)
RgTAL, At4CL, AtCHS, AtCHI	Glu/Xyl	Overexpression of ACC and Pex10 <sup>p</sup>	NAR	715	Wei et al., (2020)

<sup>a</sup>Abbreviations are indicated in **Table 4**. Codon of underlined enzymes are optimized for each host strain.

<sup>b</sup>Glu, Gly, Suc, Xyl, Phe, Tyr, tCA, pCA, and CA represent glucose, glycerol, xylose, phenylalanine, tyrosine, trans-cinnamic acid, p-coumaric acid, and caffeic acid respectively.

<sup>c</sup>NAR, PIN, EDT, and HEDT represent naringenin, pinocembrin, eriodictyol and homoeriodictyol respectively.

<sup>d</sup>Knocked out genes are *sdhABCD*, *citE* (TCA, cycle), and *adhE* (ethanol synthesis) that encode succinate dehydrogenase, citrate lyase, and alcohol dehydrogenase respectively.

<sup>e</sup>Downregulated genes in TCA cycle are *acnA/acnB*, *fumB/C*, *mdh*, *scpC*, and *sucC + sucD* that encode aconitase, fumarase, malate dehydrogenase, propionyl-CoA, succinate CoA transferase, and succinyl-CoA synthetase respectively.

<sup>f</sup>Upregulated genes in glycolysis are *gapA* and *pgk* that encode glyceraldehyde-3-phosphate dehydrogenase and phosphoglycerate kinase respectively.

<sup>g</sup>Target genes are *ppsA*, *eno*, *adhE*, *mdh*, *fumC*, *sdhA*, *sucC*, *citE*, *fabH*, *fabB*, *fabF*, and *fabI*. *ppsA* and *eno* encode phosphoenolpyruvate synthase and phosphopyruvate hydratase respectively. *fabH*, *fabB*, *fabF*, and *fabI* are genes involved in fatty acid synthesis.

<sup>h</sup>Target genes are *fabD*, *fabH*, *fabB*, *fabF*, and *fabI*.

<sup>i</sup>Target genes are *eno*, *adhE*, *mdh*, *fumC*, *sucC*, *fabB*, and *fabF*.

<sup>j</sup>Target genes are ACC and ACS in malonyl-CoA synthesis and *fabD*, *acpS*, *acpT*, *acpP*, and *acpH* in fatty acid synthesis. *acpS*, *acpT*, *acpP*, and *acpH* are involved in acyl carrier protein formation.

<sup>k</sup>Target genes are ACC and ACS in malonyl-CoA synthesis and *fabD*, *acpS*, *acpT*, *acpP*, and *acpH* in fatty acid synthesis. *acpS*, *acpT*, *acpP*, and *acpH* are involved in acyl carrier protein formation.

<sup>l</sup>*fasR*, *sdhCAB*, and *gltA* encode transcriptional repressor of fatty acid synthesis, succinate dehydrogenase complex and citrate synthase respectively.

<sup>m</sup>PEX11 encodes peroxisomal proliferation and fatty acids transport pathway protein and overexpression of Pex11 enhances  $\beta$ -oxidation.

<sup>n</sup>ARO1 encodes pentafunctional AroM that catalyzes enzymatic reactions from DAHP to 5-enolpyruvoyl-shikimate 3-phosphate.

<sup>o</sup>Target genes are *fas1*, *fas2*, and *fabD*.

<sup>p</sup>PEX10 encodes peroxisome biogenesis factor and overexpression of Pex10 enhances  $\beta$ -oxidation.

**TABLE 2 |** Co-culturing system for production of naringenin and its derivatives.

Production host (function of strain)	Precursor <sup>a</sup>	Enzymes for flavonoid biosynthesis <sup>b</sup> and Pathway engineering for increasing precursors	Flavonoid product	Yield (mg/l)	References
<i>E. coli</i> (Supplying tyrosine)	Xyl	Knockout of <i>tyrR</i> and <i>pheA</i> ; Expression of AroG <sup>D146N</sup> and TyrA <sup>M53I/A354V</sup>	Naringenin	21.2	Zhang et al., (2017)
<i>S. cerevisiae</i> (Synthesis of naringenin)	Tyr	<u>RtTAL</u> , <u>Pc4CL</u> , <u>PhCHS</u> , <u>PhCHI</u>			
<i>E. coli</i> <sup>c</sup> ( <i>p</i> -Coumaric acid synthesis)	Glu	<u>RgTAL</u> ; Expression of AroG <sup>D146N</sup> and TyrA <sup>M53I/A354V</sup> ; Deletion of <i>tyrR</i> and <i>pheA</i>	—	—	Jones et al., (2017)
<i>E. coli</i> <sup>d</sup> (Naringenin synthesis)	<i>p</i> -CA	<u>Pc4CL</u> , <u>PhCHS</u> , <u>MsCHI</u> ; Deletion of <i>fumC</i> and <i>sucC</i> ; Bypass pathway from malonate by MatB and MatC	Naringenin	—	
<i>E. coli</i> <sup>e</sup> (Flavane-3-ol synthesis)	NAR	<u>F3H</u> , <u>DFR</u> , and <u>LAR</u>	Afzelechin	26	
<i>E. coli</i> <sup>f</sup> (Anthocyanin synthesis)	Afzelechin	<u>ANS</u> and <u>3-O-GT</u> (glucose as sugar donor)	Callistephin	9.5	
<i>S. cerevisiae</i> (Synthesis of naringenin)	Glu	<u>FjTAL</u> , <u>At4CL</u> , <u>AtCHS</u> , <u>AtCHI</u> ; Deletion of <i>ARO10</i> and <i>PDC5</i> ; Expression of Aro4 <sup>G226S</sup> and Aro7 <sup>G141S</sup> ; Overexpression of ACC with <i>E. coli</i> PDH	Naringenin	144	Du et al., (2020)
<i>S. cerevisiae</i> (Modification of naringenin)	NAR	<u>F3H</u> , <u>FLS</u>	Kaempferol	168	
	<u>F3H</u> , <u>F3'H</u> , <u>FLS</u>	<u>F3H</u> , <u>FLS</u>	Quercetin	154	
	<u>F3H</u> , <u>F3'H</u> , <u>F3'5'H</u> , <u>FLS</u>	Myricetin	145		
	<u>F3H</u> , <u>DFR</u> , <u>ANS</u>	Pelargonidin	33		
	<u>F3H</u> , <u>DFR</u> , <u>F3'H</u> , <u>ANS</u>	Cyanidin	32		

<sup>a</sup>Glu, Xyl, Tyr, *p*CA, and NAR represent glucose, xylose, tyrosine, *p*-coumaric acid and naringenin respectively.

<sup>b</sup>Naringenin biosynthetic and modification enzymes are represented by double- and dotted-underline respectively. Abbreviations for naringenin biosynthetic enzymes are indicated in **Table 4**.

<sup>c</sup>Derived from Santos et al. Santos, C.N., Koffas, M., and Stephanopoulos, G. (2011). Optimization of a heterologous pathway for the production of flavonoids from glucose. *Metab Eng* 13, 392–400.

<sup>d</sup>Derived from Xu et al. Xu, P., Ranganathan, S., Fowler, Z.L., Maranas, C.D., and Koffas, M.A., *Ibid*. Genome-scale metabolic network modeling results in minimal interventions that cooperatively force carbon flux towards malonyl-CoA. 578–587.

<sup>e</sup>Derived from Zhao et al. Zhao, S., Jones, J.A., Lachance, D.M., Bhan, N., Khalidi, O., Venkataraman, S., Wang, Z., and Koffas, M.A.G. (2015). Improvement of catechin production in *Escherichia coli* through combinatorial metabolic engineering. *Ibid*. 28, 43–53.

<sup>f</sup>Derived from Cress et al. Cress, B.F., Leitz, Q.D., Kim, D.C., Amore, T.D., Suzuki, J.Y., Linhardt, R.J., and Koffas, M.A. (2017). CRISPRi-mediated metabolic engineering of *E. coli* for *O*-methylated anthocyanin production. *Microb Cell Fact* 16, 10.

3'5'-hydroxylase (F3'5'H) respectively. For example, eriodictyol can be synthesized by hydroxylating C3' of naringenin (Amor et al., 2010) as well as synthesized from caffeic acid described above (**Figure 2**). Similarly, myricetin, one of flavonol, is biosynthesized from naringenin by combination of F3H, FLS, and F3'5'H (Leonard et al., 2006). Additionally, isoflavone synthase (IFS), a P450 monooxygenase, catalyzes the transposition of the B-ring from C2 to C3 in the C-ring.

Glycosylation and prenylation convert naringenin-derived flavonoids into more complex flavonoids in plants (**Figure 1**). Glycosylation increases the solubility and bioactivity of flavonoids, with the increased solubility contributing to their utilization upon eating the plants. Plant glycosyltransferase produces flavonoid glycosides mainly by linking the activated sugar moiety to the oxygen atom of the flavonoid backbone (*O*-glycosylation) or by directly forming a C-C linkage between the sugar and the flavonoid (*C*-glycosylation). An alternative modification to glycosylation, prenylation, attaches one or more prenyl groups, such as dimethylallyl (C<sub>5</sub>) and geranyl (C<sub>10</sub>) moieties, to the flavonoid backbone. Prenylation increases the

liposolubility of flavonoids, contributing to the maintenance of these compounds in cell membranes compared with non-prenylated flavonoids (Mukai et al., 2012). Additionally, the attachment position and chain length of the prenyl moieties, and further modification after prenylation, can alter and enhance the biological activity of flavonoids. These modifications are essential for the various biological activities of flavonoids and their structural diversity.

In this review, we summarize current engineering strategies for improving the microbial production of plant flavonoids. First, we describe metabolic engineering strategies, including the screening of host organisms and plant-derived enzymes, and pathway designs using these organisms and enzymes mainly focusing on naringenin. Next, we describe evolutionary strategies utilizing high-throughput screening techniques to identify cells with improved production of naringenin and its derivatives. Moreover, we explain how genetic circuits that regulate pathway gene expression improve the production of flavonoids and their precursors. Finally, we describe recent progress in the microbial production of structurally complicated flavonoids using various modification enzymes.

**TABLE 3 |** Microbial production of diversified flavonoids.

Production host	Modification enzyme	Substrate	Product	Reference
<i>Hydroxylation/dehydration</i>				
<i>E. coli</i>	F3H, DFR and ANS	Naringenin	Cyanidin	Yan et al., (2005a)
	F3H, FLS	Eriodictyol	Pelargonidin	
		<i>p</i> -Coumaric acid <sup>a</sup>	Dihydrokaempferol, Kaempferol	Leonard et al., (2006)
		Naringenin		
		Caffeic acid <sup>a</sup>	Eriodictyol	
		Eriodictyol	Dihydroquercetin, Quercetin	
	F3H, FLS, and F3'5'H with CPR	<i>p</i> -Coumaric acid <sup>a</sup>	Dihydrokaempferol, Kaempferol, Dihydroquercetin, Quercetin	
		Naringenin	Dihydrokaempferol, Kaempferol, Dihydroquercetin, Quercetin, Myricetin	
		Eriodictyol	Dihydroquercetin, Quercetin, Myricetin	
		FLS	Dihydrokaempferol	Kaempferol
	F3H, DFR, and LAR	Naringenin	Afzelechin	Zhao et al., (2015)
		Eriodictyol	Catechin	
	F3H and FLS	Tyrosine <sup>p</sup>	Kaempferol	Boada et al., (2020)
<i>C. glutamicum</i>	F3H and FLS	<i>p</i> -Coumaric acid <sup>a</sup>	Kaempferol	Kallscheuer et al., (2017)
		Caffeic acid <sup>a</sup>	Quercetin	
<i>Lactococcus lactis</i>	ANS	Catechin	Cyanidin	Solopova et al., (2019)
	F3H and FLS	Glucose <sup>p</sup>	Kaempferol	Marin et al., (2018)
<i>S. albus</i>	F3H, FLS, and F3'H	Quercetin		
	F3H, FLS, F3'H, and F3'5'H	Myricetin		
<i>Streptomyces</i> bacteria	Endogenous hydroxylase	Naringenin; Daidzein; Genistein	Hydroxynaringenin	Shrestha et al., (2021)
		7,3',4'-trihydroxyisoflavone		
		Hydroxygenistein		
<i>S. cerevisiae</i>	FNS and CPR	<i>trans</i> -Cinnamic acid <sup>a</sup>	Chrysin	Leonard et al., (2005)
		<i>p</i> -Coumaric acid <sup>a</sup>	Apigenin	
		Caffeic acid <sup>a</sup>	Luteolin	
	FNS <sup>c</sup>	Naringenin chalcone	Apigenin	Ralston et al., (2005)
		Isoliquiritigenin	7,4'-Dihydroxyflavone	
	F3H <sup>c</sup>	Naringenin chalcone	Dihydroxykaempferol	
		Isoliquiritigenin	3,7,4'-Trihydroxyflavonol	
	IFS <sup>c</sup>	Naringenin chalcone	Genistein, 4'-tetrahydroxyisoflavanone	
		Isoliquiritigenin	Daizein and other isoflavones	
	IFS	Phenylalanine <sup>d</sup>	Genistein	Trantas et al., (2009)
	FLS and F3H	<i>p</i> -Coumaric acid <sup>a</sup>	Kaempferol	
	FLS, F3H, and F3'H	Naringenin	Quercetin	
	F3'H	Naringenin	Eriodictyol	Amor et al., (2010)
	F3H and FLS <sup>e</sup>	Glucose	Kaempferol	Duan et al., (2017)
	F3H, FLS, and F3'H/CPR <sup>b</sup>		Kaempferol Quercetin	Rodriguez et al., (2017)
FNS and F6H <sup>e</sup>		Scutellarein	Liu et al., (2018)	
F3H and FLS <sup>b</sup>		Kaempferol	Lyu et al., (2019)	
F3H, F3'H <sup>b</sup>		dihydromyricetin	Li et al., (2020)	
FNS and F6H <sup>e</sup>		Balclein	Liu et al., (2020)	
FNS and F8H <sup>e</sup>		Norwogonin		
<i>Y. lipolytica</i>	F3'H/CPR and F3H <sup>b</sup>	Glucose	Taxifolin	Lv et al., (2019)
<i>Pichia pastoris</i> ( <i>Komagataella phaffii</i> )	P450 from <i>Aspergillus oryzae</i> with CPR	Genistein	3'-Hydroxygenistein	Wang et al., (2016b)
<i>Rhodotorula glutinis</i>	Endogenous hydroxylase	Naringenin	Carthamidin, Isocarthamidin	Sordon et al., (2016)
		Apigenin	8-Hydroxyapigenin	
		Hesperetin	8-Hydroxyhesperetin	
		Luteolin	Hypoletin	
		Chrysin	Norwogonin	
<i>Glycosylation</i> <sup>f</sup>				
<i>E. coli</i>	O-glycosyltransferase (GT) (Glucose, 7-OH and 3'-OH)	Quercetin	Quercetin 7,3'-di-O-glucosides	Lim et al., (2004)
		Quercetin 7-O-glucosides		
		Quercetin 3'-O-glucosides		
	O-GT (Rhamnose, 3-OH)	Quercetin	Quercetin 3-O-rhamnosides	Kim et al., (2012)
		Kaempferol	Kaempferol 3-O-rhamnosides	
	O-GT (Glucose, 3-OH)	Quercetin	Quercetin 3-O-glucoside-7-O-rhamnoside	Kim et al., (2013)
	O-GT (Rhamnose, 7-OH)	Quercetin 3,7-O-bisrhamnoside		
O-GT (Rhamnose, 3-OH) O-GT (Rhamnose, 7-OH)				
O-GT (Rhamnose, 3'-OH)	Hesperetin	Hesperetin 3'-O-rhamnosides	Ruprecht et al., (2019)	

(Continued on following page)



**TABLE 3** | (Continued) Microbial production of diversified flavonoids.

Production host	Modification enzyme	Substrate	Product	Reference
<i>C. glutamicum</i> <i>L. lactis</i> <i>S. cerevisiae</i>	O-GT (Glucose, 3-OH) with F3H, DFR and ANS	Quercetin	Quercetin 3-O-rhamnosides	Yan et al., (2005a)
		Kaempferol	Kaempferol 3-O-rhamnosides	
		Naringenin	Cyanidin 3-O-glucoside	
	C-GT (Glucose, C6)	Eriodictyol	Pelargonidin 3-O-glucoside	Ito et al., (2014)
		2-Hydroxy-pinocembrin	2-Hydroxy-pinocembrin C-glucosides	
		2-Hydroxy-naringenin	2-Hydroxy-naringenin C-glucosides	
	O-GT (Glucose, 3-OH) with ANS	Phloretin	Nothofagin	Zha et al., (2018)
		Catechin	Cyanidin 3-O-glucoside	
	O-GT (Glucose, 3-OH) with ANS	Catechin	Cyanidin 3-O-glucoside	Solopova et al., (2019)
		Naringenin	Naringenin 7-O-glucosides	
<i>S. cerevisiae</i>	O-GT (Glucose, 7-OH and 4'-OH)	Naringenin 4'-O-glucosides	Werner and Morgan, (2009)	
		Phloretin		
		Naringenin		
	O-GT (Glucose, 7-OH)	Phleretin 2'-O-glucoside	Werner and Morgan, (2010)	
		Naringenin 7-O-glucosides		
	O-GT (Glucuronic acid, 7-OH) with FNS and F6H <sup>e</sup>	Naringenin 4'-O-glucosides	Wang et al., (2016a)	
		Scutellarein 7-O-glucoside		
	O-GT (Glucuronic acid, 7-OH) with FNS and F6H <sup>e</sup>	Apigenin 7-O-glucuronides	Liu et al., (2018)	
		Scutellarein 7-O-glucuronides		
	O-GT (Glucose, 3-OH) with F3H, F3'H, F3'5'H, DFR, LAR, and ANS <sup>d</sup>	Baicalin	Liu et al., (2020)	
Glucose		Pelargonidin 3-O-glucoside Cyanidin 3-O-glucoside Delphinidin 3-O-glucoside Kaempferol 3-O-glucoside Quercetin 3-O-glucoside Myricetin 3-O-glucoside Pelargonidin 3-O-glucoside	Eichenberger et al., (2018)	
O-GT (Glucose, 3-OH) with F3H, DFR, and ANS <sup>g</sup>	Naringenin	2-Hydroxynaringenin-C-glucoside	Brazier-Hicks and Edwards, (2013)	
		2-Hydroxynaringenin-C-glucoside		
		2-Hydroxynaringenin-C-glucoside		
C-GT (Glucose, C6) with flavanone 2-hydroxylase (F2H)	Glucose	Vitexin (8-C-glucoside)	Vanegas et al., (2018)	
		Isovitexin (8-C-glucoside)		
		Orientin (8-C-glucoside)		
C-GTs (Glucose, C6 and C8) with FNS, F3'H, and F2H <sup>d</sup>	Glucose	Isoorientin (6-C-glucoside)	Vanegas et al., (2018)	
		Isoorientin (6-C-glucoside)		
		Isoorientin (6-C-glucoside)		
Prenylation <i>S. cerevisiae</i>	SfN8DT-1	Naringenin	Sasaki et al., (2009)	
		<i>p</i> -Coumaric acid <sup>a</sup>		
	Fungal promiscuous prenyltransferase	Glucose <sup>g</sup>	8-Prenylnaringenin	Li et al., (2015)
Phenylalanine <sup>d</sup>		3'-Prenylnaringenin	Levisson et al., (2019) Isogai et al., (2021)	

<sup>a</sup>4CL, CHS, and CHI are co-expressed for flavanone synthesis from phenylpropanoic acids.

<sup>b</sup>TAL, 4CL, CHS, and CHI are co-expressed for naringenin synthesis.

<sup>c</sup>Co-expressed with CHI.

<sup>d</sup>PAL, C4H, CPR, 4CL, CHS, and CHI are co-expressed for naringenin synthesis.

<sup>e</sup>PAL, C4H, 4CL, CHS, and CHI are co-expressed for flavanone synthesis.

<sup>f</sup>Sugar donor and modification site of flavonoids for glycosyltransferase (GT) are indicated in parenthesis of the "Modification enzyme" column.

<sup>g</sup>PAL, C4H, CPR, 4CL, CHS, CHI, and TAL are co-expressed for naringenin synthesis.

## 2 FLAVANONES PRODUCTION IN MICROORGANISMS

Most structurally and functionally diverse flavonoids are biosynthesized from flavanones such as naringenin, pinocembrin and eriodictyol (Figure 1). Since naringenin is the most basic skeleton for flavanone, there have been many studies on reconstructing naringenin biosynthetic pathways in microorganisms. In this section, we introduce various aspects of microbial flavanone production, including metabolic engineering strategies and synthetic biology approaches, with a focus mainly on naringenin as well as some studies on pinocembrin and eriodictyol (Tables 1, 2).

### 2.1 Production Host

Bacteria and yeasts, especially *Escherichia coli* and *Saccharomyces cerevisiae*, have generally been used for the microbial production of flavonoids (Tables 1–3). Differences between these hosts significantly influence the strategy for reconstructing flavanone biosynthetic pathways. This section summarizes the characteristics of these host strains and examples of their development with respect to flavanone production.

#### 2.1.1 Bacteria

In general, bacteria grow faster than yeasts. In addition, heterogeneous genes can be expressed in a polycistronic manner in bacteria, decreasing the complexity of designing genetic systems. Furthermore, some bacteria represented by

*E. coli* have extraordinarily powerful expression systems for proteins, which often leads to the clear phenotype change by the introduced metabolic enzymes. In some cases, however, functional expression of eukaryotic enzymes is difficult in bacteria. For example, Watts et al. (2004) has reported to fail a functional expression of the plant C4H enzyme, a cytochrome P450 monooxygenase, in *E. coli*, resulting in the accumulation of *trans*-cinnamic acid. The lack of C4H enzyme significantly influenced the conversion efficiency of phenylalanine to *p*-coumaric acid when only PAL from the yeast *Rhodotorula mucilaginosa*, which catalyzes the deamination of phenylalanine and tyrosine, was expressed, resulting in low naringenin production compared with tyrosine supplementation (Hwang et al., 2003). Reconstruction of the naringenin biosynthetic pathway in *E. coli* thus utilized microbial TAL (usually a yeast enzyme), bacterial/plant 4CL, and plant CHS and CHI using tyrosine (Watts et al., 2004; Wu et al., 2014a; Wu et al., 2014b; Wu et al., 2015; Yang et al., 2018; Zhou et al., 2019; Zhou et al., 2020; Zhou et al., 2021). Expressing these heterogenous genes by engineering precursor metabolic pathways provided the highest levels of naringenin production in *E. coli* to date: 1,073 mg/L from tyrosine (Wu et al., 2021) and 588 mg/L from glucose (Zhou et al., 2020). As well as the naringenin production, expressing 4CL, CHS and CHI enzymes enabled pinocembrin production from *trans*-cinnamic acid (Leonard et al., 2007). An additional expression of PAL allowed to produce pinocembrin from glucose (Hwang et al., 2003; Wu et al., 2013; Kim et al., 2014; Cao et al., 2016a; Wu et al., 2016a; Cao et al., 2016b; Wu et al., 2016b) or phenylalanine (Dunstan et al., 2020). Similarly, the engineered *E. coli* produced eriodictyol from caffeic acid by expressing 4CL, CHS and CHI (Leonard et al., 2007; Fowler et al., 2009; Dunstan et al., 2020) or tyrosine by additional expression of TAL, F3'H and CPR with those pathway enzymes (Zhu et al., 2014).

Recently, several studies employed *Corynebacterium glutamicum* as the host strain instead of *E. coli* (Kallscheuer et al., 2016b; Milke et al., 2019). *C. glutamicum* is used for the several million ton-scale industrial production of amino acids such as glutamate and lysine (Koffas et al., 2003; Becker et al., 2011), and therefore might have a great potential to supply the precursor tyrosine through its endogenous pathway. Kallscheuer et al. (2016a) have previously found the potential of this organism to grow on several phenylpropanoids and identified the phenylpropanoid degradation pathway (Phd pathway), reporting an engineered *C. glutamicum* that produces 35 mg/L of naringenin from *p*-coumaric acid with deletion of the pathways for degradation and unspecific conversion of the phenylpropanoid substrates (Kallscheuer et al., 2016b). They also achieved 32 mg/L of naringenin from glucose followed by increasing the intracellular availability of precursors (tyrosine and malonyl-CoA) by additional engineering of the shikimate pathway and supplementation of cerulenin, an inhibitor of fatty acid synthesis (see Section 2.4) (Kallscheuer et al., 2016b). In a subsequent study by the same group, the authors modulated gene expressions of the fatty acid synthesis and the tricarboxylic acid (TCA) cycle of the host strain to enhance the malonyl-CoA supply without the supplementation of any

inhibitors, achieving 24 mg/L of naringenin from glucose (Milke et al., 2019).

Furthermore, some cyanobacteria species, such as *Synechocistis* and *Synechococcus*, have been used as production host for plant secondary metabolites including flavonoids and their pathway intermediates (Xue et al., 2014; Ni et al., 2016), achieving 7.1 mg/L of naringenin with cerulenin supplementation (Ni et al., 2016). Although further improvement in productivity and pathway design is required, cyanobacterium is an attractive host for flavonoid production owing to following features (Jin et al., 2021). They can use carbon dioxides (CO<sub>2</sub>) and sunlight as carbon and energy sources, respectively, contributing sustainable and low-cost microbial production of flavonoids. They can also supply large quantities of energies and reductants, ATP and NADPH, by photosynthetic system, facilitating the reaction steps of energy consumption (e.g., 4CL) and P450 enzymes (e.g., C4H, F3H and IFS) in the biosynthesis of flavonoids.

### 2.1.2 Yeasts

Leveraging the eukaryotic nature of yeasts facilitates the functional expression of plant-derived genes. For example, heterogeneous production of naringenin in *S. cerevisiae* has been achieved by introducing a five- or six-step pathway from phenylalanine (Trantas et al., 2009; Koopman et al., 2012) or a four-step pathway from tyrosine (Lyu et al., 2017; Rodriguez et al., 2017; Du et al., 2020). Furthermore, additional expression of microbial TAL in yeast harboring the naringenin biosynthetic pathway from phenylalanine improved the *p*-coumaric acid supply, leading to a 1.5-fold improvement in naringenin production (Koopman et al., 2012). To our knowledge, the highest reported production of naringenin from phenylalanine and glucose in *S. cerevisiae* is 8.9 mg/L (Trantas et al., 2009) and 1,129 mg/L (Zhang et al., 2021), respectively. Furthermore, pinocembrin production in *S. cerevisiae* was achieved by expressing PAL, 4CL and CHS (Jiang et al., 2005) and additional expression of CHI (Liu et al., 2020).

Recent research used the non-conventional yeast *Yarrowia lipolytica* as a host strain instead of *S. cerevisiae*. Although the genetic background of *Y. lipolytica* is less well understood in terms of gene manipulation than laboratory yeast, this yeast strain is expected to have superior malonyl-CoA cell availability, with the potential to enhance naringenin biosynthesis (Beopoulos et al., 2009). Indeed, the production of naringenin by *Y. lipolytica* was achieved by reconstructing the naringenin biosynthetic pathway and enhancing precursor supply (Lv et al., 2019; Palmer et al., 2020; Wei et al., 2020). The engineered *Y. lipolytica* generated 898 mg/L of naringenin from glucose (Palmer et al., 2020). Nonetheless, the introduction of several genes into yeasts to construct a multi-step pathway is laborious due to the monocistronic nature of their gene expression; thus, recent studies employed the four-step pathway from tyrosine, a simpler genetic design compared with that from phenylalanine (Lyu et al., 2017; Rodriguez et al., 2017; Du et al., 2020). To address this drawback, Zhang et al. reported a co-culture system of *S. cerevisiae* and *E. coli*, in which the former is engineered to efficiently produce naringenin from tyrosine and the latter is engineered to supply the precursor tyrosine from D-xylose to the former. This system resulted in the

**TABLE 4** | Enzyme list for flavanone biosynthesis.

Enzyme	Function	Source organism	Abbreviation
PAL	Phenylalanine ammonia lyase	<i>Populus trichocarpa</i> × <i>Populus deltoides</i> <i>Rhodotorula mucilaginosa</i> <i>Rhodotorula rubra</i> <i>Rhodospiridium toruloides</i>	Ptr-PdePAL RmPAL RrPAL RtPAL
TAL	Tyrosine ammonia lyase	<i>Bambusa oldhamii</i> <i>Flavobacterium johnsoniae</i> <i>Rhodobacter capsulatus</i> <i>Rhodotorula glutinis</i> <i>Rhodobacter sphaeroides</i> <i>Saccharothrix espanaensis</i> <i>Trichosporon cutaneum</i>	BoTAL FJTAL RcTAL RgTAL RsTAL SeTAL TcTAL
C4H	<i>trans</i> -Cinnamate 4-hydroxylase	<i>Arabidopsis thaliana</i>	AtC4H
C3H	<i>p</i> -Coumarate 3-hydroxylase	<i>Saccharothrix espanaensis</i>	SeC3H
CPR	Cytochrome P450 reductase	<i>Populus trichocarpa</i> × <i>Populus deltoides</i> <i>Catharanthus roseus</i>	Ptr-PdeCPR CrCPR
4CL	4-Coumaroyl-CoA ligase	<i>Glycine max</i> <i>Nicotiana tabacum</i> <i>Oryza sativa</i> <i>Petroselinum crispum</i> <i>Streptomyces coelicolor</i> A3(2)	Gm4CL Nt4CL Os4CL Pc4CL Sc4CL
CHS	Chalcone synthase	<i>Arabidopsis thaliana</i> <i>Glycyrrhiza echinate</i> <i>Glycyrrhiza uralensis</i> <i>Huperiza serrata</i> <i>Hypericum androsaemum</i> <i>Petunia hybrida</i> <i>Populus euramericana</i>	AtCHS GeCHS GuCHS HsCHS HaCHS PhCHS PeCHS
CHI	Chalcone isomerase	<i>Cattleya maxima</i> <i>Glycine max</i> <i>Medicago sativa</i>	CmCHS GmCHS MsCHS
F3'H 3OMT	Flavonoid 3'-hydroxylase 3-O-methyltransferase	<i>Gerbera hybrida</i> <i>Populus kitakamiensis</i>	GmF3'H Pk3OMT

production of 21.16 mg/L naringenin (Zhang et al., 2017). More recently, a co-culture system of two different *S. cerevisiae* strains was designed to efficiently produce naringenin derivatives (Du et al., 2020) (Table 2).

In addition to the above-mentioned bacteria and yeasts, the bacteria *Lactococcus lactis* (Solopova et al., 2019) and *Streptomyces albus* (Marin et al., 2018) and the non-conventional yeast *Pichia pastoris* (*Komagataella phaffii*) (Wang et al., 2016) have recently been used as alternative hosts for flavonoid production, but further engineering is necessary to improve the productivity and design pathways (Tables 1, 3).


## 2.2 Heterologous Genes for Flavanone Biosynthesis

Various plants produce flavonoids as secondary metabolites with different productivities and diversities. It is thus likely that genes from high-producing organisms encode pathway enzymes (Figure 2) with high catalytic activity. However, genes from a limited number of plants have been tested to date, as listed in Table 4 due to a lack of genetic information on these natural producers. Because of this situation, *Arabidopsis thaliana*, one of the well-characterized plant species, has been widely used as the source organism for naringenin and flavonoid biosynthesis genes despite of its lower productivity. Recently, Mark et al. (2019)

demonstrated that using 4CL and CHS from different plants such as *Medicago truncatula* and *Vitis vinifera* instead of *A. thaliana* resulted in the increased production of naringenin. In addition, expressing PAL from *Bambusa oldhamii* and 4CL from *Petroselinum crispum* instead of those from *Rhodotorula mucilaginosa* and *Streptomyces coelicolor* A3(2) increased intracellular *trans*-cinnamic acid, resulting in a 3.5-fold increase in pinocembrin production (Cao et al., 2016a). Similarly, Dunstan et al. (2020) evaluated the different combinations of three TAL, five 4CL and two CHS genes from various plants, from which they identified the pathway constructs that enabled efficient flavanone production in *E. coli*; 484 mg/L of naringenin and 17 mg/L of homoeriodictyol from glycerol, 198 mg/L of pinocembrin from phenylalanine and 88 mg/L of eriodictyol from caffeic acid with cerulenin supplementation, respectively. Taken together, reconstruction of the flavanone biosynthetic pathways with different combinations of pathway genes screened from various plants will be promising for increasing flavanone production by engineered microorganisms.

## 2.3 Biosynthesis From Pathway Intermediates

Supplementation of precursor aromatic amino acids and their derivative phenylpropanoids (e.g., tyrosine and *p*-coumaric acid)



**FIGURE 3 |** Metabolic engineering of endogenous biosynthetic pathways for precursors. **(A,B)** Biosynthesis of aromatic amino acids in *E. coli* **(A)** and *S. cerevisiae* **(B)**. Phenylalanine and tyrosine are synthesized via the shikimate pathway. DAHP synthase and chorismate mutase, which catalyze the first and last step of the shikimate pathway, respectively, are subjected to feedback regulation by aromatic amino acids (indicated by gray-dashed lines). Chorismate is bypassed to tryptophan. Red and blue arrows indicate the enzymatic reactions catalyzed by DAHP synthase and chorismate mutase, respectively. **(A)** The transcriptional regulator TyrR represses the expression of genes involved in the biosynthesis of phenylalanine and tyrosine, such as *aroF* and *tyrA* (indicated by black-dashed lines). **(B)** The enzymatic activity of Aro7 is activated by tryptophan. Green arrow indicates degradation of phenylpyruvate and *p*-hydroxyphenylpyruvate, precursors of phenylalanine and tyrosine, respectively, to the corresponding alcohols by decarboxylases, Aro10, and Pdc5, and alcohol dehydrogenase (ADH). PEP: phosphoenolpyruvate, E4P: D-erythrose-4-phosphate, DAHP: 3-deoxy-D-arabino-heptulosonate 7-phosphate. **(C)** Biosynthesis of malonyl-CoA. Endogenous malonyl-CoA is synthesized from acetyl-CoA by ACC. MCS catalyzes the formation of malonyl-CoA from malonate, which is used as the bypass pathway in malonyl-CoA synthesis, in addition to the natural metabolic pathway. Acetyl-CoA is synthesized in the PDH complex from pyruvate and by ACS from acetate in *S. cerevisiae*. The Ack-Pta pathway from acetate is also used in addition to these biosynthetic pathways in *E. coli*. PanK catalyzes the phosphorylation of pantothenate to 4-phospho pantothenate, a precursor of CoA. Red arrows represent two key enzymatic reactions for directly increasing intracellular malonyl-CoA, which are catalyzed by ACC and MCS. Whereas, blue arrows indicate pathways for acetyl-CoA consumption by glycolysis, the TCA cycle, and fatty acid synthesis, and malonyl-CoA consumption by the synthesis of fatty acids and phospholipids. Cerulenin inhibits fatty acid synthesis as indicated by grey dotted line. Pyruvate, a precursor for acetyl-CoA, is catabolized to acetate, another precursor of acetyl-CoA, and ethanol via acetaldehyde. The reaction catalyzed by ADH is bi-directional; thus, overexpression of ADH that prefers to convert ethanol to acetaldehyde, such as Adh2 in *S. cerevisiae*, was used for increasing acetate availability. ACC, acetyl-CoA carboxylase; MCS, malonyl-CoA synthetase; PDH, pyruvate dehydrogenase; ACS, acetyl-CoA synthetase; Ack, acetate kinase; Pta, phosphate acetyltransferase; PDC, pyruvate decarboxylase; ADH, alcohol dehydrogenase; ALDH, acetaldehyde dehydrogenase; PanK, pantothenate kinase.

to the culture medium increases their intracellular availability in engineered microorganisms, improving flavanone production without modulating their biosynthetic pathways. For example, phenylalanine supplementation increased *p*-coumaric acid production in *S. cerevisiae*, enabling the production of 8.9 mg/L naringenin from phenylalanine (Trantas et al., 2009). Moreover, the addition of tyrosine and phenylalanine to the culture medium resulted in a five-fold increase in production of naringenin and pinocembrin in *E. coli*, respectively (Hwang et al., 2003). More recent research reported naringenin production using engineered *E. coli* expressing the genes responsible for naringenin biosynthesis from tyrosine (i.e., TAL, 4CL, CHS, and CHI) upon tyrosine supplementation, achieving 1,073 mg/L naringenin from tyrosine (Wu et al., 2014b; Wu et al., 2015; Wu et al., 2021). Furthermore, supplementation of *trans*-cinnamic acid or *p*-coumaric acid improved naringenin production in both *E. coli* (Leonard et al., 2007; Fowler et al., 2009; Xu et al., 2011) and *S. cerevisiae* (Yan et al., 2005b; Trantas et al., 2009). Similarly, supplementation of *trans*-cinnamic acid enabled pinocembrin production in *E. coli* expressing 4CL, CHS, and CHI (Leonard et al., 2007), while eriodictyol production in *E. coli* was achieved by addition of tyrosine (Zhu et al., 2014) or caffeic acid (Leonard et al., 2007; Fowler et al., 2009; Dunstan et al., 2020), respectively. The addition of these pathway intermediates can decrease the necessity of introducing the genes responsible for intermediate biosynthesis, therefore simplifying the expression system. However, the use of these intermediates often negatively influences the microbial production of flavanone due to the increase in precursor cost, the toxicity (e.g., acidic nature) of these compounds, and the low solubility of these intermediates.

## 2.4 Pathway Design for *De Novo* Synthesis From a Sole Carbon Source

*De novo* synthesis from a simple sole carbon source such as glucose is the most cost-effective approach for microbial flavanone production, but would lead to an insufficient supply of precursors (the aromatic amino acids phenylalanine and tyrosine, and malonyl-CoA) due to tight regulation of the biosynthetic pathways for these compounds in a host microorganism. In the following subsections, we summarize efforts to date to overcome these limitations by the metabolic engineering of biosynthetic pathways for precursors (Table 1) (Santos et al., 2011; Koopman et al., 2012; Wu et al., 2014b; Kim et al., 2014; Zhu et al., 2014; Cao et al., 2016a; Wu et al., 2016a; Cao et al., 2016b; Wu et al., 2016b; Rodriguez et al., 2017; Yang et al., 2018; Zhou et al., 2019; Du et al., 2020; Dunstan et al., 2020; Zhou et al., 2020; Zhou et al., 2021).

### 2.4.1 Biosynthesis of Aromatic Amino Acids

Phenylalanine and tyrosine are biosynthesized from phosphoenolpyruvate (PEP) and D-erythrose-4-phosphate (E4P) via the shikimate pathway (Figures 3A,B). 3-Deoxy-D-arabino-heptulosonate 7-phosphate (DAHP) synthase catalyzes the first step of the shikimate pathway by condensing

PEP and E4P to form DAHP. Next, DAHP is converted to chorismate through shikimate via two-step enzymatic reactions. Then, chorismate mutase catalyzes the rearrangement of chorismate to prephenate, which is the last step in the shikimate pathway. Finally, prephenate is converted to tyrosine and phenylalanine via dehydrogenation and transamination in parallel pathways. The biosynthesis of phenylalanine and tyrosine via chorismate (the shikimate pathway) is tightly regulated at both the biochemical and genetic levels. DAHP synthases are feedback-inhibited by phenylalanine, tyrosine, and tryptophan, whereas chorismate mutases are subjected to feedback inhibition by phenylalanine and tyrosine. In addition, the expression of these biosynthetic enzymes is regulated in response to these aromatic amino acids (Braus, 1991; Rodriguez et al., 2014). These regulations of the pathway enzymes inhibit the overproduction of phenylalanine and tyrosine in the cell, leading to insufficient supply of these compounds. Therefore, removing the negative regulation of aromatic amino acid biosynthesis is a promising approach for enhancing the endogenous biosynthesis of these precursors.

In *E. coli*, there are three isozymes of DAHP synthase (AroF, AroG, and AroH) and two isoforms of chorismate mutase (TyrA, and PheA) (Figure 3A). *tyrA* and *pheA* encode the bifunctional enzyme that catalyzes the conversion of chorismate to prephenate (chorismate mutase) and the dehydrogenation of prephenate (prephenate dehydrogenase). AroF, AroG, and AroH are feedback-inhibited by tyrosine, phenylalanine and tryptophan, respectively, whereas the enzymatic activities of TyrA and PheA are subjected to feedback inhibition by tyrosine and phenylalanine, respectively. Previous studies described AroG and TyrA mutants that are insensitive to such feedback inhibition (Kikuchi et al., 1997; Lutke-Eversloh and Stephanopoulos, 2005). Since AroG contributes 80% of total cellular DAHP synthase activity, the expression of these AroG and TyrA mutants (AroG<sup>D146N</sup> and TyrA<sup>M531/A354V</sup>) increases intracellular tyrosine in *E. coli* even in the presence of tyrosine-insensitive AroF (Lütke-Eversloh and Stephanopoulos, 2007). Introducing the *aroG* and *tyrA* mutants (AroG<sup>D146N</sup>, TyrA<sup>M531/A354V</sup>) resulted in increased tyrosine supply in *E. coli*, enabling the production of over 80 mg/L of naringenin from glucose (Santos et al., 2011; Wu et al., 2014b; Yang et al., 2018). Because tyrosine biosynthesis in *E. coli* is transcriptionally regulated by TyrR (Pittard et al., 2005), the removal of downregulation by TyrR also increases intracellular tyrosine (Kim et al., 2018), contributing to the increase in naringenin production (Yang et al., 2018). Furthermore, Dunstan et al. (2020) demonstrated that deletion of *pheA* together with the introduction of *ppsA* that catalyzes the generation of PEP from pyruvate enabled 484 mg/L of naringenin production from glucose in combination with above mentioned strategies (i.e., the introduction of *aroG* and *tyrA* mutants and the knockout of *tyrR*). In contrast, the similar approach was employed to increase intracellular phenylalanine availability for heterologous pinocembrin production from phenylalanine via *trans*-cinnamic acid. Previous study reported that co-expression of PheA mutant (PheA<sup>T326P</sup>) that is insensitive to feedback inhibition by phenylalanine with the wild-type AroF

increased intracellular phenylalanine pool (Zhou et al., 2010). Introducing the wild-type *aroF* and the *pheA* mutant resulted in 40 mg/L of pinocembrin production from glucose (Wu et al., 2013), subsequently achieving 432 mg/L along with optimization of PAL expression level by modifying its mRNA secondary structure (Wu et al., 2016a).

In *S. cerevisiae*, two DAHP synthases, Aro3 and Aro4, are subjected to feedback inhibition by phenylalanine and tyrosine, respectively, while one chorismate mutase, Aro7, is inhibited by tyrosine but activated by tryptophan (**Figure 3B**) (Schnappauf et al., 1997; Helmstaedt et al., 2005). Koopman et al. (2012) deleted the *ARO3* gene and introduced the tyrosine-insensitive *ARO4* mutant (Aro4<sup>G226S</sup>) (Hartmann et al., 2003) into engineered *S. cerevisiae* to enhance the flux from glucose to phenylalanine. They also deleted genes encoding decarboxylase (Aro10, Pdc5, and Pdc6) to alleviate the loss of phenylpyruvate, a precursor of phenylalanine. Next, they introduced both PAL and TAL to the resultant strain to form *p*-coumaric acid from phenylalanine and tyrosine, enabling the production of 113 mg/L of naringenin from glucose. Later, tyrosine-insensitive Aro4 and/or Aro7 mutants (Aro4<sup>K229L</sup> and Aro7<sup>G141S</sup>) were used to increase naringenin production together with the heterologous expression of TAL in *S. cerevisiae* (Lyu et al., 2017; Rodriguez et al., 2017; Du et al., 2020). This strategy would improve the production of chorismate and prephenate; however, these compounds are also converted to tryptophan (by Trp2/Trp3) and phenylalanine (by Pha2), respectively, resulting in the loss of tyrosine. Lyu et al. (2017) revealed that weakening *PHA2* expression resulted in a 20% improvement in naringenin production due to enhanced tyrosine availability. Additionally, *p*-hydroxy-phenylpyruvate, a precursor of tyrosine, is degraded by endogenous decarboxylases, such as Aro10 and Pdc5. Deletion of these decarboxylases aided the accumulation of intracellular tyrosine (Rodriguez et al., 2015; Rodriguez et al., 2017), resulting in an approximately two-fold improvement in naringenin production (Lyu et al., 2017; Du et al., 2020).

#### 2.4.2 Biosynthesis of Malonyl-CoA

The metabolic engineering of pathways responsible for malonyl-CoA biosynthesis is another effective strategy for enhancing naringenin production (**Figure 3C**). Endogenous synthesis of malonyl-CoA depends on the conversion from acetyl-CoA by acetyl-CoA carboxylase (ACC) using adenosine triphosphate (ATP) (Milke and Marienhagen, 2020). Additional expression (overexpression) of ACC is therefore the most common way to enhance the intracellular malonyl-CoA pool and thus increase productivities of naringenin (Leonard et al., 2007; Xu et al., 2011; Du et al., 2020; Wu et al., 2021), pinocembrin (Leonard et al., 2007; Kim et al., 2014; Cao et al., 2016a), and eriodictyol (Leonard et al., 2007; Zhu et al., 2014). Another strategy is the addition of a synthetic bypass pathway for malonyl-CoA synthesis. For example, malonyl-CoA synthetase (MCS) encoded by *matB* can directly synthesize malonyl-CoA from malonate (An and Kim, 1998). Co-expression of MatB and the malonate carrier protein MatC in engineered *E. coli* would enhance intracellular malonyl-CoA by supplementing malonate (Wu et al., 2014a; Wu

et al., 2014b), resulting in between a 1.6–2.7-fold increase in naringenin production (Leonard et al., 2008; Santos et al., 2011). Similarly, expression of bypass pathway from malonate was employed to increase malonyl-CoA supply for pinocembrin production (Wu et al., 2013; Wu et al., 2016a). Further improvement of the intracellular malonyl-CoA pool can be accomplished by increasing acetyl-CoA availability and deleting the branching pathway from malonyl-CoA to fatty acids and phospholipids. In *S. cerevisiae*, acetyl-CoA is synthesized by the pyruvate dehydrogenase (PDH) complex from pyruvate and by acetyl-CoA synthetase (ACS) from acetate using ATP, whereas acetate kinase-phosphate acetyltransferase catalyzes the formation of acetyl-CoA in addition to PDH and ACS pathways in *E. coli*. Therefore, the overexpression of these enzymes can increase acetyl-CoA availability. For example, PDH (Xu et al., 2011), ACS (Leonard et al., 2007; Fowler et al., 2009), and ACC and acetate kinase-phosphate acetyltransferase (Ack-Pta) (Leonard et al., 2007) were overexpressed in engineered *E. coli*. More recently, Zhu et al. (2014) overexpressed the genes encoding endogenous ACS (*acs*) and heterologous ACC (*accBC* and *dtsR1* from *C. glutamicum*), and removed the acetate competition pathway by deletion of acetate kinase gene (*ackA*), achieving 107 mg/L of eriodictyol production from tyrosine in *E. coli* (Zhu et al., 2014). Similarly, a heterologous ACS mutant (Acs<sup>L641P</sup> from *Salmonella enteric*) was co-expressed with endogenous acetaldehyde dehydrogenase (Ald6) to bypass acetyl-CoA synthesis from PDH pathway, producing 2.9 mg/L of naringenin from *p*-coumaric acid in engineered *S. cerevisiae*. They also achieved 10.5 mg/L of naringenin followed by increasing CoA availability by additional expression of endogenous pantothenate kinase (PanK encoded by *CAB1*). Pantothenate supplementation to this yeast led to another 19% increase in naringenin production (12.5 mg/L) (Liu et al., 2017). In the subsequent studies, Adh2, an enzyme converting ethanol to acetaldehyde, or Adh2 and ACC were additionally expressed to further increase malonyl-CoA availability in engineered *S. cerevisiae* which produce derivatives of naringenin and pinocembrin (see **Section 3.1** and **3.2**) (Liu et al., 2018; Liu et al., 2020). Additionally, the downregulation of acetyl-CoA and pyruvate consuming pathways enhanced acetyl-CoA availability by, for example, deleting the genes involved in the TCA cycle and ethanol synthesis in engineered *E. coli* (Fowler et al., 2009) and deleting or downregulating ergosterol biosynthetic enzymes in engineered *S. cerevisiae* (Du et al., 2020). Furthermore, downregulation of the TCA cycle combined with the upregulation of glycolysis and malonyl-CoA synthesis achieved a four-fold increase in intracellular malonyl-CoA, enabling the production of 474 mg/L of naringenin by engineered *E. coli* (Xu et al., 2011). Interception of malonyl-CoA from fatty acid synthesis was also reported to increase malonyl-CoA availability in cells. Transacylases that catalyze the condensation of acyl-CoA (acetyl-CoA and malonyl-CoA) with acyl carrier protein were downregulated by the inhibitor cerulenin (Leonard et al., 2008; Cao et al., 2016b; Dunstan et al., 2020), antisense RNA (Wu et al., 2014a; Yang et al., 2015) and knockdown of the regulator (Yang et al., 2018) in engineered

*E. coli*. This strategy was also adopted for optimizing malonyl-CoA allocation in *Y. lipolytica* using CRISPRi-based genetic circuits (see **Section 2.6**) (Lv et al., 2020). Furthermore, CRISPRi-based downregulation of the various pathways for malonyl-CoA consumption, TCA cycle, ethanol synthesis and fatty acid synthesis, increased intracellular malonyl-CoA supply in *E. coli*, achieving the production of naringenin up to 1,073 mg/L from tyrosine (Wu et al., 2015; Lv et al., 2020; Wu et al., 2021) and 526 mg/L of pinocembrin from glucose (Wu et al., 2016b). These engineering efforts resulted in a 1.14- to 24-fold increased production of naringenin, 4- to 12-fold of pinocembrin and 2- to 3-fold of eriodictyol compared with the parental strains of *E. coli*, *S. cerevisiae* and *Y. lipolytica*, respectively. In addition, these engineering strategies were employed to modulate intracellular availability in *C. glutamicum*, resulting in the production of 24 mg/L of naringenin by the overexpression of ACC and downregulation of the TCA cycle (Milke et al., 2019). More recently, overexpression of peroxisomal proliferation and free fatty-acid transport pathway protein (Pex11) increased acetyl-CoA availability via  $\beta$ -oxidation, enabling the production of 1,129 mg/l of naringenin in *S. cerevisiae* (Zhang et al., 2021). The similar approach enabled the production of 898 mg/l of naringenin from glucose in *Y. lipolytica* (Palmer et al., 2020).

## 2.5 Directed Evolution/Pathway Tuning

The expression of heterogenous genes such as flavonoid biosynthetic genes in microorganisms often utilizes an inducible or constitutive expression system, but the resulting high-level expression of heterogenous genes often causes metabolic flux imbalance, a reduction in cell growth, an accumulation of (toxic) intermediates, and the synthesis of undesired by-products, resulting in reduced carbon influx into flavonoid biosynthesis. To overcome this problem, fine-tuning of the expression level of pathway gene(s) and directed evolution of pathway enzyme(s) have been used to optimize the introduced biosynthetic pathway (Jones et al., 2015).

The expression of pathway genes can be efficiently balanced by constructing strains with different expression levels of each pathway gene, followed by screening for high naringenin production. Zhou et al. (2019) established a high-throughput method to evaluate naringenin production using ultraviolet spectrophotometry–fluorescence spectrophotometry. In this method, cells are incubated in the presence of aluminum ion ( $Al^{3+}$ ) to allow the binding of  $Al^{3+}$  to naringenin, and the resulting  $Al^{3+}$ -bound naringenin is measured using fluorescence excitation at 382 nm and emission at 505 nm and/or UV absorbance at 373 nm. This approach was used to fine-tune the pathway enzymes to optimize promoter strength for naringenin pathway genes in both *E. coli* (Zhou et al., 2019) and *S. cerevisiae* (Gao et al., 2020). Another example utilized the naringenin-inducible transcription factor FdeR and its cognate binding site (*fdeO*) to construct a naringenin-responsive transcription switch in *S. cerevisiae* (Wang et al., 2019). This biosensor enabled the high-throughput screening of as many as 972 combinations of expression cassette with different promoter-terminator pairs and plasmid vectors for three naringenin

biosynthetic genes 4CL, CHS, and CHI, leading to identify the yeast strain producing 52.0 mg/L naringenin. Additionally, naringenin-responsive riboswitches have been developed (Jang et al., 2017; Xiu et al., 2017) and utilized for fine-tuning the expression of multiple enzymes in *E. coli*, enabling a 3-fold improvement in naringenin production (Hwang et al., 2021).

The high-throughput screening approaches using genetically encoded biosensors described above can be used not only to balance the expression of pathway enzymes, but also to perform directed evolution of the catalytic activity of these enzymes. To this end, several genetically encoded biosensors that translate intracellular concentrations of naringenin and malonyl-CoA into reporter gene expression levels have been described. For example, the transcription factor TtgR from *Pseudomonas putida* and its target promoter have been evolutionarily engineered in *E. coli* to respond to resveratrol, which is synthesized from *p*-coumaroyl-CoA by stilbene synthase (resveratrol synthase) catalysis (Xiong et al., 2017). The resultant biosensor was used to screen 4CL1 (*A. thaliana* 4CL) mutants with improved activity from a randomly mutagenized 4CL1 library, leading to the enhanced production of naringenin and resveratrol with *p*-coumaric acid supplementation. Furthermore, the repressor QdoR from *Bacillus subtilis* responsive for downstream flavonoids (flavonols; quercetin and kaempferol, see **Section 3.1** for more information on these compounds) can be used for the high-throughput screening of functional FLS mutants that convert dihydrokaempferol into kaempferol with different efficiencies (Siedler et al., 2014).

As mentioned above, biosensor-assisted metabolic engineering is a powerful technique to obtain high production of flavonoids, but this strategy requires well-optimized biosensors for each host species. Both rational and irrational engineering toolboxes have been described in detail for *E. coli* and *S. cerevisiae*, but not for non-conventional microbes such as *Y. lipolytica*. Recently, Bowman et al. (2021) addressed this problem by developing a droplet-based high-throughput screening technique that enables the co-encapsulation of two different microbes in a single droplet: one is a mutant cell that produces naringenin, and the other is a sensor cell that translates the naringenin concentration inside the droplet into an altered level of gene expression. Different flavonoids can be monitored by simply changing the sensor cells using this method.

## 2.6 Dynamic Regulation for Optimizing the Allocation of Precursors in Flavonoid Synthesis

Since flavonoids share the same precursors as lipids during biosynthesis, reallocation of the precursors could increase flavonoid production. To this end, metabolite sensors that respond to flavonoids or their precursors have been assembled into genetic circuits to dynamically control the expression of branching pathway enzymes. Lv et al. (2020) constructed CRISPRi-based, fatty acid-responsive genetic circuits in *Y. lipolytica* to improve naringenin production by optimizing malonyl-CoA allocation. They used a synthetic, fatty acid inducible promoter, based on the regulatory components from

natural fatty acids inducible promoter, *POX2* (Shabbir Hussain et al., 2017), to regulate the expression of three gRNAs that repress three genes responsible for fatty acids synthesis [*FAS1*, *FAS2*, and *YAL10E18590g* (equivalent to bacterial *fabD* or yeast *MCT1*)], together with constitutively expressed dCas9. This approach repressed fatty acid synthesis in response to excessive fatty acid synthesis, enabling an increase in naringenin production by up to 74.8%. The expression of pathway genes can be regulated in response not only to intracellular metabolite concentrations but also to cell density. For example, Dinh and Prather created an autonomous genetic circuit that leads to the activation of TAL and 4CL gene expression along with the CRISPRi-based downregulation of malonyl-CoA consuming pathway genes (*fabF* and *sucC*) in response to high cell density in *E. coli* (Dinh and Prather, 2019). This dynamic regulation resulted in a 6-fold increase in naringenin production compared with the control strain without the dynamic control and the CRISPRi-based downregulation even in a bioreactor fermentation.

In addition to pathway gene regulation, metabolite sensors can be used for the selective growth of cells producing target metabolites to increase productivity. Lv et al. (2020) created a naringenin-responsive transcription switch based on the transcription factor FdeR and its binding sites (*fdeO*) fused to the *TEF1* promoter, without an upstream activation sequence controlling the *LEU2* gene. In this system, only cells producing sufficient naringenin to induce *LEU2* expression can grow in medium lacking leucine. This strategy, together with the aforementioned fatty-acid responsive CRISPRi system, enabled stable naringenin production by up to 324 generations.

Despite the wide utility of metabolite sensors described above, the sensors are often saturated by the metabolite to be sensed. In naringenin production, the FdeR-based naringenin sensor could not detect naringenin over 40 mg/L. Boada et al. (2020) addressed this problem by indirectly detecting large amounts of naringenin by converting a small fraction of the naringenin into kaempferol (see Section 3.1 for more information on this compound), which can be detected using the kaempferol-responsive transcription factor QdoR.

### 3 MICROBIAL PRODUCTION OF DIVERSIFIED FLAVONOIDS

In plants, structurally complicated flavonoids are synthesized using flavanones as starting compounds (Figure 1) (Winkel, 2006). Introducing plant-derived genes encoding modification enzymes into microbes allows microbial production of these complicated flavonoids. This section introduces the microbial production of structurally complicated flavonoids derived from naringenin, pinocembrin and eriodictyol using various types of modifications (Table 3).

#### 3.1 Hydroxylation/Dehydration

Flavone, dihydroflavonol, flavonol, anthocyanidin, flavane-3-ol, and isoflavone can be synthesized *via* flavanone (Figure 1). This has prompted many studies on the bioconversion of naringenin

and eriodictyol by the expression of one or several modification enzymes or the partial reconstruction of the biosynthetic pathway in *E. coli* (Yan et al., 2005a; Leonard et al., 2006; Zhao et al., 2015) and *S. cerevisiae* (Leonard et al., 2005; Ralston et al., 2005; Trantas et al., 2009; Amor et al., 2010). In *E. coli*, kaempferol, a flavonol, was produced by expressing FLS (FLS1) from *A. thaliana* to convert dihydrokaempferol, a dihydroflavonol. The concentration of kaempferol produced was monitored by a QdoR-based metabolite sensor responsive to kaempferol and quercetin (flavonols) but not to dihydrokaempferol (see Section 2.5) (Siedler et al., 2014). Furthermore, kaempferol was produced in *E. coli* from tyrosine by expressing F3H and FLS in addition to the naringenin biosynthetic pathway *via* feedback control of CHS expression (downregulation) using a genetic circuit incorporating the QdoR-based kaempferol biosensor (see Section 2.6) (Boada et al., 2020). Recently, the production of these derivatives using bacteria and yeasts other than *E. coli* and *S. cerevisiae* has been reported: kaempferol and quercetin production from phenylpropanoic acids by *C. glutamicum* expressing F3H and FLS together with 4CL, CHS and CHI (Kallscheuer et al., 2017), hydroxylation of genistein (an isoflavone) by *P. pastoris* (Wang et al., 2016), and conversion of catechin (a flavane-3-ol) to cyanidin (a anthocyanidin) by *L. lactis* expressing ANS (Solopova et al., 2019). This engineered *L. lactis* further glycosylated cyanidin to anthocyanin (see Section 3.2).

The *de novo* synthesis of various naringenin derivatives, including kaempferol and quercetin, was recently achieved by expressing suitable modification enzymes in *S. cerevisiae* carrying the reconstructed naringenin biosynthetic pathway, namely, the production of kaempferol (Duan et al., 2017; Rodriguez et al., 2017; Lyu et al., 2019) and quercetin (Rodriguez et al., 2017) by expression of FLS, and dihydromyricetin (a dihydroflavonol) by multiple hydroxylases, including F3H (Li et al., 2020). In addition, Liu et al. (2018) identified P450 enzymes responsible for hydroxylation of C6 (F6H) and C8 (F8H) of flavanone backbone. Expression of these hydroxylases with FNS achieved the production of hydroxylated flavones from glucose in engineered *S. cerevisiae*: expression of F6H, FNS and naringenin biosynthetic enzymes produced scutellarein, whereas that of F6H or F8H with FNS and genes responsible for pinocembrin biosynthesis enabled the production of balclein and norwogonin respectively (Liu et al., 2020). The engineered *S. cerevisiae* further glycosylated these flavones by additional expression of glycosyltransferase (see section 3.2). Furthermore, engineered *S. albus* and *Y. lipolytica de novo* synthesized flavonols (myricetin, kaempferol and quercetin) (Marin et al., 2018) and taxifolin (a dihydroflavonol) (Lv et al., 2019), respectively. More recently, the co-culture system of two different *S. cerevisiae* strains has been developed, in which one strain synthesized naringenin and the other converted it to its derivatives (Table 2) (Du et al., 2020). This platform enabled the production of different flavonols and anthocyanidins simply by exchanging naringenin-converting strain with that expressing different modification enzymes. Another approach for the hydroxylation of flavonoids involves the bioconversion of flavonoids using yeast or *Streptomyces* bacteria without the use of heterogeneous genes (Sordon et al., 2016; Shrestha et al., 2021).



### 3.2 Glycosylation

Several studies reported the bioconversion of flavonoid scaffolds to *O*-glycosylated derivatives using *E. coli* (Lim et al., 2004; Kim et al., 2012; Kim et al., 2013; Ruprecht et al., 2019) and *S. cerevisiae* (Werner and Morgan, 2009, 2010; Wang et al., 2016) by expressing botanical *O*-glycosyltransferase (Figure 1). *E. coli* (Ito et al., 2014) and *S. cerevisiae* (Brazier-Hicks and Edwards, 2013) expressing *C*-glycosyltransferase converted flavone and dihydrochalcone to the corresponding *C*-glycosylated derivatives. Moreover, anthocyanins, which are pigments synthesized via the glycosylation of anthocyanidins, were biosynthesized from flavanone (eriodictyol and naringenin) in engineered *E. coli* expressing *O*-glycosyltransferase with F3H, DFR, ANS (Yan et al., 2005a), and from catechin by *C. glutamicum* (Zha et al., 2018) and *L. lactis* (Solopova et al., 2019) expressing *O*-glycosyltransferase and ANS.

More recent studies reported the *de novo* synthesis of anthocyanins by the expression of *O*-glycosyltransferase, F3H, DFR, and ANS with reconstruction of the naringenin biosynthetic pathway in engineered *S. cerevisiae* (Eichenberger et al., 2018; Levisson et al., 2018). Additionally, a co-culture system comprising four different *E. coli* strains was developed for the *de novo* synthesis of anthocyanin. Each *E. coli* strain was designed to provide one of the following modules: synthesis of phenylpropanoic acids from glucose, formation of flavanones, conversion of flavanones to flavane-3-ols, and production of anthocyanins from flavane-3-ols (Table 2) (Jones et al., 2017). Furthermore, the *de novo* synthesis of flavonoid-7-*O*-glucuronides, such as apigenin-7-*O*-glucuronide, scutellarin and baicalin, were reported by the expression of 7-*O*-glucuronosyltransferase with FNS in engineered *S. cerevisiae* that produce naringenin (Liu et al., 2018) or pinocembrin (Liu et al., 2020). Similarly, *C*-glycosylated flavones were *de novo* synthesized using *C*-glycosyltransferase together with the reconstruction of the naringenin biosynthetic pathway in yeast (Vanegas et al., 2018).

### 3.3 Prenylation

Flavonoid prenyltransferases, a class of enzymes responsible for flavonoid prenylation, are membrane-bound and strictly control the stereoselectivity of prenyl transfer (Figure 1). The prenyltransferase SfN8DT-1, isolated from *Sophora flavescens*, is responsible for the C8-prenylation of naringenin. Expression of this enzyme in *S. cerevisiae* resulted in the bioconversion of naringenin to 8-prenylnaringenin (Sasaki et al., 2009). Moreover, SfN8DT expression in naringenin-producing *S. cerevisiae* resulted in the synthesis of 8-prenylnaringenin from glucose (Levisson et al., 2019), phenylalanine (Isogai et al., 2021) and *p*-coumaric acid (Li et al., 2015).

However, to date only 13 flavonoid prenyltransferases have been identified from plants, limiting the microbial production of prenylated flavonoids. For example, three kinds of prenylnaringenins have been isolated from plants: 8-, 6-, and 3'-prenylnaringenin carrying the dimethylallyl moiety at C8 or C6 in the A-ring, and at C3' in the B-ring, respectively. However, only the prenyltransferase SfN8DT-1 was identified, and enzymes responsible for the C6- and C3'-prenylation of naringenin remain

unidentified. Recently, the *in vivo* production of 3'-prenylnaringenin was achieved by the expression of fungal prenyltransferase with broad substrate specificity (Isogai et al., 2021). Such microbial promiscuous enzyme may also be able to catalyze prenylation of other positions of flavonoids, for which specific enzymes have not yet been identified.

## 4 CONCLUSION AND FUTURE PERSPECTIVES

Flavonoids are promising as pharmaceuticals and food supplements due to their various biological activities; however, their supply is not stable or environmentally friendly. Therefore, the development of microbial production platform for flavonoids is desired. In this review, we focused on the strategies to increase microbial production of flavanones, specifically naringenin, pinocembrin, and eriodictyol, that serve as a starting compound for diverse flavonoids. The strategies are mainly the pathway engineering for the biosynthesis of flavanones and their precursors (aromatic amino acids and malonyl-CoA) by selecting host organism, gene source and optimizing enzyme expression levels.

As described in Section 2.6, several recent studies have demonstrated that genetically encoded biosensors for flavonoids can be useful for improving naringenin production by dynamically controlling the expression of pathway enzymes in response to cellular conditions. However, such strategies largely depend on the specifications of the biosensors, thereby necessitating the reconstruction of biosensors for functional fine-tuning or improvement. To this end, both rational and evolutionary methodologies for engineering synthetic promoters (Snoek et al., 2020; Cazier and Blazeck, 2021; Tominaga et al., 2021) can be adapted. The assembly of well-optimized biosensors into genetic circuits to control the important pathway enzymes will further sophisticate the flavonoids production as well as balancing their expression levels *via* translational control using a ribosome binding site (RBS) library (Wang et al., 2021).

We also described recent progress in the microbial production of structurally and functionally diverse flavonoids *via* flavanones such as naringenin, some of which have complex structures. Diverse flavonoids can be efficiently produced by introducing modification enzymes into flavanone-producing microbes. Although thousands of flavonoids are known to be produced in plants, there are few reports describing the microbial production of such flavonoids, and these were summarized in this review. Progress faces two possible bottlenecks: only a few modification enzymes from plants have been identified, and these enzymes are membrane-bound, complicating their functional expression in bacteria. Methodologies for the rapid identification and microbial expression of these enzymes need to be developed. The massive increase in genomic information due to the expansion of next-generation sequencing will help to uncover novel microbial enzymes, as well as botanical enzymes suitable for various modification steps (Unamba et al., 2015; Medema and Osbourn, 2016). Alternatively, microbial promiscuous enzymes can be used not only as modification enzymes but also as easy-to-express counterparts of botanical enzymes. Altogether, advanced

methodologies combining metabolic engineering and synthetic biology will enable microbial production of an incredibly diverse range of plant flavonoids.

## AUTHOR CONTRIBUTIONS

SI: conceptualization, investigation, original draft preparation, visualization, writing and editing, MT: conceptualization, investigation, original draft preparation, visualization, writing and editing, AK: conceptualization, supervision and JI: conceptualization, investigation, editing.

## REFERENCES

- Amor, I. L., Hehn, A., Guedone, E., Ghedira, K., Engasser, J. M., Chekir-Ghedira, L., et al. (2010). Biotransformation of Naringenin to Eriodictyol by *Saccharomyces Cerevisiae* Functionally Expressing Flavonoid 3' Hydroxylase. *Nat. Prod. Commun.* 5, 1893–1898. doi:10.1177/1934578x1000501211
- An, J. H., and Kim, Y. S. (1998). A Gene Cluster Encoding Malonyl-CoA Decarboxylase (MatA), Malonyl-CoA Synthetase (MatB) and a Putative Dicarboxylate Carrier Protein (MatC) in *Rhizobium Trifolii*. Cloning, Sequencing, and Expression of the Enzymes in *Escherichia coli*. *Eur. J. Biochem.* 257, 395–402. doi:10.1046/j.1432-1327.1998.2570395.x
- Barros, J., and Dixon, R. A. (2020). Plant Phenylalanine/Tyrosine Ammonia-Lyases. *Trends Plant Sci.* 25, 66–79. doi:10.1016/j.tplants.2019.09.011
- Becker, J., Zelder, O., Häfner, S., Schröder, H., and Wittmann, C. (2011). From Zero to Hero-Design-Based Systems Metabolic Engineering of *Corynebacterium Glutamicum* for L-Lysine Production. *Metab. Eng.* 13, 159–168. doi:10.1016/j.jymben.2011.01.003
- Beopoulos, A., Cescut, J., Haddouche, R., Uribelarrea, J.-L., Molina-Jouve, C., and Nicaud, J.-M. (2009). *Yarrowia Lipolytica* as a Model for Bio-Oil Production. *Prog. Lipid Res.* 48, 375–387. doi:10.1016/j.plipres.2009.08.005
- Boada, Y., Vignoni, A., Picó, J., and Carbonell, P. (2020). Extended Metabolic Biosensor Design for Dynamic Pathway Regulation of Cell Factories. *iScience* 23, 101305. doi:10.1016/j.isci.2020.101305
- Bowman, E. K., Wagner, J. M., Yuan, S. F., Deaner, M., Palmer, C. M., D'oeisnitz, S., et al. (2021). Sorting for Secreted Molecule Production Using a Biosensor-In-Microdroplet Approach. *Proc. Natl. Acad. Sci. U. S. A.* 118. doi:10.1073/pnas.2106818118
- Braus, G. H. (1991). Aromatic Amino Acid Biosynthesis in the Yeast *Saccharomyces cerevisiae*: a Model System for the Regulation of a Eukaryotic Biosynthetic Pathway. *Microbiol. Rev.* 55, 349–370. doi:10.1128/mr.55.3.349-370.1991
- Brazier-Hicks, M., and Edwards, R. (2013). Metabolic Engineering of the Flavone-C-Glycoside Pathway Using Polyprotein Technology. *Metab. Eng.* 16, 11–20. doi:10.1016/j.jymben.2012.11.004
- Cao, W., Ma, W., Wang, X., Zhang, B., Cao, X., Chen, K., et al. (2016a). Enhanced Pinocembrin Production in *Escherichia coli* by Regulating Cinnamic Acid Metabolism. *Sci. Rep.* 6, 32640. doi:10.1038/srep32640
- Cao, W., Ma, W., Zhang, B., Wang, X., Chen, K., Li, Y., et al. (2016b). Improved Pinocembrin Production in *Escherichia coli* by Engineering Fatty Acid Synthesis. *J. Ind. Microbiol. Biotechnol.* 43, 557–566. doi:10.1007/s10295-015-1725-3
- Cazier, A. P., and Blazeck, J. (2021). Advances in Promoter Engineering: Novel Applications and Predefined Transcriptional Control. *Biotechnol. J.* 16, e2100239. doi:10.1002/biot.202100239
- Dinh, C. V., and Prather, K. L. J. (2019). Development of an Autonomous and Bifunctional Quorum-sensing Circuit for Metabolic Flux Control in Engineered *Escherichia coli*. *Proc. Natl. Acad. Sci. U.S.A.* 116, 25562–25568. doi:10.1073/pnas.1911144116

## FUNDING

This work was partially supported by the Commission for Development of Artificial Gene Synthesis Technology for Creating Innovative Biomaterials from the Ministry of Economy, Trade and Industry (METI), the CREST program (JPMJCR21N2) and the JST-Mirai Program (JPMJMI17EJ) from the Japan Science and Technology Agency (JST), JSPS KAKENHI (Grant Number 18K14374), and the Intensive Support for Young Promising Researchers program (JPNP20004) from the New Energy and Industrial Technology Development Organization (NEDO).

- Dixon, R. A., and Pasinetti, G. M. (2010). Flavonoids and Isoflavonoids: from Plant Biology to Agriculture and Neuroscience. *Plant Physiol.* 154, 453–457. doi:10.1104/pp.110.161430
- Du, Y., Yang, B., Yi, Z., Hu, L., and Li, M. (2020). Engineering *Saccharomyces cerevisiae* Coculture Platform for the Production of Flavonoids. *J. Agric. Food Chem.* 68, 2146–2154. doi:10.1021/acs.jafc.9b07916
- Duan, L., Ding, W., Liu, X., Cheng, X., Cai, J., Hua, E., et al. (2017). Biosynthesis and Engineering of Kaempferol in *Saccharomyces cerevisiae*. *Microb. Cell. Fact.* 16, 165. doi:10.1186/s12934-017-0774-x
- Dunstan, M. S., Robinson, C. J., Jervis, A. J., Yan, C., Carbonell, P., Hollywood, K. A., et al. (2020). Engineering *Escherichia coli* towards De Novo Production of Gatekeeper (2S)-Flavanones: Naringenin, Pinocembrin, Eriodictyol and Homoeriodictyol. *Synth. Biol. (Oxf)* 5, ysaa012. doi:10.1093/synbio/ysaa012
- Eichenberger, M., Hansson, A., Fischer, D., Dürr, L., and Naesby, M. (2018). De Novo biosynthesis of Anthocyanins in *Saccharomyces cerevisiae*. *FEMS Yeast Res.* 18. doi:10.1093/femsyr/foy046
- Fowler, Z. L., Gikandi, W. W., and Koffas, M. A. G. (2009). Increased Malonyl Coenzyme A Biosynthesis by Tuning the *Escherichia coli* Metabolic Network and its Application to Flavanone Production. *Appl. Environ. Microbiol.* 75, 5831–5839. doi:10.1128/aem.00270-09
- Fraga, C. G., Croft, K. D., Kennedy, D. O., and Tomás-Barberán, F. A. (2019). The Effects of Polyphenols and Other Bioactives on Human Health. *Food Funct.* 10, 514–528. doi:10.1039/c8fo01997e
- Gao, S., Zhou, H., Zhou, J., and Chen, J. (2020). Promoter-Library-Based Pathway Optimization for Efficient (2S)-Naringenin Production from *P-Coumaric Acid* in *Saccharomyces cerevisiae*. *J. Agric. Food Chem.* 68, 6884–6891. doi:10.1021/acs.jafc.0c01130
- Harborne, J. B., and Williams, C. A. (2000). Advances in Flavonoid Research since 1992. *Phytochemistry* 55, 481–504. doi:10.1016/s0031-9422(00)00235-1
- Hartmann, M., Schneider, T. R., Pfeil, A., Heinrich, G., Lipscomb, W. N., and Braus, G. H. (2003). Evolution of Feedback-Inhibited  $\beta/\alpha$  Barrel Isoenzymes by Gene Duplication and a Single Mutation. *Proc. Natl. Acad. Sci. U.S.A.* 100, 862–867. doi:10.1073/pnas.0337566100
- Helmstaedt, K., Strittmatter, A., Lipscomb, W. N., and Braus, G. H. (2005). Evolution of 3-deoxy-D-Arabinose-7-Phosphate Synthase-Encoding Genes in the Yeast *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. U.S.A.* 102, 9784–9789. doi:10.1073/pnas.0504238102
- Hwang, E. I., Kaneko, M., Ohnishi, Y., and Horinouchi, S. (2003). Production of Plant-specific Flavanones by *Escherichia coli* Containing an Artificial Gene Cluster. *Appl. Environ. Microbiol.* 69, 2699–2706. doi:10.1128/aem.69.5.2699-2706.2003
- Hwang, H. G., Noh, M. H., Koffas, M. A. G., Jang, S., and Jung, G. Y. (2021). Multi-level Rebalancing of the Naringenin Pathway Using Riboswitch-Guided High-Throughput Screening. *Metab. Eng.* 67, 417–427. doi:10.1016/j.jymben.2021.08.003
- Isogai, S., Okahashi, N., Asama, R., Nakamura, T., Hasunuma, T., Matsuda, F., et al. (2021). Synthetic Production of Prenylated Naringenins in Yeast Using Promiscuous Microbial Prenyltransferases. *Metab. Eng. Commun.* 12, e00169. doi:10.1016/j.mec.2021.e00169
- Ito, T., Fujimoto, S., Shimosaka, M., and Taguchi, G. (2014). Production of C-Glucosides of Flavonoids and Related Compounds by *Escherichia coli*

- Expressing Buckwheat C-Glucosyltransferase. *Plant Biotechnol.* 31, 519–524. doi:10.5511/plantbiotechnology.14.1016a
- Jang, S., Jang, S., Xiu, Y., Kang, T. J., Lee, S.-H., Koffas, M. A. G., et al. (2017). Development of Artificial Riboswitches for Monitoring of Naringenin *In Vivo*. *ACS Synth. Biol.* 6, 2077–2085. doi:10.1021/acssynbio.7b00128
- Jiang, H., Wood, K. V., and Morgan, J. A. (2005). Metabolic Engineering of the Phenylpropanoid Pathway in *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* 71, 2962–2969. doi:10.1128/aem.71.6.2962-2969.2005
- Jin, H., Wang, Y., Zhao, P., Wang, L., Zhang, S., Meng, D., et al. (2021). Potential of Producing Flavonoids Using Cyanobacteria as a Sustainable Chassis. *J. Agric. Food Chem.* 69, 12385–12401. doi:10.1021/acs.jafc.1c04632
- Jones, J. A., Vernacchio, V. R., Collins, S. M., Shirke, A. N., Xiu, Y., Englaender, J. A., et al. (2017). Complete Biosynthesis of Anthocyanins Using *E. coli* Polycultures. *mBio* 8, e00621–00617. doi:10.1128/mBio.00621-17
- Jones, J. A., Toparlak, Ö. D., and Koffas, M. A. (2015). Metabolic Pathway Balancing and its Role in the Production of Biofuels and Chemicals. *Curr. Opin. Biotechnol.* 33, 52–59. doi:10.1016/j.copbio.2014.11.013
- Kallscheuer, N., Vogt, M., Bott, M., and Marienhagen, J. (2017). Functional Expression of Plant-Derived O-methyltransferase, Flavanone 3-hydroxylase, and Flavonol Synthase in *Corynebacterium glutamicum* for Production of Pterostilbene, Kaempferol, and Quercetin. *J. Biotechnol.* 258, 190–196. doi:10.1016/j.jbiotec.2017.01.006
- Kallscheuer, N., Vogt, M., Kappelmann, J., Krumbach, K., Noack, S., Bott, M., et al. (2016a). Identification of the Phd Gene Cluster Responsible for Phenylpropanoid Utilization in *Corynebacterium glutamicum*. *Appl. Microbiol. Biotechnol.* 100, 1871–1881. doi:10.1007/s00253-015-7165-1
- Kallscheuer, N., Vogt, M., Stenzel, A., Gätgens, J., Bott, M., and Marienhagen, J. (2016b). Construction of a *Corynebacterium glutamicum* Platform Strain for the Production of Stilbenes and (2S)-Flavanones. *Metab. Eng.* 38, 47–55. doi:10.1016/j.ymben.2016.06.003
- Kikuchi, Y., Tsujimoto, K., and Kurahashi, O. (1997). Mutational Analysis of the Feedback Sites of Phenylalanine-Sensitive 3-Deoxy-D-Arabinose-Heptulosonate-7-Phosphate Synthase of *Escherichia coli*. *Appl. Environ. Microbiol.* 63, 761–762. doi:10.1128/aem.63.2.761-762.1997
- Kim, B.-G., Kim, H. J., and Ahn, J.-H. (2012). Production of Bioactive Flavonol Rhamnosides by Expression of Plant Genes in *Escherichia coli*. *J. Agric. Food Chem.* 60, 11143–11148. doi:10.1021/jf302123c
- Kim, B., Binkley, R., Kim, H. U., and Lee, S. Y. (2018). Metabolic Engineering of *Escherichia coli* for the Enhanced Production of L-tyrosine. *Biotechnol. Bioeng.* 115, 2554–2564. doi:10.1002/bit.26797
- Kim, B. G., Lee, H., and Ahn, J.-H. (2014). Biosynthesis of Pinocembrin from Glucose Using Engineered *Escherichia coli*. *J. Microbiol. Biotechnol.* 24, 1536–1541. doi:10.4014/jmb.1406.06011
- Kim, H. J., Kim, B.-G., and Ahn, J.-H. (2013). Regioselective Synthesis of Flavonoid Bisglycosides Using *Escherichia coli* Harboring Two Glycosyltransferases. *Appl. Microbiol. Biotechnol.* 97, 5275–5282. doi:10.1007/s00253-013-4844-7
- Koffas, M. A. G., Jung, G. Y., and Stephanopoulos, G. (2003). Engineering Metabolism and Product Formation in *Corynebacterium glutamicum* by Coordinated Gene Overexpression. *Metab. Eng.* 5, 32–41. doi:10.1016/s1096-7176(03)00002-8
- Koopman, F., Beekwilder, J., Crimi, B., Van Houwelingen, A., Hall, R. D., Bosch, D., et al. (2012). De Novo production of the Flavonoid Naringenin in Engineered *Saccharomyces cerevisiae*. *Microb. Cell. Fact.* 11, 155. doi:10.1186/1475-2859-11-155
- Leonard, E., Lim, K.-H., Saw, P.-N., and Koffas, M. A. G. (2007). Engineering Central Metabolic Pathways for High-Level Flavonoid Production in *Escherichia coli*. *Appl. Environ. Microbiol.* 73, 3877–3886. doi:10.1128/aem.00200-07
- Leonard, E., Yan, Y., Fowler, Z. L., Li, Z., Lim, C.-G., Lim, K.-H., et al. (2008). Strain Improvement of Recombinant *Escherichia coli* for Efficient Production of Plant Flavonoids. *Mol. Pharm.* 5, 257–265. doi:10.1021/mp7001472
- Leonard, E., Yan, Y., and Koffas, M. (2006). Functional Expression of a P450 Flavonoid Hydroxylase for the Biosynthesis of Plant-specific Hydroxylated Flavonols in *Escherichia coli*. *Metab. Eng.* 8, 172–181. doi:10.1016/j.ymben.2005.11.001
- Leonard, E., Yan, Y., Lim, K. H., and Koffas, M. A. G. (2005). Investigation of Two Distinct Flavone Synthases for Plant-specific Flavone Biosynthesis in *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* 71, 8241–8248. doi:10.1128/aem.71.12.8241-8248.2005
- Levisson, M., Araya-Cloutier, C., De Bruijn, W. J. C., Van Der Heide, M., Salvador López, J. M., Daran, J.-M., et al. (2019). Toward Developing a Yeast Cell Factory for the Production of Prenylated Flavonoids. *J. Agric. Food Chem.* 67, 13478–13486. doi:10.1021/acs.jafc.9b01367
- Levisson, M., Patinios, C., Hein, S., De Groot, P. A., Daran, J.-M., Hall, R. D., et al. (2018). Engineering De Novo Anthocyanin Production in *Saccharomyces cerevisiae*. *Microb. Cell. Fact.* 17, 103. doi:10.1186/s12934-018-0951-6
- Li, G., Li, H., Lyu, Y., Zeng, W., and Zhou, J. (2020). Enhanced Biosynthesis of Dihydromyricetin in *Saccharomyces cerevisiae* by Coexpression of Multiple Hydroxylases. *J. Agric. Food Chem.* 68, 14221–14229. doi:10.1021/acs.jafc.0c05261
- Li, H., Ban, Z., Qin, H., Ma, L., King, A. J., and Wang, G. (2015). A Heteromeric Membrane-Bound Prenyltransferase Complex from Hop Catalyzes Three Sequential Aromatic Prenylations in the Bitter Acid Pathway. *Plant Physiol.* 167, 650–659. doi:10.1104/pp.114.253682
- Lim, E.-K., Ashford, D. A., Hou, B., Jackson, R. G., and Bowles, D. J. (2004). *Arabidopsis* Glycosyltransferases as Biocatalysts in Fermentation for Regioselective Synthesis of Diverse Quercetin Glucosides. *Biotechnol. Bioeng.* 87, 623–631. doi:10.1002/bit.20154
- Liu, W., Zhang, B., and Jiang, R. (2017a). Improving Acetyl-CoA Biosynthesis in *Saccharomyces cerevisiae* via the Overexpression of Pantothenate Kinase and PDH Bypass. *Biotechnol. Biofuels* 10, 41. doi:10.1186/s13068-017-0726-z
- Liu, X., Cheng, J., Zhang, G., Ding, W., Duan, L., Yang, J., et al. (2018). Engineering Yeast for the Production of Breviscapine by Genomic Analysis and Synthetic Biology Approaches. *Nat. Commun.* 9, 448. doi:10.1038/s41467-018-02883-z
- Liu, X., Cheng, J., Zhu, X., Zhang, G., Yang, S., Guo, X., et al. (2020). De Novo Biosynthesis of Multiple Pinocembrin Derivatives in *Saccharomyces cerevisiae*. *ACS Synth. Biol.* 9, 3042–3051. doi:10.1021/acssynbio.0c00289
- Liu, X., Ding, W., and Jiang, H. (2017b). Engineering Microbial Cell Factories for the Production of Plant Natural Products: from Design Principles to Industrial-Scale Production. *Microb. Cell. Fact.* 16, 125. doi:10.1186/s12934-017-0732-7
- Lütke-Eversloh, T., and Stephanopoulos, G. (2005). Feedback Inhibition of Chorismate Mutase/prephenate Dehydrogenase (TyrA) of *Escherichia coli*: Generation and Characterization of Tyrosine-Insensitive Mutants. *Appl. Environ. Microbiol.* 71, 7224–7228. doi:10.1128/AEM.71.11.7224-7228.2005
- Lütke-Eversloh, T., and Stephanopoulos, G. (2007). L-tyrosine Production by Deregulated Strains of *Escherichia coli*. *Appl. Microbiol. Biotechnol.* 75, 103–110. doi:10.1007/s00253-006-0792-9
- Lv, Y., Gu, Y., Xu, J., Zhou, J., and Xu, P. (2020). Coupling Metabolic Addiction with Negative Autoregulation to Improve Strain Stability and Pathway Yield. *Metab. Eng.* 61, 79–88. doi:10.1016/j.ymben.2020.05.005
- Lv, Y., Marsafari, M., Koffas, M., Zhou, J., and Xu, P. (2019). Optimizing Oleaginous Yeast Cell Factories for Flavonoids and Hydroxylated Flavonoids Biosynthesis. *ACS Synth. Biol.* 8, 2514–2523. doi:10.1021/acssynbio.9b00193
- Lyu, X., Ng, K. R., Lee, J. L., Mark, R., and Chen, W. N. (2017). Enhancement of Naringenin Biosynthesis from Tyrosine by Metabolic Engineering of *Saccharomyces cerevisiae*. *J. Agric. Food Chem.* 65, 6638–6646. doi:10.1021/acs.jafc.7b02507
- Lyu, X., Zhao, G., Ng, K. R., Mark, R., and Chen, W. N. (2019). Metabolic Engineering of *Saccharomyces cerevisiae* for De Novo Production of Kaempferol. *J. Agric. Food Chem.* 67, 5596–5606. doi:10.1021/acs.jafc.9b01329
- Marín, L., Gutiérrez-del-Río, I., Entrialgo-Cadierno, R., Villar, C. J., and Lombó, F. (2018). De Novo biosynthesis of Myricetin, Kaempferol and Quercetin in *Streptomyces albus* and *Streptomyces coelicolor*. *PLoS One* 13, e0207278. doi:10.1371/journal.pone.0207278
- Mark, R., Lyu, X., Ng, K. R., and Chen, W. N. (2019). Gene Source Screening as a Tool for Naringenin Production in Engineered *Saccharomyces cerevisiae*. *ACS Omega* 4, 12872–12879. doi:10.1021/acsomega.9b00364
- Medema, M. H., and Osbourn, A. (2016). Computational Genomic Identification and Functional Reconstitution of Plant Natural Product Biosynthetic Pathways. *Nat. Prod. Rep.* 33, 951–962. doi:10.1039/c6np00035e
- Milke, L., Ferreira, P., Kallscheuer, N., Braga, A., Vogt, M., Kappelmann, J., et al. (2019). Modulation of the Central Carbon Metabolism of *Corynebacterium glutamicum* Improves malonyl-CoA Availability and Increases Plant Polyphenol Synthesis. *Biotechnol. Bioeng.* 116, 1380–1391. doi:10.1002/bit.26939

- Milke, L., and Marienhagen, J. (2020). Engineering Intracellular Malonyl-CoA Availability in Microbial Hosts and its Impact on Polyketide and Fatty Acid Synthesis. *Appl. Microbiol. Biotechnol.* 104, 6057–6065. doi:10.1007/s00253-020-10643-7
- Mukai, R., Horikawa, H., Fujikura, Y., Kawamura, T., Nemoto, H., Nikawa, T., et al. (2012). Prevention of Disuse Muscle Atrophy by Dietary Ingestion of 8-prenylnaringenin in Denervated Mice. *PLoS One* 7, e45048. doi:10.1371/journal.pone.0045048
- Ni, J., Tao, F., Wang, Y., Yao, F., and Xu, P. (2016). A Photoautotrophic Platform for the Sustainable Production of Valuable Plant Natural Products from CO<sub>2</sub>. *Green Chem.* 18, 3537–3548. doi:10.1039/c6gc00317f
- Palmer, C. M., Miller, K. K., Nguyen, A., and Alper, H. S. (2020). Engineering 4-Coumaroyl-CoA Derived Polyketide Production in *Yarrowia Lipolytica* through a  $\beta$ -oxidation Mediated Strategy. *Metab. Eng.* 57, 174–181. doi:10.1016/j.ymben.2019.11.006
- Pittard, J., Camakaris, H., and Yang, J. (2005). The TyrR Regulon. *Mol. Microbiol.* 55, 16–26. doi:10.1111/j.1365-2958.2004.04385.x
- Ralston, L., Subramanian, S., Matsuno, M., and Yu, O. (2005). Partial Reconstruction of Flavonoid and Isoflavonoid Biosynthesis in Yeast Using Soybean Type I and Type II Chalcone Isomerases. *Plant Physiol.* 137, 1375–1388. doi:10.1104/pp.104.054502
- Rodriguez, A., Kildegaard, K. R., Li, M., Borodina, I., and Nielsen, J. (2015). Establishment of a Yeast Platform Strain for Production of P-Coumaric Acid through Metabolic Engineering of Aromatic Amino Acid Biosynthesis. *Metab. Eng.* 31, 181–188. doi:10.1016/j.ymben.2015.08.003
- Rodriguez, A., Martnez, J. A., Flores, N., Escalante, A., Gosset, G., and Bolivar, F. (2014). Engineering *Escherichia coli* to Overproduce Aromatic Amino Acids and Derived Compounds. *Microb. Cell. Fact.* 13, 126. doi:10.1186/s12934-014-0126-z
- Rodriguez, A., Strucko, T., Stahlhut, S. G., Kristensen, M., Svendsen, D. K., Forster, J., et al. (2017). Metabolic Engineering of Yeast for Fermentative Production of Flavonoids. *Bioresour. Technol.* 245, 1645–1654. doi:10.1016/j.biortech.2017.06.043
- Ruprecht, C., Bönisch, F., Ilmberger, N., Heyer, T. V., Haupt, E. T. K., Streit, W. R., et al. (2019). High Level Production of Flavonoid Rhamnosides by Metagenome-Derived Glycosyltransferase C in *Escherichia coli* Utilizing Dextrins of Starch as a Single Carbon Source. *Metab. Eng.* 55, 212–219. doi:10.1016/j.ymben.2019.07.002
- Santos, C. N. S., Koffas, M., and Stephanopoulos, G. (2011). Optimization of a Heterologous Pathway for the Production of Flavonoids from Glucose. *Metab. Eng.* 13, 392–400. doi:10.1016/j.ymben.2011.02.002
- Sasaki, K., Tsurumaru, Y., and Yazaki, K. (2009). Prenylation of Flavonoids by Biotransformation of Yeast Expressing Plant Membrane-Bound Prenyltransferase SfN8DT-1. *Biosci. Biotechnol. Biochem.* 73, 759–761. doi:10.1271/bbb.80729
- Schnappauf, G., Sträter, N., Lipscomb, W. N., and Braus, G. H. (1997). A Glutamate Residue in the Catalytic Center of the Yeast Chorismate Mutase Restricts Enzyme Activity to Acidic Conditions. *Proc. Natl. Acad. Sci. U.S.A.* 94, 8491–8496. doi:10.1073/pnas.94.16.8491
- Shabbir Hussain, M., Wheeldon, I., and Blenner, M. A. (2017). A Strong Hybrid Fatty Acid Inducible Transcriptional Sensor Built from *Yarrowia Lipolytica* Upstream Activating and Regulatory Sequences. *Biotechnol. J.* 12. doi:10.1002/biot.201700248
- Shrestha, L., Marasini, B. P., Pradhan, S. P., Shrestha, R. K., Shrestha, S., Regmi, K. P., et al. (2021). Biotransformation of Daidzein, Genistein, and Naringenin by *Streptomyces* Species Isolated from High-Altitude Soil of Nepal. *Int. J. Microbiol.* 2021, 9948738. doi:10.1155/2021/9948738
- Siedler, S., Stahlhut, S. G., Malla, S., Maury, J., and Neves, A. R. (2014). Novel Biosensors Based on Flavonoid-Responsive Transcriptional Regulators Introduced into *Escherichia coli*. *Metab. Eng.* 21, 2–8. doi:10.1016/j.ymben.2013.10.011
- Snoek, T., Chaberski, E. K., Ambri, F., Kol, S., Bjørn, S. P., Pang, B., et al. (2020). Evolution-guided Engineering of Small-Molecule Biosensors. *Nucleic Acids Res.* 48, e3. doi:10.1093/nar/gkz954
- Solopova, A., Van Tilburg, A. Y., Foito, A., Allwood, J. W., Stewart, D., Kulakauskas, S., et al. (2019). Engineering *Lactococcus Lactis* for the Production of Unusual Anthocyanins Using Tea as Substrate. *Metab. Eng.* 54, 160–169. doi:10.1016/j.ymben.2019.04.002
- Sordon, S., Madej, A., Popłoński, J., Bartmańska, A., Tronina, T., Brzezowska, E., et al. (2016). Regioselective Ortho-Hydroxylations of Flavonoids by Yeast. *J. Agric. Food Chem.* 64, 5525–5530. doi:10.1021/acs.jafc.6b02210
- Tohge, T., De Souza, L. P., and Fernie, A. R. (2017). Current Understanding of the Pathways of Flavonoid Biosynthesis in Model and Crop Plants. *J. Exp. Bot.* 68, 4013–4028. doi:10.1093/jxb/erx177
- Tominaga, M., Nozaki, K., Umeno, D., Ishii, J., and Kondo, A. (2021). Robust and Flexible Platform for Directed Evolution of Yeast Genetic Switches. *Nat. Commun.* 12, 1846. doi:10.1038/s41467-021-22134-y
- Trantas, E., Panopoulos, N., and Ververidis, F. (2009). Metabolic Engineering of the Complete Pathway Leading to Heterologous Biosynthesis of Various Flavonoids and Stilbenoids in *Saccharomyces cerevisiae*. *Metab. Eng.* 11, 355–366. doi:10.1016/j.ymben.2009.07.004
- Unamba, C. I. N., Nag, A., and Sharma, R. K. (2015). Next Generation Sequencing Technologies: The Doorway to the Unexplored Genomics of Non-model Plants. *Front. Plant Sci.* 6, 1074. doi:10.3389/fpls.2015.01074
- Vanegas, K. G., Larsen, A. B., Eichenberger, M., Fischer, D., Mortensen, U. H., and Naesby, M. (2018). Indirect and Direct Routes to C-Glycosylated Flavones in *Saccharomyces cerevisiae*. *Microb. Cell. Fact.* 17, 107. doi:10.1186/s12934-018-0952-5
- Wang, H., Yang, Y., Lin, L., Zhou, W., Liu, M., Cheng, K., et al. (2016a). Engineering *Saccharomyces cerevisiae* with the Deletion of Endogenous Glucosidases for the Production of Flavonoid Glucosides. *Microb. Cell. Fact.* 15, 134. doi:10.1186/s12934-016-0535-2
- Wang, R., Cress, B. F., Yang, Z., Hordines, J. C., 3rd, Zhao, S., Jung, G. Y., et al. (2019). Design and Characterization of Biosensors for the Screening of Modular Assembled Naringenin Biosynthetic Library in *Saccharomyces cerevisiae*. *ACS Synth. Biol.* 8, 2121–2130. doi:10.1021/acssynbio.9b00212
- Wang, T. Y., Tsai, Y. H., Yu, I. Z., and Chang, T. S. (2016b). Improving 3'-Hydroxygenistein Production in Recombinant *Pichia pastoris* Using Periodic Hydrogen Peroxide-Shocking Strategy. *J. Microbiol. Biotechnol.* 26, 498–502. doi:10.4014/jmb.1509.09013
- Wang, Y., Cheng, H., Liu, Y., Liu, Y., Wen, X., Zhang, K., et al. (2021). In-situ Generation of Large Numbers of Genetic Combinations for Metabolic Reprogramming via CRISPR-Guided Base Editing. *Nat. Commun.* 12, 678. doi:10.1038/s41467-021-21003-y
- Watts, K. T., Lee, P. C., and Schmidt-Dannert, C. (2004). Exploring Recombinant Flavonoid Biosynthesis in Metabolically Engineered *Escherichia coli*. *Chembiochem* 5, 500–507. doi:10.1002/cbic.200300783
- Wei, W., Zhang, P., Shang, Y., Zhou, Y., and Ye, B.-C. (2020). Metabolically Engineering of *Yarrowia Lipolytica* for the Biosynthesis of Naringenin from a Mixture of Glucose and Xylose. *Bioresour. Technol.* 314, 123726. doi:10.1016/j.biortech.2020.123726
- Werner, S. R., and Morgan, J. A. (2010). Controlling Selectivity and Enhancing Yield of Flavonoid Glycosides in Recombinant Yeast. *Bioprocess Biosyst. Eng.* 33, 863–871. doi:10.1007/s00449-010-0409-7
- Werner, S. R., and Morgan, J. A. (2009). Expression of a Dianthus Flavonoid Glycosyltransferase in *Saccharomyces cerevisiae* for Whole-Cell Biocatalysis. *J. Biotechnol.* 142, 233–241. doi:10.1016/j.jbiotec.2009.05.008
- Winkel, B. S. J. (2006). "The Biosynthesis of Flavonoids," in *The Science of Flavonoids*. Editor E. Grotenow (New York, NY: Springer New York), 71–95. doi:10.1007/978-0-387-28822-2\_3
- Wu, J., Du, G., Chen, J., and Zhou, J. (2015). Enhancing Flavonoid Production by Systematically Tuning the Central Metabolic Pathways Based on a CRISPR Interference System in *Escherichia coli*. *Sci. Rep.* 5, 13477. doi:10.1038/srep13477
- Wu, J., Du, G., Zhou, J., and Chen, J. (2013). Metabolic Engineering of *Escherichia coli* for (2S)-Pinocembrin Production from Glucose by a Modular Metabolic Strategy. *Metab. Eng.* 16, 48–55. doi:10.1016/j.ymben.2012.11.009
- Wu, J., Yu, O., Du, G., Zhou, J., and Chen, J. (2014a). Fine-tuning of the Fatty Acid Pathway by Synthetic Antisense RNA for Enhanced (2 S)-Naringenin Production from L-Tyrosine in *Escherichia coli*. *Appl. Environ. Microbiol.* 80, 7283–7292. doi:10.1128/aem.02411-14
- Wu, J., Zhang, X., Dong, M., and Zhou, J. (2016a). Stepwise Modular Pathway Engineering of *Escherichia coli* for Efficient One-step Production of (2S)-Pinocembrin. *J. Biotechnol.* 231, 183–192. doi:10.1016/j.jbiotec.2016.06.007

- Wu, J., Zhang, X., Zhou, J., and Dong, M. (2016b). Efficient Biosynthesis of (2S)-Pinocembrin from D-Glucose by Integrating Engineering Central Metabolic Pathways with a pH-Shift Control Strategy. *Bioresour. Technol.* 218, 999–1007. doi:10.1016/j.biortech.2016.07.066
- Wu, J., Zhou, L., Duan, X., Peng, H., Liu, S., Zhuang, Q., et al. (2021). Applied Evolution: Dual Dynamic Regulations-Based Approaches in Engineering Intracellular Malonyl-CoA Availability. *Metab. Eng.* 67, 403–416. doi:10.1016/j.ymben.2021.08.004
- Wu, J., Zhou, T., Du, G., Zhou, J., and Chen, J. (2014b). Modular Optimization of Heterologous Pathways for De Novo Synthesis of (2S)-Naringenin in *Escherichia coli*. *PLoS One* 9, e101492. doi:10.1371/journal.pone.0101492
- Xiong, D., Lu, S., Wu, J., Liang, C., Wang, W., Wang, W., et al. (2017). Improving Key Enzyme Activity in Phenylpropanoid Pathway with a Designed Biosensor. *Metab. Eng.* 40, 115–123. doi:10.1016/j.ymben.2017.01.006
- Xiu, Y., Jang, S., Jones, J. A., Zill, N. A., Linhardt, R. J., Yuan, Q., et al. (2017). Naringenin-responsive Riboswitch-Based Fluorescent Biosensor Module for *Escherichia coli* Co-cultures. *Biotechnol. Bioeng.* 114, 2235–2244. doi:10.1002/bit.26340
- Xu, P., Ranganathan, S., Fowler, Z. L., Maranas, C. D., and Koffas, M. A. G. (2011). Genome-scale Metabolic Network Modeling Results in Minimal Interventions that Cooperatively Force Carbon Flux towards Malonyl-CoA. *Metab. Eng.* 13, 578–587. doi:10.1016/j.ymben.2011.06.008
- Xue, Y., Zhang, Y., Cheng, D., Daddy, S., and He, Q. (2014). Genetically Engineering *Synechocystis* Sp. Pasteur Culture Collection 6803 for the Sustainable Production of the Plant Secondary Metabolite P-coumaric Acid. *Proc. Natl. Acad. Sci. U.S.A.* 111, 9449–9454. doi:10.1073/pnas.1323725111
- Yan, Y., Chemler, J., Huang, L., Martens, S., and Koffas, M. A. G. (2005a). Metabolic Engineering of Anthocyanin Biosynthesis in *Escherichia coli*. *Appl. Environ. Microbiol.* 71, 3617–3623. doi:10.1128/aem.71.7.3617-3623.2005
- Yan, Y., Kohli, A., and Koffas, M. A. G. (2005b). Biosynthesis of Natural Flavanones in *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* 71, 5610–5613. doi:10.1128/aem.71.9.5610-5613.2005
- Yang, D., Kim, W. J., Yoo, S. M., Choi, J. H., Ha, S. H., Lee, M. H., et al. (2018). Repurposing Type III Polyketide Synthase as a Malonyl-CoA Biosensor for Metabolic Engineering in Bacteria. *Proc. Natl. Acad. Sci. U.S.A.* 115, 9835–9844. doi:10.1073/pnas.1808567115
- Yang, Y., Lin, Y., Li, L., Linhardt, R. J., and Yan, Y. (2015). Regulating Malonyl-CoA Metabolism via Synthetic Antisense RNAs for Enhanced Biosynthesis of Natural Products. *Metab. Eng.* 29, 217–226. doi:10.1016/j.ymben.2015.03.018
- Zha, J., Zang, Y., Mattozzi, M., Plassmeier, J., Gupta, M., Wu, X., et al. (2018). Metabolic Engineering of *Corynebacterium Glutamicum* for Anthocyanin Production. *Microb. Cell. Fact.* 17, 143. doi:10.1186/s12934-018-0990-z
- Zhang, Q., Yu, S., Lyu, Y., Zeng, W., and Zhou, J. (2021). Systematically Engineered Fatty Acid Catabolite Pathway for the Production of (2S)-Naringenin in *Saccharomyces cerevisiae*. *ACS Synth. Biol.* 10, 1166–1175. doi:10.1021/acssynbio.1c00002
- Zhang, W., Liu, H., Li, X., Liu, D., Dong, X.-T., Li, F.-F., et al. (2017). Production of Naringenin from D-Xylose with Co-culture of *E. coli* and *S. Cerevisiae*. *Eng. Life Sci.* 17, 1021–1029. doi:10.1002/elsc.201700039
- Zhao, S., Jones, J. A., Lachance, D. M., Bhan, N., Khalidi, O., Venkataraman, S., et al. (2015). Improvement of Catechin Production in *Escherichia coli* through Combinatorial Metabolic Engineering. *Metab. Eng.* 28, 43–53. doi:10.1016/j.ymben.2014.12.002
- Zhou, H., Liao, X., Wang, T., Du, G., and Chen, J. (2010). Enhanced L-Phenylalanine Biosynthesis by Co-expression of pheA<sub>br</sub> and aroF<sub>w</sub>t. *Bioresour. Technol.* 101, 4151–4156. doi:10.1016/j.biortech.2010.01.043
- Zhou, S., Hao, T., and Zhou, J. (2020). Fermentation and Metabolic Pathway Optimization to De Novo Synthesize (2S)-Naringenin in *Escherichia coli*. *J. Microbiol. Biotechnol.* 30, 1574–1582. doi:10.4014/jmb.2008.08005
- Zhou, S., Lyu, Y., Li, H., Koffas, M. A. G., and Zhou, J. (2019). Fine-tuning the (2S)-naringenin Synthetic Pathway Using an Iterative High-throughput Balancing Strategy. *Biotechnol. Bioeng.* 116, 1392–1404. doi:10.1002/bit.26941
- Zhou, S., Yuan, S.-F., Nair, P. H., Alper, H. S., Deng, Y., and Zhou, J. (2021). Development of a Growth Coupled and Multi-Layered Dynamic Regulation Network Balancing Malonyl-CoA Node to Enhance (2S)-Naringenin Biosynthesis in *Escherichia coli*. *Metab. Eng.* 67, 41–52. doi:10.1016/j.ymben.2021.05.007
- Zhu, S., Wu, J., Du, G., Zhou, J., and Chen, J. (2014). Efficient Synthesis of Eriodictyol from L-Tyrosine in *Escherichia coli*. *Appl. Environ. Microbiol.* 80, 3072–3080. doi:10.1128/aem.03986-13

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Isogai, Tominaga, Kondo and Ishii. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.