



Implications of interleukin-17 in psoriatic lesions as Koebner phenomenon caused by recurrent occupational burns

Sato, Ayumi
Fukumoto, Takeshi
Yoshioka, Ai
Nishigori, Chikako

(Citation)

Dermatology Reports, 15(2):9567

(Issue Date)

2023

(Resource Type)

journal article

(Version)

Version of Record

(Rights)

©Copyright: the Author(s), 2023

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

(URL)

<https://hdl.handle.net/20.500.14094/0100482602>



Implications of interleukin-17 in psoriatic lesions as Koebner phenomenon caused by recurrent occupational burns

Ayumi Sato, Takeshi Fukumoto, Ai Yoshioka, Chikako Nishigori

Division of Dermatology, Department of Internal Related, Kobe University Graduate School of Medicine, Kobe, Japan

Abstract

The Koebner phenomenon (KP) is the emergence of new lesions in an uninvolved skin area caused by different types of stimulations, including mechanical stress, chemical stress, trauma, or injury. KP affects patients with certain skin diseases and is frequently observed in patients with psoriasis. We report the case of a 43-year-old obese male welder who developed psoriatic lesions only in areas of repeated burns due to his occupation. He was repeatedly exposed to mild burns in his anterior neck and the peri-orbital region as he was welding without shield protection. Subsequently, erythema appeared in the same region. Skin appearance and skin biopsy suggested psoriasis vulgaris (PV), and

immunohistochemical analysis of anti-interleukin (IL)-17, a crucial element in the development of PV, showed the positive-stained cells. The anti-IL-17 staining was prominent around the thickened epidermis as psoriatic lesions. IL-17 produced by T helper 17 cells stimulates keratinized cells and promotes chemokine secretion involved in neutrophil migration. Our case showed that patients, even without a history of PV, may have a risk of developing KP and PV via the enhanced production of IL-17 locally in the repeated burn area. No recurrence of skin symptoms was observed when the patient used a fully defensive shield during welding.

Introduction

Koebner phenomenon (KP) has been reported to occur in various diseases including psoriasis and some autoimmune vesicular diseases.^{1,2} It has been described as the emergence of psoriatic lesions in uninvolved skin areas after trauma in approximately 25-30% of psoriatic patients.¹⁻³ Many triggers such as tattooing, radiation, viral infections, and surgical incisions have been reported to develop KP.¹ We report a case of a welder who developed psoriatic lesions only in areas repeatedly exposed to mild occupational burns.

Case Report

A 43-year-old obese male working as a welder, with a high body mass index of 26.5, without a history of psoriasis vulgaris (PV) has repeatedly suffered from erythema with scaling on his unshielded neck and face during welding for more than 10 years. Scale erythema was observed on bilateral eyebrows, superior eyelids, temples, and the right anterolateral region of the neck (Figure 1a and b). Skin biopsy from the erythematous lesion on his neck revealed hyperkeratosis and parakeratosis of the epidermis, regular elongation of epidermal rete ridge, thinning or loss of the granular layer, and intracorneal micro-abscesses (Figure 1c and d). The immunohistochemical analysis of the anti-interleukin (IL)-17 antibody showed the positive-stained cells similar to previous reports, that were prominent around the thickened epidermis (Figure 1e).^{4,5} Notably, IL-17 produced by T helper 17 cells stimulates keratinized cells and promotes chemokine secretions involved in neutrophil migration.⁴ The patient was diagnosed with PV by clinical manifestations and histopathological examination. With topical application of maxacalcitol and dexamethasone valerate, the erythema had improved by the 2-month follow-up point (Figure 1f and g). Using a fully defensive shield over the face during welding prevented relapse of the lesions, and the patient has been in remission for 19 months.

Correspondence: Takeshi Fukumoto, Division of Dermatology, Department of Internal Related, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe 650-0017, Japan. Tel.: +81.783826134 - Fax: +81.783826149. E-mail: fuku@med.kobe-u.ac.jp

Key words: Koebner phenomenon, psoriasis, burn, IL-17.

Contributions: AS, TF, AY, CN, study design and draft of the manuscript; AS, TF, CN, data collection and interpretation of the results. All authors have read and approved the final manuscript.

Conflict of interest: the authors declare no potential conflict of interest.

Patient consent for publication: written informed consent was obtained from the patient.

Availability of data and materials: data and materials are available by the authors.

Acknowledgments: this work was supported by Kinoshita memorial enterprise subsidy.

Received for publication: 30 July 2022.

Accepted for publication: 16 September 2022.

Early view: 27 October 2022.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Dermatology Reports 2023; 15:9567

doi:10.4081/dr.2022.9567

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Discussion

This case developed psoriatic lesions selectively in areas of repeated minor occupational burns, suggesting that the burns may have caused KP and PV. Initially, KP was reported as the emergence of psoriatic lesions after skin trauma in the uninvolved skin of psoriatic patients.¹ Recently, the definition was extended to include lesions emerging after trauma in people without pre-existing skin disease.² KP is generally developed in 10 to 20 days after irritation, but some cases have been reported in a minimum of 3 days and a maximum of 2 years.³

Although the mechanism of KP has not been fully elucidated,

its involvement in epidermal cell damage and skin inflammation has been suggested in the pathophysiology.⁶ It has been reported that inflammatory mediators, including IL-17 produced by gamma-delta ($\gamma\delta$) T-cells, are involved in inducing the development of new psoriatic lesions.^{2,7} Moreover, $\gamma\delta$ T-cells are activated to enhance IL-17 production during the inflammation and the healing response after burn injury.^{8,9} We think that changes in the cytokine milieu in the burn area may have triggered KP. Guttate psoriasis and pityriasis rubra pilaris were reportedly caused by KP in burn areas.^{6,10} To clarify the mechanism of cytokine interactions related to KP in the burn area and the induced skin diseases, future accumulation of cases is required.




Figure 1. Clinical and histopathological features of the patient. a,b) Erythema on both upper eyelids, temples, and neck before the treatment; c,d) skin biopsy of the skin lesions on the patient's neck showing hyperkeratosis and parakeratosis of the epidermis, regular elongation of epidermal rete ridge, thinning of the granular layer, and intracorneal micro-abscess. Scale bar=100 μ m; e) the immunohistochemical analysis of anti-interleukin (IL)-17; rabbit polyclonal to IL-17 (Abcam, ab79056), showed the positive-stained cells in the skin; f,g) improvement in skin lesions 2 months after the treatment.

Conclusions

Our case showed that the patient, even without a history of PV, may have a risk to develop PV by enhanced production of IL-17 locally in the repeated burn area. PV, a chronic inflammatory disease, is associated with complications such as obesity, diabetes mellitus, metabolic syndrome, hypertension, and dyslipidemia.¹¹ In this patient, obesity is the only risk factor for developing PV. We speculate that PV was caused by the changes in the cytokine milieu in the skin following repeated trauma in a patient with predisposing factors for PV, even without a history of psoriasis. This is a rare case of PV occurring only in specific areas of repeated mild burns.

References

1. Boyd AS, Neldner KH. The isomorphic response of Koebner. *Int J Dermatol* 1990;29:401-10.
2. Ji YZ, Liu SR. Koebner phenomenon leading to the formation of new psoriatic lesions: evidences and mechanisms. *Biosci Rep* 2019;39:BSR20193266.
3. Arias-Santiago S, Espineira-Carmona MJ, Aneiros-Fernandez J. The Koebner phenomenon: psoriasis in tattoos. *CMAJ* 2013;185:585.
4. Picciani BLS, Domingos TA, Teixeira-Souza T, et al. Evaluation of the Th17 pathway in psoriasis and geographic tongue. *An Bras Dermatol* 2019;94:677-83.
5. Sun L, Liu Z, Lin Z, et al. Comparison of normal versus imiquimod-induced psoriatic skin in mice for penetration of drugs and nanoparticles. *Int J Nanomedicine* 2018;13:5625-35.
6. Acar EM, Has S, Kilitci A, Kemeriz F. Circumscribed juvenile pityriasis rubra pilaris (type 4) koebnerising after a hot water burn: mild disease with maximum Koebner response. *Acta Dermatovenerol Croat* 2019;27:47-9.
7. Cai Y, Shen X, Ding C, et al. Pivotal role of dermal IL-17-producing $\gamma\delta$ T cells in skin inflammation. *Immunity* 2011;35:596-610.
8. Sasaki JR, Zhang Q, Schwacha MG. Burn induces a Th-17 inflammatory response at the injury site. *Burns* 2011;37:646-51.
9. Rani M, Zhang Q, Schwacha MG. Burn wound $\gamma\delta$ T-cells support a Th2 and Th17 immune response. *J Burn Care Res* 2014;35:46-53.
10. Schicchi A, Tresoldi MM, Petrolini VM, et al. Guttate psoriasis: a case of unusual evolution of an occupation-related skin chemical burns. *Med Lav* 2021;20;112:168-70.
11. Yan D, Gudjonsson JE, Le S, et al. New frontiers in psoriatic disease research, part I: genetics, environmental triggers, immunology, pathophysiology, and precision medicine. *J Invest Dermatol* 2021;141:2112-22.