



# Bombardiers and assassins: mimetic interactions between unequally defended insects

Sugiura, Shinji  
Hayashi, Masakazu

---

(Citation)

PeerJ, 11:e15380

(Issue Date)

2023-06-06

(Resource Type)

journal article

(Version)

Version of Record

(Rights)

© 2023 Sugiura and Hayashi

This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, reproduction and adaptation in any medium and for any purpose provided that it is properly attributed...

(URL)

<https://hdl.handle.net/20.500.14094/0100482617>



# Bombardiers and assassins: mimetic interactions between unequally defended insects

Shinji Sugiura<sup>1</sup> and Masakazu Hayashi<sup>2</sup>

<sup>1</sup> Graduate School of Agricultural Science, Kobe University, Kobe, Hyogo, Japan

<sup>2</sup> Hoshizaki Green Foundation, Izumo, Shimane, Japan

## ABSTRACT

In defensive mimicry, resemblance between unequally defended species can be parasitic; this phenomenon has been termed quasi-Batesian mimicry. Few studies have used real co-mimics and their predators to test whether the mimetic interactions were parasitic. Here, we investigated the mimetic interaction between two well-defended insect species, the bombardier beetle *Pheropsophus occipitalis jessoensis* (Coleoptera: Carabidae) and the assassin bug *Sirthenia flavipes* (Hemiptera: Reduviidae), using their potential predator, the pond frog *Pelophylax nigromaculatus* (Anura: Ranidae), which coexists with these insect species in the same habitat in Japan. We observed behavioural responses of this frog species (adults and juveniles) to adult *Ph. occipitalis jessoensis* and adult *S. flavipes* under laboratory conditions. Among the frogs, 100% and 75% rejected *Ph. occipitalis jessoensis* and *S. flavipes*, respectively, suggesting that, compared with the assassin bug *S. flavipes*, the bombardier beetle *Ph. occipitalis jessoensis* is more well-defended against frogs. An assassin bug or a bombardier beetle was provided to a frog that had encountered the other insect species. Frogs with a history of assassin bug encounter demonstrated a lower rate of attack toward bombardier beetles. Similarly, frogs with a history of bombardier beetle encounter demonstrated a lower rate of attack toward assassin bugs. Therefore, both the bombardier beetle *Ph. occipitalis jessoensis* and the assassin bug *S. flavipes* benefit from the mimetic interaction.

**Subjects** Animal Behavior, Ecology, Entomology, Evolutionary Studies, Zoology

**Keywords** Assassin bugs, Bombardier beetles, Brachinini, Carabidae, Chemical defences, Co-mimics, Frogs, Müllerian mimicry, Quasi-Batesian mimicry, Reduviidae

## INTRODUCTION

Animals have evolved diverse anti-predator strategies such as chemical, morphological, physical, and behavioural defences (Edmunds, 1974; Ruxton, Sherratt & Speed, 2004; Eisner, Eisner & Siegler, 2005; Sugiura, 2020a). Some well-defended animals have aposematic body colours, which signal distaste and danger to their predators (i.e., warning signals; Ruxton, Sherratt & Speed, 2004). Well-defended species frequently share warning signals that serve to deter predation (i.e., Müllerian mimicry; Müller, 1878; Müller, 1879; Ruxton, Sherratt & Speed, 2004; Sherratt, 2008), while some non-defended species mimic well-defended species (i.e., Batesian mimicry; Bates, 1862; Ruxton, Sherratt & Speed, 2004). Müllerian mimicry is the mutualistic interaction between equally defended species, while Batesian mimicry

Submitted 28 April 2022

Accepted 18 April 2023

Published 6 June 2023

Corresponding author

Shinji Sugiura,  
sugiura.shinji@gmail.com,  
ssugiura@people.kobe-u.ac.jp

Academic editor

Joseph Gillespie

Additional Information and  
Declarations can be found on  
page 18

DOI 10.7717/peerj.15380

© Copyright

2023 Sugiura and Hayashi


Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

is the parasitic or commensal interaction between non-defended and well-defended species (Ruxton, Sherratt & Speed, 2004; Sherratt, 2008; Balogh, Gamberale-Stille & Leimar, 2008; Honma, Takakura & Nishida, 2008). These two types of mimicry are traditionally considered extreme ends of a defensive mimicry spectrum (Balogh, Gamberale-Stille & Leimar, 2008). On the mimicry spectrum, mimetic interactions between unequally defended species can be parasitic—this phenomenon is termed quasi-Batesian mimicry (Speed, 1993; Speed, 1999; Speed & Turner, 1999; Rowland et al., 2010). However, there is controversy regarding whether mimetic interactions between unequally defended species are truly parasitic (Speed et al., 2000; Rowland et al., 2007; Rowland et al., 2010; Aubier, Joron & Sherratt, 2017). Some experimental studies have indicated such interactions are parasitic (Speed et al., 2000; Rowland et al., 2010), while other studies have indicated that they are mutualistic (Rowland et al., 2007) or not always parasitic (Lindström et al., 2006). Previous experimental studies used bird predators and artificial prey to investigate the nature of quasi-Batesian mimicry (Speed et al., 2000; Lindström et al., 2006; Ihalainen, Lindström & Mappes, 2007; Rowland et al., 2007; Ihalainen et al., 2008; Rowland et al., 2010). Although resemblance between unequally defended species is commonly found in the natural environment (Marples, Brakefield & Cowie, 1989; Marples, 1993; Winters et al., 2018; Chouteau et al., 2019; Soukupová, Veselý & Fuchs, 2021), few studies have used real co-mimics and their natural predators to determine whether the mimetic interactions are parasitic or mutualistic (Pekár et al., 2017; Raška et al., 2020).

Bombardier beetles (Coleoptera: Carabidae: Brachininae: Brachinini) are chemically defended; their adults eject toxic chemicals at a temperature of 100 °C when they are attacked by predators (Aneshansley et al., 1969; Dean, 1979; Eisner, Eisner & Siegler, 2005; Arndt et al., 2015). The discharge of hot chemicals—namely, bombing—can protect beetles from various groups of predators such as birds (Kojima & Yamamoto, 2020), reptiles (Bonacci et al., 2008), amphibians (Eisner & Meinwald, 1966; Dean, 1980; Sugiura & Sato, 2018; Sugiura, 2018; Sugiura & Date, 2022), and arthropods (Eisner, 1958; Eisner & Meinwald, 1966; Eisner & Dean, 1976; Eisner et al., 2006; Sugiura, 2021). Many bombardier beetle species have similar aposematic body colours (Schaller et al., 2018; Anichtchenko et al., 2022) and are visually mimicked by some insect species that coexist with them in the same habitats (Shelford, 1902; Bonacci et al., 2008; Kojima & Yamamoto, 2020). For example, in Italy, the carabid species *Anchomenus dorsalis* (Pontoppidan) and the bombardier beetle species *Brachinus sclopeta* (Fabricius) have a similar body colour pattern (green-blue and red-brown) (Bonacci et al., 2008; Bonacci, Brandmayr & Brandmayr, 2011). In Borneo, a raspy cricket species (Orthoptera: Gryllacrididae) shares a black and orange body colour with the bombardier beetle *Pheropsophus* (*Stenaptinus*) *agnatus* (Chaudoir) (Shelford, 1902). In Japan, the assassin bug *Sirthenia flavipes* (Stål) (Hemiptera: Reduviidae: Peiratinae) has a black and yellow body colour similar to the colour of the bombardier beetle *Pheropsophus* (*Stenaptinus*) *occipitalis jessoensis* Morawitz (formerly called *Pheropsophus jessoensis*; Figs. 1A and 1B; Kojima & Yamamoto, 2020). Although the interaction between the carabid species *A. dorsalis* and the bombardier beetle *B. sclopeta* has been suggested to constitute Müllerian mimicry (Bonacci et al., 2008; Bonacci, Brandmayr



**Figure 1** A bombardier beetle, an assassin bug, and their potential predator. (A) An adult bombardier beetle *Pheropsophus occipitalis jessoensis*. (B) An adult assassin bug *Sirthenia flavipes*. (C) An adult pond frog *Pelophylax nigromaculatus*. The photos were taken in the same grassland (in Hyogo Prefecture) on 13 October 2021. Photo credit: Shinji Sugiura.

[Full-size](#) DOI: 10.7717/peerj.15380/fig-1

& Brandmayr, 2011), few studies have examined whether mimetic interactions between bombardier beetles and distantly related insects are parasitic or mutualistic.

To investigate mimetic interactions that involve bombardier beetles, we used two well-defended insect species (the bombardier beetle *Ph. occipitalis jessoensis* and the assassin bug *S. flavipes*) and their potential predator, the black-spotted pond frog *Pelophylax nigromaculatus* (Hallowell) (Anura: Ranidae), which coexists with these insects in the same habitat in Japan (Fig. 1).



The bombardier beetle *Ph. occipitalis jessoensis* (Fig. 1A) is distributed in East Asia (Japan, Korea, and China) and Southeast Asia (Vietnam) (Fedorenko, 2021). In Japan, adult *Ph. occipitalis jessoensis* are commonly found on the ground in farmland, grassland, and forest edges (Habu & Sadanaga, 1965; Yahiro et al., 1992; Ishitani & Yano, 1994; Fujisawa, Lee & Ishii, 2012; Ohwaki, Kaneko & Ikeda, 2015; Sugiura, 2018; Sugiura, 2021; Sugiura & Date, 2022). Female adults lay eggs in soil and the hatched larvae feed exclusively on egg masses of the mole cricket *Gryllotalpa orientalis* Burmeister (Orthoptera: Gryllotalpidae) (Akino, Sasaki & Okamoto, 1956; Habu & Sadanaga, 1965). Adults feed on live and dead insects of various species on the ground (Habu & Sadanaga, 1965; Sugiura, 2018). The bombardier beetle *Ph. occipitalis jessoensis* can eject quinones (1,4-benzoquinone and 2-methyl-1,4-benzoquinone) and water (vapor) at a temperature of approximately 100 °C from the end of its abdomen (Video S1; Kanehisa & Murase, 1977; Kanehisa, 1996) to repel predators such as the pond frog *Pe. nigromaculatus* (Sugiura, 2018), the bullfrog *Lithobates catesbeianus* (Shaw) (Anura: Ranidae) (Sugiura & Date, 2022), quails (Kojima & Yamamoto, 2020), and praying mantises (Sugiura, 2021).

Similar to bombardier beetles, assassin bugs are considered well-defended insects; assassin bugs kill prey insects and defend against their predators by using their proboscis (i.e., labium) to inject them with painful venoms (Eisner, Eisner & Siegler, 2005; Schmidt, 2009; Walker et al., 2016). In addition, assassin bugs have a variety of scent glands that can act as chemical defences (Louis, 1974; Staddon, 1979). The assassin bug *S. flavipes* (Fig. 1B) is distributed in East Asia (Japan, Korea, and China), Southeast Asia (Cambodia, Indonesia, Laos, Malesia, Myanmar, Philippines, Thailand, and Vietnam), South Asia (Bangladesh, India, Nepal, Pakistan, and Sri Lanka), and West Asia (Afghanistan and Iran) (Chlond, 2018). *Sirthenia flavipes* is found on the ground in grassland and farmland (Ito, Okutani & Hiura, 1977; Tomokuni et al., 1993; Takahashi, 1996; Hirashima & Morimoto, 2008). Similar to adults of the North American species *Sirthenia carinata* (Fabricius) (Hudson, 1987), juveniles and adults of *S. flavipes* exclusively prey on mole crickets (Hayashi, 2023); *S. flavipes* uses a long proboscis (labium) to inject paralyzing venoms into prey, then feeds on the prey (Video S2; Fig. 2). *Sirthenia flavipes* aggressively stabs attackers with its proboscis when it is caught (Ito, Okutani & Hiura, 1977; Yasunaga et al., 2018). A bite (stab) by adult *S. flavipes* reportedly causes severe pain to humans (Takara, 1957; Ito, Okutani & Hiura, 1977; Tomokuni et al., 1993; Takahashi, 1996; Yasunaga et al., 2018). The assassin bug *S. flavipes* shares the microhabitat (on the ground in grassland), prey (mole crickets), and body colour pattern (black and yellow colour) with the bombardier beetle *Ph. occipitalis jessoensis* (Fig. 1). However, the relationship between the bombardier beetle *Ph. occipitalis jessoensis* and assassin bug *S. flavipes* remains unexplored.

The black-spotted pond frog *Pe. nigromaculatus* is distributed in East Asia, including Japan, Korea, and China (Komaki et al., 2015; Matsui & Maeda, 2018). Postmetamorphic juveniles and adults of *Pe. nigromaculatus* are found in paddy fields and surrounding grasslands (Matsui & Maeda, 2018). Postmetamorphic juveniles and adults prey on animals smaller than themselves, especially terrestrial insects (Hirai & Matsui, 1999; Hirai, 2002; Honma, Oku & Nishida, 2006; Sano & Shinohara, 2012; Sarashina, Yoshihisa & Yoshida, 2011; Sugiura, 2018). *Pelophylax nigromaculatus* uses its tongue to catch and swallow prey



**Figure 2** An assassin bug *Sirthenea flavipes* feeding on the mole cricket *Gryllotalpa orientalis*. The arrow indicates the proboscis of the adult assassin bug. Photo credit: Shinji Sugiura.

Full-size DOI: [10.7717/peerj.15380/fig-2](https://doi.org/10.7717/peerj.15380/fig-2)

(Sugiura, 2018; Sugiura, 2020b; Sugiura & Tsujii, 2022). *Pelophylax nigromaculatus* is found with *Ph. occipitalis jessoensis* and *S. flavipes* on grasslands in central Japan (Fig. 1). Sugiura (2018) observed that almost all *Pe. nigromaculatus* juveniles and adults could attack the bombardier beetle *Ph. occipitalis jessoensis*, but ultimately rejected it under laboratory conditions. Therefore, the pond frog *Pe. nigromaculatus* was used as a model predator to investigate the mimetic interaction between *Ph. occipitalis jessoensis* and *S. flavipes*.

To elucidate the nature of the mimetic interaction between bombardier beetles and assassin bugs, we observed behavioural responses of the frog *Pe. nigromaculatus* to *Ph. occipitalis jessoensis* and *S. flavipes* under laboratory conditions. Specifically, we compared rates of rejection by *Pe. nigromaculatus* between *Ph. occipitalis jessoensis* and *S. flavipes*. To determine whether *Ph. occipitalis jessoensis* and/or *S. flavipes* benefits from the mimetic interaction, we experimentally investigated whether a frog would attack an insect species after it had encountered the other insect species. The rate of attack by frogs that had encountered bombardier beetles (or assassin bugs) was compared with the rate of attack by frogs that had not encountered the indicated species. Finally, we discuss the adaptive significance of mimetic interactions between bombardier beetles and assassin bugs.

## MATERIALS AND METHODS

### Sampling

Forty-three adults of the bombardier beetle *Ph. occipitalis jessoensis* were collected from Honshu (Hyogo, Shiga, and Shimane Prefectures) in July–September 2020 and May–October 2021. Adult beetles were housed separately in plastic cases (diameter: 85 mm; height: 25 mm) under laboratory conditions (25 °C; cf. Sugiura, 2018; Sugiura & Sato, 2018; Sugiura, 2021; Sugiura & Date, 2022) and fed dead larvae of *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) (cf. Sugiura, 2018; Sugiura & Sato, 2018; Sugiura, 2021; Sugiura & Date, 2022). Prior to experiments, body length and weight were measured to the nearest 0.01 mm and 0.1 mg using electronic slide callipers (CD-15AX, Mitsutoyo, Kawasaki, Japan) and an electronic balance (CPA64, Sartorius Japan K.K., Tokyo, Japan), respectively (Table 1). The same beetles were not used for multiple experiments.

**Table 1** Body sizes of bombardier beetles, assassin bugs, and pond frogs used in this study.

	Bombardier beetle <i>Pheropsophus occipitalis jessoensis</i>	Assassin bug <i>Sirthenea flavipes</i>	Pond frog <i>Pelophylax nigromaculatus</i>
Body length (mm) <sup>a</sup>	18.7 ± 0.2 (15.9–20.9)	20.3 ± 0.2 (18.6–21.9)	51.2 ± 0.9 (39.0–62.0)
Body weight (mg) <sup>a</sup>	291.9 ± 8.9 (160.2–412.9)	133.9 ± 6.7 (92.9–201.7)	13428.8 ± 828.0 (5497.4–29761.3)
<i>n</i>	43	20	48

**Notes.**

<sup>a</sup>Mean ± SE (range: minimum–maximum).

Twenty adults of the assassin bug *S. flavipes* were collected from Honshu (Hyogo and Shimane Prefectures) in August–October 2021. Adult bugs were housed separately in plastic cases (diameter: 85 mm; height: 25 mm) under laboratory conditions (25 °C). Prior to experiments, body length and weight were measured to the nearest 0.01 mm and 0.1 mg using electronic slide callipers and an electronic balance, respectively (Table 1). Adults and nymphs of the mole cricket *G. orientalis* were provided as prey (cf. Hayashi, 2023). Some assassin bugs were used repeatedly in different experiments.

Forty-eight adults and juveniles of the pond frog *Pe. nigromaculatus* were collected from Honshu (Hyogo Prefecture) in June–October 2021. Frogs were housed separately in plastic cages (length: 120 mm; width: 85 mm; height: 130 mm) in the laboratory at 25 °C (cf. Sugiura, 2018; Sugiura, 2020b; Sugiura & Tsujii, 2022). Live mealworms—larvae of *Tenebrio molitor* Linnaeus (Coleoptera: Tenebrionidae)—were provided as prey (cf. Sugiura, 2018; Sugiura, 2020b; Sugiura & Tsujii, 2022). Snout–vent length and body weight were measured to the nearest 0.01 mm and 0.1 mg using electronic slide callipers and an electronic balance, respectively (Table 1). Some frogs were used repeatedly in different experiments.

### Experiment 1: initial response

We used the predator *Pe. nigromaculatus* to test whether the bombardier beetle *Ph. occipitalis jessoensis* or the assassin bug *S. flavipes* is better defended under laboratory conditions. In accordance with the method established by Sugiura (2018), we experimentally investigated behavioural responses of *Pe. nigromaculatus* to *Ph. occipitalis jessoensis* and *S. flavipes* in the laboratory (Graduate School of Agricultural Science, Kobe University) between September 2021 and October 2021. We used frogs that had fed on mealworms >24 h before experiments. First, a frog was placed in a plastic cage (length: 120 mm; width: 85 mm; height: 130 mm). Next, a bombardier beetle (or an assassin bug) was transferred to the cage containing the frog. The behaviours of the frog and the bombardier beetle (or assassin bug) were recorded using a digital camera (iPhone 12 Pro Max; Apple Inc., Cupertino, CA, USA) and a digital video camera (Handycam HDR-PJ790V, Sony, Tokyo, Japan). The footage of recorded behaviour was reviewed to investigate how each insect could defend. The bombing sounds of bombardier beetles were checked to investigate whether bombing forced the frogs to reject the beetles. Stabbing by assassin bugs was investigated to determine whether stabbing forced the frogs to reject the bugs. When a frog did not attack a bombardier beetle or an assassin bug within 2 min, we considered it to have ignored the insect. When a frog swallowed a bombardier beetle or an assassin bug, we observed


whether the frog subsequently vomited the insect (cf. [Sugiura & Sato, 2018](#); [Sugiura, 2018](#); [Sugiura & Date, 2022](#)). When the frog did not vomit the insect, we inferred that the frog had digested the insect. In Experiment 1, we used 20 bombardier beetles, 20 assassin bugs, and 40 frogs. The body size (snout–vent length and body weight) of frogs that attacked bombardier beetles did not significantly differ from the body size of frogs that attacked assassin bugs ( $t$ -test; snout–vent length,  $t = -0.77455$ ,  $df = 37.923$ ,  $P = 0.4434$ ; body weight,  $t = 0.98556$ ,  $df = 35.179$ ,  $P = 0.3311$ ). The same individuals of bombardier beetles, assassin bugs, and frogs were not used repeatedly in this experiment. The sample size was determined based on the number of assassin bugs collected in this study. Experiment 1 was part of the following experiment; specifically, the initial responses of 35 frogs observed in Experiment 2 were used as the data for Experiment 1.

## Experiment 2: generalisation tests

We experimentally investigated the interaction between the bombardier beetle *Ph. occipitalis jessoensis* and the assassin bug *S. flavipes* via the potential predator *Pe. nigromaculatus* under laboratory conditions. Specifically, we investigated how a frog responded to a bombardier beetle or an assassin bug after the frog had encountered the other insect species (*i.e.*, generalisation test; [Fig. 3](#)). The same plastic cages and video cameras (see Experiment 1 for details) were used in this experiment. A bombardier beetle was provided to a frog that had encountered (attacked or ignored) an assassin bug ( $n = 23$ ; [Fig. 3A](#)). We tested whether the frog attacked or ignored the bombardier beetle approximately 6 min (median: 6 min; range: 5–14 min) after the frog had encountered an assassin bug. The rate of attack on bombardier beetles by frogs that had encountered assassin bugs was compared to the rate of attack by frogs that had not encountered assassin bugs. Similarly, an assassin bug was provided to a frog that had encountered a bombardier beetle ( $n = 20$ ; [Fig. 3B](#)). We tested whether the frog attacked or ignored the assassin bug approximately 6 min (median: 6 min; range: 5–7 min) after the frog had encountered a bombardier beetle. The rate of attack on assassin bugs by frogs that had encountered bombardier beetles was compared to the rate of attack by frogs that had not encountered bombardier beetles. Although the duration used in our generalisation tests (5–14 min) was shorter than the generalisation time of a spider (1 h; [Raška et al., 2020](#)) and the memory retention time of a bird (35 days; [Kojima & Yamamoto, 2020](#)), our field observations ([Fig. 1](#)) suggest that pond frogs frequently encounter bombardier beetles and assassin bugs under field conditions. Survivals of bombardier beetles, assassin bugs, and frogs used in this study were checked within 24 h after experiments. When a frog died within 24 h after the experiment, we dissected the frog to investigate the cause of death. In Experiment 2, we used 43 bombardier beetles, 17 assassin bugs, and 43 frogs. The same individuals of bombardier beetles and frogs were not used repeatedly in this experiment. The initial behavioural responses of 35 frogs observed in this experiment were also used as the data for Experiment 1. In addition, eight frogs were exclusively used in Experiment 2.

All experiments were performed in accordance with Kobe University Animal Experimentation Regulations (Kobe University's Animal Care and Use Committee, No. 30–01). Only one pair of insect species was provided to each frog to minimise the negative





**Figure 3** Experimental procedures and results of generalisation tests. (A) A bombardier beetle was provided to a frog that had encountered an assassin bug. (B) An assassin bug was provided to a frog that had encountered a bombardier beetle. Photo credit: Shinji Sugiura.


[Full-size](#) DOI: [10.7717/peerj.15380/fig-3](https://doi.org/10.7717/peerj.15380/fig-3)

impacts of well-defended insects on frogs. Healthy frogs were released after the experiments had been completed. No frogs were euthanised in this study.

## Data analysis

All analyses were performed using R version 3.5.2 ([R Core Team, 2018](#)).

Fisher's exact test was used to compare the rates of rejection by frogs between bombardier beetles and assassin bugs. Welch's *t*-test was used to compare the body size (body length and weight) of bombardier beetles and assassin bugs; it was also used to compare the body size (snout–vent length and body weight) of pond frogs that attacked bombardier beetles and assassin bugs. A generalised linear mixed model (GLMM) with a binomial error distribution and a logit link was used to investigate the effects of insect species and frog encounter history on the rate of attack by frogs. The frog response (attack, 1; or ignore, 0) was used as a response variable. The insect species (the bombardier beetle *Ph. occipitalis jessoensis* or the assassin bug *S. flavipes*), frog encounter history (an initial response or a response after encountering the other species), and the interaction between insect species and frog encounter history were used as fixed factors. Individual assassin bugs and frogs were used as random effects. The GLMM was conducted using the lme4 package version 1.1.19 in R ([Bates et al., 2015](#)). A significance threshold of 0.05 was used for all statistical tests.



**Figure 4** Behavioural responses of the frog *Pelophylax nigromaculatus* to the bombardier beetle *Pheropsophus occipitalis jessoensis* and the assassin bug *Sirthenaea flavipes*. Frogs that had not encountered the bombardier beetle or assassin bug were used in this study. Swallow: frogs successfully swallowed beetles (or bugs). Spit out: frogs spat out beetles (or bugs) immediately after taking the indicated insects into their mouths. Stop attack: frogs stopped their attacks immediately after their tongues had contacted beetles (or bugs). Ignore: frogs did not attack beetles (or bugs). Bombed: frogs were bombed by beetles. Stabbed: frogs were bitten (stabbed) by bugs. Photo credit: Shinji Sugiura. [Full-size DOI: 10.7717/peerj.15380/fig-4](https://doi.org/10.7717/peerj.15380/fig-4)


## RESULTS

### Experiment 1: initial response

Among the 20 frogs in this experiment, none successfully swallowed bombardier beetles (Fig. 4). Five frogs (25%) captured beetles in their mouths but spat out the beetles immediately after bombing had occurred in their mouths (Video S3; Figs. 4 and 5). Ten frogs (50%) stopped attacking beetles immediately after their tongues had contacted the beetles (Fig. 4). Five frogs (25.0%) ignored the beetles (Fig. 4).

Five (25%) of 20 frogs successfully swallowed assassin bugs (Fig. 3), although these frogs were stabbed by assassin bugs in their mouths (Fig. 4). Fifteen of the 20 frogs (75%) rejected assassin bugs. Five frogs (25%) captured assassin bugs in their mouths but spat out the bugs immediately after stabbing had occurred in their mouths (Video S4; Figs. 4 and 6). Eight frogs (40%) stopped attacking assassin bugs immediately after their tongues had contacted the bugs (Fig. 4). Two frogs (10.0%) ignored the bugs (Fig. 4).


The rate of bombardier beetle rejection by frogs (100.0%) significantly differed from the rate of assassin bug rejection by frogs (75.0%; Fisher's exact test,  $P = 0.0471$ ). The body size (body length and weight) significantly differed between bombardier beetles and assassin bugs; bombardier beetles were shorter but heavier than assassin bugs ( $t$ -test; body



**Figure 5** Temporal sequence of the frog *Pelophylax nigromaculatus* rejecting an adult bombardier beetle *Pheropsophus occipitalis jessoensis*. (A) 0 ms. (B) 929 ms. (C) 1,337 ms. (D) 3,780 ms. (E) 3,861 ms. (F) 3,918 ms. (G) 4,646 ms. (H) Close-up view (E), with the arrow indicating bombing from the tip of the abdomen of the adult *Ph. occipitalis jessoensis*. The frog spat out the bombardier beetle after bombing had occurred in its mouth (see [Video S3](#)). Credit: Shinji Sugiura.

Full-size [DOI: 10.7717/peerj.15380/fig-5](#)





**Figure 6** Temporal sequence of the frog *Pelophylax nigromaculatus* rejecting an adult assassin bug *Sirthenia flavipes*. (A) 0 ms. (B) 346 ms. (C) 1,332 ms. (D) 2,188 ms. (E) 2,256 ms. (F) 2,760 ms. (G) 2,959 ms. (H) Close-up view (E), with the arrow indicating the proboscis of the adult assassin bug. The frog spat out the assassin bug after stabbing had occurred in its mouth (see [Video S4](#)). Credit: Shinji Sugiura.

[Full-size](#) DOI: [10.7717/peerj.15380/fig-6](https://doi.org/10.7717/peerj.15380/fig-6)



**Table 2** Results of generalisation tests: responses of frogs to assassin bugs (first) and bombardier beetles (second).

	Frog behaviour <sup>a</sup>	Second trial: frog responses to bombardier beetles				
		Swallow	Spit out (bombed)	Stop attack	Ignore	Total
First trial: frog responses to assassin bugs	Swallow	0	0	0	5	5
	Spit out (stabbed)	0	1	1	2	4
	Stop attack	0	0	3	9	12
	Ignore	0	0	0	2	2
	Total	0	1	4	18	23

**Notes.**  
 Values: numbers of frogs.  
<sup>a</sup>Frog behaviour: Swallow: frogs successfully swallowed beetles (or bugs). Spit out: frogs spat out beetles (or bugs) immediately after capturing the insects in their mouths (frogs were bombed or stabbed). Stop attack: frogs stopped their attacks immediately after their tongues had contacted beetles (or bugs). Ignore: frogs did not attack beetles (or bugs).

**Table 3** Results of generalisation tests: responses of frogs to bombardier beetles (first) and assassin bugs (second).

	Frog behaviour <sup>a</sup>	Second trial: frog responses to assassin bugs				
		Swallow	Spit out (stabbed)	Stop attack	Ignore	Total
First trial: frog responses to bombardier beetles	Swallow	0	0	0	0	0
	Spit out (bombed)	0	1	0	4	5
	Stop attack	0	1	5	4	10
	Ignore	0	0	1	4	5
	Total	0	2	6	12	20

**Notes.**  
 Values: numbers of frogs.  
<sup>a</sup>Frog behaviour: Swallow: frogs successfully swallowed beetles (or bugs). Spit out: frogs spat out beetles (or bugs) immediately after capturing the insects in their mouths (frogs were bombed or stabbed). Stop attack: frogs stopped their attacks immediately after their tongues had contacted beetles (or bugs). Ignore: frogs did not attack beetles (or bugs).


length,  $t = -4.8737$ ,  $df = 35.532$ ,  $P < 0.0001$ ; body weight,  $t = 11.255$ ,  $df = 30.211$ ,  $P < 0.0001$ ).

## Experiment 2: generalisation tests

Twenty-one (91.3%) of 23 frogs attacked assassin bugs (Figs. 3A and 7; Table 2). Bombardier beetles were provided to the frogs that had encountered assassin bugs ( $n = 23$ ); five frogs (21.7%) attacked bombardier beetles (Figs. 3A and 7; Table 2), while 18 frogs (78.3%) ignored bombardier beetles (Table 2).

Fifteen (75.0%) of 20 frogs attacked bombardier beetles (Figs. 3B and 7; Table 3). Assassin bugs were provided to the frogs that had encountered bombardier beetles ( $n = 20$ ); eight frogs (40.0%) attacked assassin bugs (Figs. 3B and 7), while 12 frogs (60.0%) ignored assassin bugs (Table 3).

A history of assassin bug encounter reduced the rate of attack on bombardier beetles by frogs from 75.0% to 21.7% (Fig. 7), although the rate of rejection by frogs did not change (100%; Fig. 8A). A history of bombardier beetle encounter reduced the rate of attack on assassin bugs by frogs from 91.3% to 40.0% (Fig. 7); the rate of rejection increased from 78.3% to 100.0% (Fig. 8B). The GLMM analysis showed that frog encounter history had a significant influence on the rate of attack by frogs, although insect species and the



**Figure 7** Rates of attack on the bombardier beetle *Pheropsophus occipitalis jessoensis* and the assassin bug *Sirthenia flavipes* by the frog *Pelophylax nigromaculatus* before and after encounters with the other insect. Photo credit: Shinji Sugiura.

Full-size [DOI: 10.7717/peerj.15380/fig-7](https://doi.org/10.7717/peerj.15380/fig-7)

**Table 4** Results of a generalised linear mixed model (GLMM) to identify factors that influenced the rates of attack on bombardier beetles and assassin bugs by frogs.

Response variable	Explanatory variable (fixed factor)	Coefficient estimate	SE	z value	P value
Attack	Intercept	3.2264	1.4013	2.302	0.0213
	Insect species (vs. bombardier beetles) <sup>a</sup>	−1.5950	1.2290	−1.298	0.1943
	Frog encounter history (vs. encounter) <sup>b</sup>	−3.8513	1.7386	−2.215	0.0267
	Insect species × frog experience	0.2856	1.6187	0.176	0.8599

**Notes.**

<sup>a</sup>Assassin bugs were used as a reference.

<sup>b</sup>Initial (no previous encounter) response was used as a reference.

interaction between insect species and frog encounter history did not have significant effects on the rate of attack (Table 4).


## Survival

None of the bombardier beetles or assassin bugs that successfully defended against frogs died within 24 h after the experiments. Of the frogs ( $n = 48$ ), one (2.1%) died within 24 h (4 h) after swallowing an assassin bug; a dead assassin bug was found in the stomach of the dead frog. Other frogs (97.9%) were not harmed by our experiments.


## DISCUSSION

There has been controversy regarding whether mimetic interactions between unequally defended species are parasitic (Speed *et al.*, 2000; Rowland *et al.*, 2007; Rowland *et al.*, 2010;

(A) Bombardier beetles



(B) Assassin bugs



**Figure 8** Responses of the frog *Pelophylax nigromaculatus* to the bombardier beetle *Pheropsophus occipitalis jessoensis* and the assassin bug *Sirthenia flavipes* after the frog encountered the other insect species. Swallow: frogs successfully swallowed beetles (or bugs). Spit out: frogs spat out beetles (or bugs) immediately after capturing the insects in their mouths (frogs were bombed or stabbed). Stop attack: frogs stopped their attacks immediately after their tongues had contacted beetles (or bugs). Ignore: frogs did not attack beetles (or bugs). Photo credit: Shinji Sugiura.

Full-size DOI: 10.7717/peerj.15380/fig-8

Aubier, Joron & Sherratt, 2017). In the present study, we showed that both the bombardier beetle *Ph. occipitalis jessoensis* and the assassin bug *S. flavipes* were well-defended against the frog *Pe. nigromaculatus* (Fig. 4). In generalisation tests (Experiment 2), frogs with a history of assassin bug encounter attacked bombardier beetles less frequently compared with frogs

that had no such encounter history (Figs. 3A, 7 and 8A). Similarly, frogs with a history of bombardier beetle encounter attacked assassin bugs less frequently compared with frogs that had no such encounter history (Figs. 3B, 7 and 8B). These results suggest that both the bombardier beetle *Ph. occipitalis jessoensis* and the assassin bug *S. flavipes* benefit from the mimetic interaction in terms of defence against the potential predator *Pe. nigromaculatus*. Although *Ph. occipitalis jessoensis* demonstrated superior defensive abilities compared to *S. flavipes* (Figs. 4 and 8), the mimetic interaction between these two unequally defended species may be mutualistic, rather than parasitic.

## Frogs as predators

Frogs have been frequently used as predators to investigate the effectiveness of anti-predator defences in insects (Taniguchi et al., 2005; Ito, Taniguchi & Billen, 2016; Matsubara & Sugiura, 2017; Sugiura, 2018; Shinohara & Takami, 2020; Sugiura, 2020a; Sugiura, 2020b; Sugiura & Date, 2022; Sugiura & Tsujii, 2022). In this study, we used pond frogs as predators of bombardier beetles and assassin bugs to show that the frog species *Pe. nigromaculatus* was unable to distinguish between the bombardier beetle *Ph. occipitalis jessoensis* and the assassin bug *S. flavipes*. This finding could be explained by the adaptive generalisation in predators, where they learn to recognise dangerous prey by generalising the appearance of previously encountered prey to that of subsequently encountered prey (Ruxton et al., 2008).

In the Experiment 2, the time used in the generalisation tests ranged from 5–14 min. The time used in previous memory and generalisation tests (1 h–35 days; Ito, Taniguchi & Billen, 2016; Kojima & Yamamoto, 2020; Raška et al., 2020) was longer than the time of our generalisation tests. For example, the tree frog *Dryophytes japonica* (Günther) reportedly retains the memory of unpalatable prey for at least 1 day (Ito, Taniguchi & Billen, 2016). In addition, the *Pe. nigromaculatus* individuals used in this study were collected from the study sites where both *Ph. occipitalis jessoensis* and *S. flavipes* were found, suggesting that some individuals of *Pe. nigromaculatus* may have already experienced *Ph. occipitalis jessoensis* and/or *S. flavipes* at the sites prior to our experiments. Therefore, the use of short experimental durations and wild-collected individuals may have influenced the results of our experiments in *Pe. nigromaculatus*. Further studies are needed to investigate detailed memory retention in *Pe. nigromaculatus*.

Some predators have evolved counter defences, such as specific skills to avoid well-defended prey by detecting toxic chemicals or recognising warning signals (Edmunds, 1974; Endler, 1991; Ruxton, Sherratt & Speed, 2004; Skelhorn & Rowe, 2006; Williams et al., 2010). In the present study, 50% and 40% of frogs stopped attacking *Ph. occipitalis jessoensis* and *S. flavipes* before they had been bombed and stabbed, respectively (Fig. 4). Because *Pe. nigromaculatus* individuals stopped attacking immediately after their tongues had contacted these insects, this frog species may quickly detect deterrent characteristics on the body surfaces of *Ph. occipitalis jessoensis* and *S. flavipes* with its tongue; this enables avoidance of damage (Sugiura, 2018). Such reactions to well-defended prey have been reported in other predators such as tree frogs (Ito, Taniguchi & Billen, 2016) and quails (Kojima & Yamamoto, 2020).



## Bombardier beetles as models and mimics

Bombardier beetles can chemically defend themselves against various types of predators (Eisner, 1958; Eisner & Meinwald, 1966; Eisner & Dean, 1976; Dean, 1980; Eisner et al., 2006; Bonacci et al., 2008; Sugiura & Sato, 2018; Sugiura, 2018; Kojima & Yamamoto, 2020; Sugiura, 2021; Sugiura & Date, 2022). Many bombardier beetle species have aposematic body colour patterns that advertise their toxicity to predators (Schaller et al., 2018; Anichtchenko et al., 2022). Therefore, bombardier beetles are visually mimicked by distantly related insects that coexist with them in the same habitats (Shelford, 1902; Bonacci et al., 2008; Kojima & Yamamoto, 2020). However, very few studies have elucidated the nature of mimetic interactions that include bombardier beetles. In the present study, we used the frog *Pe. nigromaculatus* as a potential predator to investigate that the mimetic interaction between the bombardier beetle *Ph. occipitalis jessoensis* and the assassin bug *S. flavipes*. We found that a history of encounter with *Ph. occipitalis jessoensis* reduced the rate of attack on *S. flavipes* (Fig. 8B), suggesting that the coexistence with *Ph. occipitalis jessoensis* is beneficial for *S. flavipes*. However, *Ph. occipitalis jessoensis* consistently repelled the frog *Pe. nigromaculatus* in our study (Figs. 4 and 8A). Therefore, a history of encounters with the assassin bug *S. flavipes* may not benefit *Ph. occipitalis jessoensis*. Nevertheless, the mortality risk from frog attacks is not zero, as a previous study reported that 3.6% of the frog *Pe. nigromaculatus* successfully ate *Ph. occipitalis jessoensis* (Sugiura, 2018). Thus, the coexistence with the assassin bug *S. flavipes* is beneficial for the bombardier beetle *Ph. occipitalis jessoensis*, although the mutualistic interaction between these two insect species may be asymmetric.

## Assassin bugs as mimics

Ground-dwelling assassin bugs that belong to the subfamily Peiratinae reportedly stab with their proboscises, causing severe pain in humans (Readio, 1927; Ito, Okutani & Hiura, 1977; Willemse, 1985; Tomokuni et al., 1993; Gil-Santana, Forero & Weirauch, 2015; Yasunaga et al., 2018). Assassin bugs can paralyze prey and repel enemies through the injection of saliva or venom (Eisner, Eisner & Siegler, 2005; Schmidt, 2009; Walker et al., 2016). Assassin bugs also have scent glands to chemically defend themselves against predators (Louis, 1974; Staddon, 1979). However, few studies have investigated the effectiveness of anti-predator defences in assassin bugs (Walker et al., 2018; Walker et al., 2019).

In this study, we showed that the frog species *Pe. nigromaculatus* frequently rejected the assassin bug *S. flavipes* (Figs. 4 and 8). Although some frogs successfully swallowed *S. flavipes* individuals (Figs. 4 and 8B), one frog died 4 h after a successful swallowing event. These results suggest that *S. flavipes* venom is sufficiently strong to repel predators. In addition, some frogs stopped attacking assassin bugs immediately after their tongues had contacted the bugs (Fig. 4), suggesting that chemicals on the body surfaces of *S. flavipes* act as a deterrent to *Pe. nigromaculatus*.

Some assassin bug species share body colour patterns with hymenopteran insects such as paper wasps, ichneumonid wasps, spider wasps, and stingless bees (Maldonado Capriles & Lozada Robles, 1992; Zhang & Weirauch, 2014), suggesting that they mimic wasps (Haviland, 1931; Forero & Giraldo-Echeverry, 2015; Gil-Santana, Forero & Weirauch,

2015) and bees (Jackson, 1973; Wattanachaiyingcharoen & Jongjitvimol, 2007; Gil-Santana, 2008; Gil-Santana, Forero & Weirauch, 2015; Alvarez, Zamudio & Melo, 2019). Although assassin bugs reportedly coexist with model wasps or bees in the same microhabitats (Alvarez, Zamudio & Melo, 2019), the mimetic interactions between assassin bugs and other insects have not been experimentally tested using predators. In this study, we used the frog *Pe. nigromaculatus* as a potential predator to investigate the mimetic interaction between the assassin bug *S. flavipes* and the bombardier beetle *Ph. occipitalis jessoensis*. Although both species were well defended against predators, *S. flavipes* showed poorer defence than did *Ph. occipitalis jessoensis* (Figs. 4 and 8). This could be explained by the differences in body size between the two insect species (Table 1), as prey weight could influence predation success by the frog *Pe. nigromaculatus* (Sugiura, 2018).

The distribution of the assassin bug *S. flavipes* overlaps with the distribution of the bombardier beetle *Ph. occipitalis jessoensis* in East and Southeast Asia (Chlond, 2018; Fedorenko, 2021). However, *S. flavipes* is also found in South and West Asia where *Ph. occipitalis jessoensis* is not distributed (Chlond, 2018; Fedorenko, 2021). In the assassin bug *S. flavipes*, the body colour pattern of the South and West Asian populations partially differs from the body colour pattern of the East and Southeast Asian populations; the pronotum of the South and West Asian populations is redder than the pronotum of the East and Southeast Asian populations, although both types share the black and yellow pattern on other body parts (Chond, Bugaj-Nawrocka & Sawka-Gadek, 2019). Notably, the body colour pattern of South and West Asian *S. flavipes* is very similar to the body colour pattern of another bombardier beetle, *Pheropsophus (Stenaptinus) catoirei* (Dejean); adult *Ph. catoirei* individuals have a reddish head and pronotum (Fedorenko, 2021). *Pheropsophus catoirei*, which is closely related to *Ph. occipitalis jessoensis*, shares its distribution area (South and West Asia) with *S. flavipes* in East and Southeast Asia (Chond, Bugaj-Nawrocka & Sawka-Gadek, 2019; Fedorenko, 2021). Therefore, the mimetic partner of *S. flavipes* could differ between East–Southeast Asia and West–South Asia.

## CONCLUSIONS

Some aposematic species form ‘mimicry rings’ (Kunte, Kizhakke & Nawge, 2021; Chatelain et al., 2023). Mimicry rings are composed of at least two Müllerian co-mimics or one aposematic species plus one Batesian mimic (Kunte, Kizhakke & Nawge, 2021); the smallest mimicry rings include only two species (Kunte, Kizhakke & Nawge, 2021), while the largest mimicry rings include >100 species (Pekár et al., 2017). Although many studies have investigated mimicry rings that are composed of closely related taxa (Kunte, Kizhakke & Nawge, 2021), fewer studies have focused on mimicry complexes that involve distantly related taxa (Linsley, Esiner & Klots, 1961; Pekár et al., 2017). Our results suggest that the bombardier beetle *Ph. occipitalis jessoensis* and the distantly related *S. flavipes* form a multi-order mimetic complex. Other insects such as the rove beetle species *Ocypus weisei* Harold (Coleoptera: Staphylinidae) may be included in the mimetic complex; *O. weisei* adults share a similar microhabitat (on the ground in grassland), body colour pattern (black and yellow pattern), and movement pattern with *Ph. occipitalis jessoensis* and *S.*

*flavipes* adults in Japan. Furthermore, the black and yellow body colour pattern of these insects is similar to the typical aposematic colour of stinging hymenopteran insects, such as paper wasps and bees ([Chatelain et al., 2023](#)). The presence of a colour pattern similar to a typical aposematic pattern likely provides *S. flavipes* and *Ph. occipitalis jessoensis* with more robust protection from predators, compared with other colour patterns.

## ACKNOWLEDGEMENTS

We thank T Date for assisting with insect and frog maintenance. We also thank J Raška and anonymous reviewers for helpful comments on an earlier version of the manuscript.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This study was supported by a Grant-in-Aid for Scientific Research (JSPS KAKENHI Grant number 19K06073) and funds from the Hoshizaki Green Foundation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Grant-in-Aid for Scientific Research: 19K06073.

Hoshizaki Green Foundation.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Shinji Sugiura conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Masakazu Hayashi conceived and designed the experiments, prepared figures and/or tables, and approved the final draft.

### Animal Ethics

The following information was supplied relating to ethical approvals (*i.e.*, approving body and any reference numbers):

The experiments were performed in accordance with the Kobe University Animal Experimentation Regulations (Kobe University's Animal Care and Use Committee, No. 30–01).

### Field Study Permissions

The following information was supplied relating to field study approvals (*i.e.*, approving body and any reference numbers):

Our study was not conducted in any protected areas. Because the study insect and frog species were not protected, no specific permissions were required to collect these species in Japan.

## Data Availability

The following information was supplied regarding data availability:

The raw data are available at figshare: Sugiura, Shinji; Hayashi, Masakazu (2023): Data from: Bombardiers and assassins: mimetic interactions between unequally defended insects. figshare. Dataset. <https://doi.org/10.6084/m9.figshare.19636617>.

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.15380#supplemental-information>.

## REFERENCES

- Akino K, Sasaki M, Okamoto D. 1956.** Studies on mole-crickets attacking rice seeds directly sown without transplanting between rows of wheat and barley. *Bulletin of the Chugoku National Agricultural Experiment Station* **3**(1):91–110 (in Japanese with English résumé).
- Alvarez LJ, Zamudio F, Melo MC. 2019.** Eating with the enemy? Mimic complex between a stingless bee and assassin bugs. *Papéis Avulsos de Zoologia* **59**:e20195927 DOI [10.11606/1807-0205/2019.59.27](https://doi.org/10.11606/1807-0205/2019.59.27).
- Aneshansley DT, Eisner T, Widom JM, Widom B. 1969.** Biochemistry at 100 °C: explosive secretory discharge of bombardier beetles (*Brachinus*). *Science* **165**(3888):61–63 DOI [10.1126/science.165.3888.61](https://doi.org/10.1126/science.165.3888.61).
- Anichtchenko A, Choi JB, Facchini S, Marrero J, Panin R, Potanin D, Roguet D, Solodovnikov I, Will KW. 2022.** Carabidae of the World. Available at <https://carabidae.org/taxa/licinini-bonelli> (accessed on 23 February 2022).
- Arndt EM, Moore W, Lee WK, Ortiz C. 2015.** Mechanistic origins of bombardier beetle (*Brachinini*) explosion-induced defensive spray pulsation. *Science* **348**(6234):563–567 DOI [10.1126/science.1261166](https://doi.org/10.1126/science.1261166).
- Aubier TG, Joron M, Sherratt TN. 2017.** Mimicry among unequally defended prey should be mutualistic when predators sample optimally. *The American Naturalist* **189**(3):267–282 DOI [10.1086/690121](https://doi.org/10.1086/690121).
- Balogh ACV, Gamberale-Stille G, Leimar O. 2008.** Learning and the mimicry spectrum: from quasi-Bates to super-Müller. *Animal Behaviour* **76**(5):1591–1599 DOI [10.1016/j.anbehav.2008.07.017](https://doi.org/10.1016/j.anbehav.2008.07.017).
- Bates HW. 1862.** Contributions to an insect fauna of the Amazon valley. Lepidoptera: Heliconidae. *Transactions of the Linnean Society of London* **23**:495–566 DOI [10.1111/j.1096-3642.1860.tb00146.x](https://doi.org/10.1111/j.1096-3642.1860.tb00146.x).
- Bates D, Mächler M, Bolker B, Walker S. 2015.** Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**:1–48 DOI [10.18637/jss.v067.i01](https://doi.org/10.18637/jss.v067.i01).



- Bonacci T, Aloise G, Brandmayr P, Brandmayr TZ, Capula M. 2008.** Testing the predatory behaviour of *Podarcis sicula* (Reptilia: Lacertidae) towards aposematic and non-aposematic preys. *Amphibia-Reptilia* **29**:449–453 DOI [10.1163/156853808785111986](https://doi.org/10.1163/156853808785111986).
- Bonacci T, Brandmayr P, Brandmayr TZ. 2011.** Predator feeding choice on conspicuous and non-conspicuous carabid beetles: first results. *ZooKeys* **100**:171–179 DOI [10.3897/zookeys.100.1525](https://doi.org/10.3897/zookeys.100.1525).
- Chatelain P, Elias M, Fontaine C, Villemant C, Dajoz I, Perrard A. 2023.** Müllerian mimicry among bees and wasps: a review of current knowledge and future avenues of research. *Biological Reviews* DOI [10.1111/brv.12955](https://doi.org/10.1111/brv.12955).
- Chłond D. 2018.** A taxonomic revision of the genus *Sirthenia* (Hemiptera: Heteroptera: Reduviidae) of the Old World. *Zootaxa* **4520**(1):1–85 DOI [10.11646/zootaxa.4520.1.1](https://doi.org/10.11646/zootaxa.4520.1.1).
- Chłond D, Bugaj-Nawrocka A, Sawka-Gądek N. 2019.** Are we witnessing speciation? A case study of the species *Sirthenia flavipes* (Stål, 1855) (Heteroptera: Reduviidae). *Austral Entomology* **58**(1):96–110 DOI [10.1111/aen.12320](https://doi.org/10.1111/aen.12320).
- Chouteau M, Dezeure J, Sherratt TN, Llaurens V, Joron M. 2019.** Similar predator aversion for natural prey with diverse toxicity levels. *Animal Behaviour* **153**:49–59 DOI [10.1016/j.anbehav.2019.04.017](https://doi.org/10.1016/j.anbehav.2019.04.017).
- Dean J. 1979.** Defensive reaction time of bombardier beetles: an investigation of the speed of a chemical defense. *Journal of Chemical Ecology* **5**(5):691–701 DOI [10.1007/BF00986554](https://doi.org/10.1007/BF00986554).
- Dean J. 1980.** Encounters between bombardier beetles and two species of toads (*Bufo americanus*, *B. marinus*): speed of prey-capture does not determine success. *Journal of Comparative Physiology* **135**(1):41–50 DOI [10.1007/BF00660180](https://doi.org/10.1007/BF00660180).
- Edmunds M. 1974.** *Defense in animals*. Harlow: Longman.
- Eisner T. 1958.** The protective role of the spray mechanism of the bombardier beetle, *Brachynus ballistarius* Lec. *Journal of Insect Physiology* **2**(3):215–220 DOI [10.1016/0022-1910\(58\)90006-4](https://doi.org/10.1016/0022-1910(58)90006-4).
- Eisner T, Aneshansley DJ, Campo MLdel, Eisner M, Frank JH, Deyrup M. 2006.** Effect of bombardier beetle spray on a wolf spider: repellency and leg autotomy. *Chemoecology* **16**(4):185–189 DOI [10.1007/s00049-006-0346-8](https://doi.org/10.1007/s00049-006-0346-8).
- Eisner T, Dean J. 1976.** Ploy and counterploy in predator–prey interactions: orb-weaving spiders versus bombardier beetles. *Proceedings of the National Academy of Sciences of the United States of America* **73**(4):1365–1367 DOI [10.1073/pnas.73.4.1365](https://doi.org/10.1073/pnas.73.4.1365).
- Eisner T, Eisner M, Siegler M. 2005.** *Secret weapons: defenses of insects, spiders, scorpions, and other many-legged creatures*. Cambridge: The Belknap Press of the Harvard University Press.
- Eisner T, Meinwald J. 1966.** Defensive secretions of arthropods. *Science* **153**(1734):1341–1350 DOI [10.1126/science.153.3742.1341](https://doi.org/10.1126/science.153.3742.1341).
- Endler JA. 1991.** Interactions between predators and prey. In: Krebs JR, Davies NB, eds. *Behavioural ecology: an evolutionary approach*. London, Paris, Berlin, Vienna: Blackwell, 169–196.

- Fedorenko DN. 2021.** *Stenaptinus* (Coleoptera: Carabidae: Brachininae) of Vietnam. Note 3. *Russian Entomological Journal* **30**(3):252–263 DOI [10.15298/rusentj.30.3.02](https://doi.org/10.15298/rusentj.30.3.02).
- Forero D, Giraldo-Echeverry NG. 2015.** First record of the assassin bug genus *Coilopus* Elkins, 1969 (Hemiptera: Heteroptera: Reduviidae) from Colombia. *Check List* **11**(3):1634 DOI [10.15560/11.3.1634](https://doi.org/10.15560/11.3.1634).
- Fujisawa T, Lee CM, Ishii M. 2012.** Species diversity of ground beetle assemblages in the distinctive landscapes of the Yodo River flowing through northern Osaka Prefecture, central Japan. *Japanese Journal of Environmental Entomology and Zoology* **23**(2):89–100 DOI [10.11257/jjeez.23.89](https://doi.org/10.11257/jjeez.23.89).
- Gil-Santana HR. 2008.** New records, and nomenclatural and biological notes on Reduviidae (Hemiptera: Heteroptera) from Bolivia and Brazil. *Zootaxa* **1785**(1):43–53 DOI [10.11646/zootaxa.1785.1.2](https://doi.org/10.11646/zootaxa.1785.1.2).
- Gil-Santana HR, Forero D, Weirauch C. 2015.** Assassin bugs (Reduviidae excluding Triatominae). In: Pannizi AR, Grazia J, eds. *True Bugs (Heteroptera) of the Neotropics*. Dordrecht, Heidelberg, New York, London: Springer, 307–351 DOI [10.1007/978-94-017-9861-7\\_12](https://doi.org/10.1007/978-94-017-9861-7_12).
- Habu A, Sadanaga K. 1965.** Illustrations for identification of larvae of the Carabidae found in cultivated fields and paddy-fields (III). *Bulletin of the National Institute of Agricultural Sciences, Series C: Plant Pathology and Entomology* **19**:81–216 (in Japanese with English summary).
- Haviland MD. 1931.** The Reduviidae of Kartabo Bartica District, British Guiana. *Zoologica* **7**(5):129–154.
- Hayashi M. 2023.** Life history of an assassin bug, *Sirthenia flavipes* (Stål, 1855): laboratory rearing and field observations. *Special Bulletin of the Hoshizaki Green Foundation* **32**:51–62.
- Hirai T. 2002.** Ontogenetic change in the diet of the pond frog, *Rana nigromaculata*. *Ecological Research* **17**(6):639–644 DOI [10.1046/j.1440-1703.2002.00521.x](https://doi.org/10.1046/j.1440-1703.2002.00521.x).
- Hirai T, Matsui M. 1999.** Feeding habits of the pond frog, *Rana nigromaculata*, inhabiting rice fields in Kyoto, Japan. *Copeia* **1999**(4):940–947 DOI [10.2307/1447969](https://doi.org/10.2307/1447969).
- Hirashima Y, Morimoto K. 2008.** *Iconographia insectorum Japonicorum colore naturali edita III*. Tokyo: Hokuryukan(in Japanese).
- Honma A, Oku S, Nishida T. 2006.** Adaptive significance of death feigning posture as a specialized inducible defence against gape-limited predators. *Proceedings of the Royal Society B* **273**(1594):1631–1636 DOI [10.1098/rspb.2006.3501](https://doi.org/10.1098/rspb.2006.3501).
- Honma A, Takakura K, Nishida T. 2008.** Optimal-foraging predator favors commensalistic Batesian mimicry. *PLOS ONE* **3**(10):e3411 DOI [10.1371/journal.pone.0003411](https://doi.org/10.1371/journal.pone.0003411).
- Hudson WG. 1987.** Ontogeny of prey selection in *Sirthenia carinata*: generalist juveniles become specialist adults. *Entomophaga* **32**(4):399–406 DOI [10.1007/BF02372449](https://doi.org/10.1007/BF02372449).
- Ihalainen E, Lindström L, Mappes J. 2007.** Investigating Müllerian mimicry: predator learning and variation in prey defences. *Journal of Evolutionary Biology* **20**(2):780–791 DOI [10.1111/j.1420-9101.2006.01234.x](https://doi.org/10.1111/j.1420-9101.2006.01234.x).

- Ihalainen E, Lindström L, Mappes J, Puolakkainen S. 2008. Butterfly effects in mimicry? Combining signal and taste can twist the relationship of Müllerian co-mimics. *Behavioral Ecology and Sociobiology* 62(8):1267–1276 DOI 10.1007/s00265-008-0555-y.
- Ishitani M, Yano K. 1994. Species composition and seasonal activities of ground beetles (Coleoptera) in a fig orchard. *Japanese Journal of Entomology* 62(1):201–210.
- Ito F, Taniguchi K, Billen J. 2016. Defensive function of petiole spines in queens and workers of the formicine ant *Polyrhachis lamellidens* (Hymenoptera: Formicidae) against an ant predator, the Japanese tree frog *Hyla japonica*. *Asian Myrmecology* 8:81–86 DOI 10.20362/am.008014.
- Ito S, Okutani T, Hiura I. 1977. *Colored illustrations of the insect of Japan II*. Osaka: Hoikusha (in Japanese).
- Jackson JF. 1973. Mimicry of *Trigona* bees by a reduviid (Hemiptera) from British Honduras. *The Florida Entomologist* 56(3):200–202 DOI 10.2307/3493124.
- Kanehisa K. 1996. Secretion of defensive substance by Carabidae and Brachinidae. *Bulletin of the Research Institute for Bioresources, Okayama University* 4(1):9–23 (in Japanese with English summary).
- Kanehisa K, Murase M. 1977. Comparative study of the pygidial defensive systems of carabid beetles. *Applied Entomology and Zoology* 12(3):225–235 DOI 10.1303/aez.12.225.
- Kojima W, Yamamoto R. 2020. Defense of bombardier beetles against avian predators. *The Science of Nature* 107:36 DOI 10.1007/s00114-020-01692-z.
- Komaki S, Igawa T, Lin SM, Tojo K, Min MS, Sumida M. 2015. Robust molecular phylogeny and palaeodistribution modelling resolve a complex evolutionary history: glacial cycling drove recurrent mtDNA introgression among *Pelophylax* frogs in East Asia. *Journal of Biogeography* 42(11):2159–2171 DOI 10.1111/jbi.12584.
- Kunte K, Kizhakke AG, Nawge V. 2021. Evolution of mimicry rings as a window into community dynamics. *Annual Review of Ecology, Evolution, and Systematics* 52:315–341 DOI 10.1146/annurev-ecolsys-012021-024616.
- Lindström L, Lyytinen A, Mappes J, Ojala K. 2006. Relative importance of taste and visual appearance for predator education in Müllerian mimicry. *Animal Behaviour* 72(2):323–333 DOI 10.1016/j.anbehav.2005.10.015.
- Linsley EG, Esiner T, Klots AB. 1961. Mimetic assemblages of sibling species of lycid beetles. *Evolution* 15(1):15–29 DOI 10.1111/j.1558-5646.1961.tb03126.x.
- Louis D. 1974. Biology of Reduviidae of cocoa farms in Ghana. *The American Midland Naturalist* 91(1):68–89 DOI 10.2307/2424512.
- Maldonado Capriles J, Lozada Robles PW. 1992. Key to the group of neotropical wasp-mimetic harpactorine genera and the description of a new species (Hemiptera: Reduviidae). *Proceedings of the Entomological Society of Washington* 94(1):162–165.
- Marples NM. 1993. Toxicity assays of ladybirds using natural predators. *Chemoecology* 4(1):33–38 DOI 10.1007/BF01245894.

- Marples NM, Brakefield PM, Cowie RJ. 1989. Differences between the 7-spot and 2-spot ladybird beetles (Coccinellidae) in their toxic effects on a bird predator. *Ecological Entomology* 14(1):79–84 DOI 10.1111/j.1365-2311.1989.tb00756.x.
- Matsubara S, Sugiura S. 2017. Chemical defence of turnip sawfly larvae against Japanese tree frogs. *Journal of Asia-Pacific Entomology* 20(1):225–227 DOI 10.1016/j.aspen.2017.01.001.
- Matsui M, Maeda N. 2018. *Encyclopedia of Japanese frogs*. Tokyo: Bun-ichi Sogo Shuppan.
- Müller F. 1878. Über die vorthteile der mimicry bei schmetterlingen. *Zoologischer Anzeiger* 1:54–55.
- Müller F. 1879. Ituna and Thyridia: a remarkable case of mimicry in butterflies. *Transactions of the Entomological Society of London* 1879:20–29.
- Ohwaki A, Kaneko Y, Ikeda H. 2015. Seasonal variability in the response of ground beetles (Coleoptera: Carabidae) to a forest edge in a heterogeneous agricultural landscape in Japan. *European Journal of Entomology* 112(1):135–144 DOI 10.14411/eje.2015.022.
- Pekár S, Petráková L, Bulbert MW, Whiting MJ, Herberstein ME. 2017. The golden mimicry complex uses a wide spectrum of defence to deter a community of predators. *eLife* 6:e22089 DOI 10.7554/eLife.22089.
- R Core Team. 2018. *R, a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing.
- Raška J, Krajíček J, Bosáková Z, Štys P, Exnerová A. 2020. Larvae of pyrrhocorid true bugs are not to spiders' taste: putative Müllerian mimicry. *Biological Journal of the Linnean Society* 129(1):199–212 DOI 10.1093/biolinlean/blz174.
- Readio PA. 1927. Studies on the biology of the Reduviidae of America North of Mexico. *The University of Kansas Science Bulletin* 17(1):5–291.
- Rowland HM, Ihalainen E, Lindström L, Mappes J, Speed MP. 2007. Co-mimics have a mutualistic relationship despite unequal defences. *Nature* 448(7149):64–67 DOI 10.1038/nature05899.
- Rowland HM, Mappes J, Ruxton GD, Speed MP. 2010. Mimicry between unequally defended prey can be parasitic: evidence for quasi-Batesian mimicry. *Ecology Letters* 13(12):1494–1502 DOI 10.1111/j.1461-0248.2010.01539.x.
- Ruxton GD, Franks DW, Balogh ACV, Leimar O. 2008. Evolutionary implications of the form of predator generalization for aposematic signals and mimicry in prey. *Evolution* 62(11):2913–2921 DOI 10.1111/j.1558-5646.2008.00485.x.
- Ruxton GD, Sherratt TN, Speed MP. 2004. *Avoiding attack: the evolutionary ecology of crypsis, aposematism, and mimicry*. Oxford: Oxford University Press.
- Sano M, Shinohara M. 2012. Species comparison of frogs food habits during mating seasons in Uenohara, Yamanashi Pref. Japan. *Bulletin of Teikyo University of Science and Technology* 8:101–111 (in Japanese with English summary).

- Sarashina M, Yoshihisa Y, Yoshida T. 2011. Stomach contents of invasive Black-spotted Pond frog (*Rana nigromaculata*) in urban landscape of Sapporo City. *Journal of Rakuno Gakuen University* 36(1):81–86 (in Japanese with English summary).
- Schaller JC, Davidowitz G, Papaj DR, Smith RL, Carrière Y, Moore W. 2018. Molecular phylogeny, ecology and multispecies aggregation behaviour of bombardier beetles in Arizona. *PLOS ONE* 13(10):e0205192 DOI 10.1371/journal.pone.0205192.
- Schmidt JO. 2009. Defensive behavior. In: Resh VH, Cardé RT, eds. *Encyclopedia of Insects*. Second Edition. Burlington, San Diego, London: Elsevier, 252–257.
- Shelford MA. 1902. Observations on some mimetic insects and spiders from Borneo and Singapore. *Proceedings of the Zoological Society of London* 2:230–284.
- Sherratt TN. 2008. The evolution of Müllerian mimicry. *Naturwissenschaften* 95(8):681–695 DOI 10.1007/s00114-008-0403-y.
- Shinohara T, Takami Y. 2020. Functional diversity and trade-offs in divergent antipredator morphologies in herbivorous insects. *Ecology and Evolution* 10(11):5089–5096 DOI 10.1002/ece3.6262.
- Skelhorn J, Rowe C. 2006. Taste-rejection by predators and the evolution of unpalatability in prey. *Behavioral Ecology and Sociobiology* 60(4):550–555 DOI 10.1007/s00265-006-0199-8.
- Soukupová TS, Veselý P, Fuchs R. 2021. Wariness to colour patterns: birds versus European red-and-black insects. *Ecological Entomology* 46(5):1157–1164 DOI 10.1111/een.13060.
- Speed MP. 1993. Muellierian mimicry and psychology of predation. *Animal Behaviour* 45(3):571–580 DOI 10.1006/anbe.1993.1067.
- Speed MP. 1999. Batesian, quasi-Batesian or Müllerian mimicry? Theory and data in mimicry research. *Evolutionary Ecology* 13(7–8):755–776 DOI 10.1023/A:1010871106763.
- Speed MP, Alderson NJ, Hardman C, Ruxton GD. 2000. Testing Müllerian mimicry: an experiment with wild birds. *Proceedings of the Royal Society of London B* 267(1444):725–731 DOI 10.1098/rspb.2000.1063.
- Speed MP, Turner JRG. 1999. Learning and memory in mimicry: II. Do we understand the mimicry spectrum? *Biological Journal of the Linnean Society* 67(3):281–312 DOI 10.1111/j.1095-8312.1999.tb01935.x.
- Staddon BW. 1979. The scent glands of Heteroptera. *Advances in Insect Physiology* 14:351–418 DOI 10.1016/S0065-2806(08)60055-7.
- Sugiura S. 2018. Anti-predator defences of a bombardier beetle: is bombing essential for successful escape from frogs? *PeerJ* 6:e5942 DOI 10.7717/peerj.5942.
- Sugiura S. 2020a. Predators as drivers of insect defenses. *Entomological Science* 23(3):316–337 DOI 10.1111/ens.12423.
- Sugiura S. 2020b. Active escape of prey from predator vent via the digestive tract. *Current Biology* 30(15):R867–R868 DOI 10.1016/j.cub.2020.06.026.
- Sugiura S. 2021. Beetle bombing always deters praying mantises. *PeerJ* 9:e11657 DOI 10.7717/peerj.11657.



- Sugiura S, Date T. 2022. Bombardier beetles repel invasive bullfrogs. *PeerJ* 10:e13805 DOI 10.7717/peerj.13805.
- Sugiura S, Sato T. 2018. Successful escape of bombardier beetles from predator digestive systems. *Biology Letters* 14(2):20170647 DOI 10.1098/rsbl.2017.0647.
- Sugiura S, Tsujii M. 2022. Male wasp genitalia as an anti-predator defense. *Current Biology* 32(24):R1336–R1337 DOI 10.1016/j.cub.2022.11.030.
- Takahashi T. 1996. Reduviidae and Nabidae in Hyogo Prefecture (2). *Parnassius* 43:3–10 (in Japanese).
- Takara T. 1957. Provisional list of Hemiptera (Heteroptera) in the Ryukyu Islands. *Science Bulletin of Agriculture & Home Economics Division, University of the Ryukyus* 4:11–90.
- Taniguchi K, Maruyama M, Ichikawa T, Ito F. 2005. A case of Batesian mimicry between myrmecophilous staphylinid beetle, *Pella comes*, and its host ant, (*Lasius Dendrolasius*) *spathepus*: an experiment using the Japanese tree frog *Hyla japonica* as a real predator. *Insectes Sociaux* 52(4):320–322 DOI 10.1007/s00040-005-0813-1.
- Tomokuni M, Yasunaga T, Takai M, Yamashita I, Kawamura M, Kawasaki T. 1993. *A field guide to Japanese bugs—Terrestrial Heteropterans*. Tokyo: Zenkoku Noson Kyoiku Kyokai(in Japanese).
- Walker AA, Mayhew ML, Jin J, Herzig V, Undheim EAB, Sombke A, Fry BG, Meritt DJ, King GF. 2018. The assassin bug *Pristhesancus plagipennis* produces two distinct venoms in separate gland lumens. *Nature Communications* 9:755 DOI 10.1038/s41467-018-03091-5.
- Walker AA, Robinson SD, Undheim EAB, Jin J, Han X, Fry BG, Vetter I, King GF. 2019. Missiles of mass disruption: composition and glandular origin of venom used as a projectile defensive weapon by the assassin bug *Platymeris rhadamanthus*. *Toxins* 11(11):673 DOI 10.3390/toxins11110673.
- Walker AA, Weirauch C, Fry BG, King GF. 2016. Venoms of heteropteran insects: a treasure trove of diverse pharmacological toolkits. *Toxins* 8(2):43 DOI 10.3390/toxins8020043.
- Wattanachaiyingcharoen W, Jongjitvimol T. 2007. First record of the predator, *Pahabengkakia piliceps* Miller, 1941 (Reduviidae, Harpactorinae) in the stingless bee, *Trigona collina* Smith, 1857 (Apidae, Meliponinae) in Thailand. *The Natural History Journal of Chulalongkorn University* 7(1):71–74.
- Willemse L. 1985. A taxonomic revision of the New World species of *Sirthena* (Heteroptera: Reduviidae: Peiratinae). *Zoologische Verhandelingen* 215(1):3–67.
- Williams BL, Hanifin CT, Brodie Jr ED, Brodie III ED. 2010. Tetrodotoxin affects survival probability of rough-skinned newts (*Taricha granulosa*) faced with TTX-resistant garter snake predators (*Thamnophis sirtalis*). *Chemoecology* 20(4):285–290 DOI 10.1007/s00049-010-0057-z.

- Winters AE, Wilson NG, Van den Berg CP, How MJ, Endler JA, Marshall NJ, White AM, Garson MJ, Cheney KL. 2018.** Toxicity and taste: unequal chemical defences in a mimicry ring. *Proceedings of the Royal Society B* **285**(1880):20180457 DOI [10.1098/rspb.2018.0457](https://doi.org/10.1098/rspb.2018.0457).
- Yahiro K, Fujimoto T, Tokuda M, Yano K. 1992.** Species composition and seasonal abundance of ground beetles (Coleoptera) in paddy fields. *Japanese Journal of Entomology* **60**(4):805–813.
- Yasunaga T, Maehara S, Ishikawa T, Takai M. 2018.** *Guidebook to the heteropteran world—Basic ecology, morphology, classification and research methodology*. Tokyo: Zenkoku Noson Kyoiku Kyokai(in Japanese).
- Zhang G, Weirauch C. 2014.** Molecular phylogeny of Harpactorini (Insecta: Reduviidae): correlation of novel predation strategy with accelerated evolution of predatory leg morphology. *Cladistics* **30**(4):339–351 DOI [10.1111/cla.12049](https://doi.org/10.1111/cla.12049).