



# Metabolic and ventilatory changes during postural change from the supine position to the reclining position in bedridden older patients

山田, 洋二

---

(Degree)

博士 (保健学)

(Date of Degree)

2023-09-25

(Date of Publication)

2024-09-01

(Resource Type)

doctoral thesis

(Report Number)

甲第8736号

(URL)

<https://hdl.handle.net/20.500.14094/0100485920>

※ 当コンテンツは神戸大学の学術成果です。無断複製・不正使用等を禁じます。著作権法で認められている範囲内で、適切にご利用ください。



# 博 士 論 文

Metabolic and ventilatory changes during postural change from the supine position to the reclining position in bedridden older patients

(寝たきり高齢者における背臥位から座位への姿勢変化時の代謝・換気変化)

令和5年7月7日

神戸大学大学院保健学研究科保健学専攻

Yoji YAMADA 山田 洋二

## **Metabolic and ventilatory changes during postural change from the supine position to the reclining position in bedridden older patients**

Yoji Yamada MS<sup>1,2</sup>, Yuji Mitani PhD<sup>2</sup>, Akio Yamamoto PhD<sup>2</sup>, Kazumo Miura MS<sup>2</sup>, Kanji Yamada PhD<sup>2</sup>, Yukari Oki PhD<sup>2</sup>, Yutaro Oki PhD<sup>2</sup>, Yasumichi Maejima MS<sup>3</sup>, Yoko Kurumatani MD<sup>4</sup>, and Akira Ishikawa PhD<sup>2</sup>

<sup>1</sup> Department of Rehabilitation, Isawa Kyoritsu Hospital, Fuefuki, Yamanashi, Japan

<sup>2</sup> Department of Public Health, Kobe University Graduate School of Health Sciences, Kobe, Hyogo, Japan

<sup>3</sup> Department of Rehabilitation, Kofu Kyoritsu Hospital, Kofu, Yamanashi, Japan

<sup>4</sup> Department of cardiology, Kofu Kyoritsu Hospital, Kofu, Yamanashi, Japan

### **Abstract**

**Background:** The prevention of pneumonia in bedridden older patients is important, and its recurrence in these patients is a relevant issue. Patients who are bedridden and inactive, and have dysphagia are considered to be at risk for pneumonia. Efforts to reduce the bedridden state and low activity may be necessary to reduce the risk of developing pneumonia in bedridden older patients. This study aimed to clarify the effects of postural change from the supine position to the reclining position on metabolic and ventilatory parameters and on safety in bedridden older patients.

**Methods:** Using a breath gas analyzer and other tools, we assessed the following three positions: lying on the back (supine), resting in the Fowler position (Fowler), and resting in an 80° recline wheelchair (80°). Measurements were oxygen uptake ( $\dot{V}O_2$ ), carbon dioxide output ( $\dot{V}CO_2$ ), gas exchange ratio, tidal volume ( $V_T$ ), minute volume, respiratory rate, inspiratory time, expiratory time, total respiratory time, mean inspiratory flow ( $V_T/Ti$ ), metabolic equivalents, end-expiratory oxygen, and end-expiratory carbon dioxide as well as various vital signs.

**Results:** The study analysis included 19 bedridden participants. The change in  $\dot{V}O_2$  driven by changing the posture from the supine position to the Fowler position was as small as 10.8 mL/min.  $V_T$  significantly increased from the supine position ( $398.4 \pm 111.2$  mL) to the Fowler position ( $426.9 \pm 106.8$  mL) ( $P = .037$ ) and then showed a decreasing trend in the 80° position ( $416.8 \pm 92.5$  mL).

**Conclusions:** For bedridden older patients, sitting in a wheelchair is a very low-impact physical

activity, similar to that in normal people. The  $V_T$  of bedridden older patients was maximal in the Fowler position, and the ventilatory volume did not increase with an increasing reclining angle, unlike that in normal people. These findings suggest that appropriate reclining postures in clinical situations can promote an increase in the ventilatory rate in bedridden older patients.

**Abbreviation list:**

$\dot{V}O_2$  = oxygen uptake,  $\dot{V}CO_2$  = carbon dioxide output,  $V_T$  = tidal volume,  $VE$  = minute volume,  $V_T/Ti$  = mean inspiratory flow, ADL = activities of daily living, FIM = Functional Independence Measure, MASA = Mann Assessment of Swallowing Ability

## Introduction

The prevention of pneumonia in bedridden older patients is important. However, rehabilitation for these patients has not yet been established. Pneumonia is the fourth leading cause of deaths in Japan (<https://www.mhlw.go.jp/toukei/itiran/eiyaku.html>, Accessed September 5, 2022), and 80.1% of patients with pneumonia aged >70 years are diagnosed with aspiration pneumonia.<sup>1</sup> Low activities of daily living (ADLs)<sup>2</sup> are reportedly associated with the development of pneumonia. To prevent the development of pneumonia, vaccination and oral care are strongly recommended.<sup>3</sup> Notably, early rehabilitation for patients with pneumonia has been reported to be effective.<sup>4</sup> In contrast, methods of rehabilitation for preventing the onset of pneumonia in older people have not yet been established.<sup>5</sup> Older patients who are bedridden and inactive and have dysphagia are considered to be at risk for pneumonia. Some previous studies have reported bedridden state, weight loss, gait disturbance, dysphagia, and presence of aspiration, impaired swallowing function, dehydration, and dementia as risk factors for aspiration pneumonia.<sup>6</sup> Recently, Functional Independence Measure (FIM) motor item scores of <20 points on ADLs scale and Mann Assessment of Swallowing Ability (MASA) scores of <171 points, indicating swallowing function, were reported to be associated with an increased the risk of pneumonia.<sup>5</sup> Based on the above evidence, efforts to reduce the bedridden state and reduce low activity may be necessary to reduce the risk of developing pneumonia in bedridden older patients.

The change of posture from the supine position to sitting upright in a wheelchair is expected to change the ventilation status. In clinical practice, for patients with mobility disorder, maintaining wheelchair sitting and reducing a bedridden state can be important rehabilitation approaches that can help with postural drainage.<sup>7</sup> It has also been previously reported that sitting upright increases functional residual capacity and prevents alveolar collapse.<sup>8</sup> Furthermore, tidal volume ( $V_T$ ) and minute volume ( $VE$ ) increased in the sitting posture without changing oxygen uptake ( $\dot{V}O_2$ ) compared with the supine posture in individuals who could sit or stand.<sup>9,10</sup> Increasing  $V_T$  and the expiratory flow rate is useful for sputum expectoration.<sup>11</sup> However, the effects of changes in the wheelchair sitting posture on metabolic and ventilatory status in bedridden older patients are not clear. Furthermore, as many bedridden older people have several comorbidities as well as communication disorders, it is necessary to examine whether the effects of changes in sitting posture on breathing and metabolism are within a safe range.

Therefore, this study aimed to clarify the safety and effects of postural change from the supine position to the reclining position on metabolic and ventilatory parameters in bedridden older

patients. By clarifying the effects of postural change in bedridden older patients, we expect to obtain useful information for identifying the optimal physical activity to prevent pneumonia.

## **Methods**

### **Ethical consideration**

This study was approved by the Ethics Committee of Kofu Kyoritsu Hospital (approval no.: 2018-2) and the Ethics Committee of Health Sciences of Kobe University Graduate School of Health Sciences (no.: 961). Before the study, explanations about the study were given to the participants and their family members, and written consent was obtained.

### **Participants**

Twenty patients who were hospitalized and treated in an acute care hospital between May and October 2018 participated in this study. The inclusion criteria were as follows: (1) being older, requiring nursing care, and having an FIM motor score of less than 20 points; (2) having cerebrovascular disease as the leading cause of a bedridden state (at least 6 months have passed since the onset of the disease); (3) not having chronic respiratory failure or chronic heart failure; (4) not having a fever (temperature less than 37.5°C) and having a stable medical condition; (5) having less than 14 hours of sitting time in a wheelchair per week<sup>12</sup>; and (6) having rehabilitation prescribed by a physician and having practiced sitting on the edge of a bed for at least 10 minutes with a physical therapist or others, during which the vital signs remained stable.

### **Measurement procedure**

Participants were placed in the following three positions: lying on the back (supine), resting in the Fowler position with a 40-degree recline and 20-degree tilt-up (Fowler), and resting in an 80-degree recline wheelchair (80°). Data on respiration and circulation were continuously obtained. The experiment was conducted at a fixed time more than 2 hours after eating for participants who took food orally or enterally. The participants were placed in the supine position for at least 10 minutes before the measurement. After checking heart rate (HR), transcutaneous oxygen saturation (SpO<sub>2</sub>), and blood pressure (BP), the participants wore a face mask and a HR sensor. We confirmed that there was no air leakage when the face mask was put on, and we instructed the participants to breathe spontaneously and not to speak or make any sound during the measurement. The participants were then seated in a reclining wheelchair in a relaxed position to allow full

respiratory function. The trunk was placed in a straight line along the back of the wheelchair, and the lower limbs were placed on a footrest. Measurements were taken with the head and neck in a 30-degree head and neck flexion position (slight flexion) as a pillow was used to stabilize the head and neck.

Measurements in each position were taken consecutively in the order of supine, Fowler, and 80°. These conditions were set up to approximate those commonly seen in rehabilitation clinical practice, in which patients are intervened in the supine position, then moved to the head-up position, and finally released in the sitting position (Fig. 1).

Measurements were taken for 6 minutes in each position, and data were recorded every 5 seconds.

### **Apparatus**

Exhaled gases were measured by the breath-by-breath method using a breath gas analyzer (Aeromonitor AE-300S; Minato Medical Science Co, Ltd, Osaka, Japan). Calibration of the gas sensor, volume transducer, and barometric pressure sensor was performed before each measurement. Exhaled gases were automatically collected via a face mask and a sample tube. The indices obtained from the expiratory gas measurements were  $\dot{V}O_2$ ,  $\dot{V}O_2$  divided by weight ( $\dot{V}O_2/W$ ), carbon dioxide output ( $\dot{V}CO_2$ ), gas exchange ratio (R),  $V_T$ , VE, respiratory rate (RR), inspiratory time (Ti), expiratory time (Te), total respiratory time (Ttot), mean inspiratory flow ( $V_T/Ti$ ), metabolic equivalents, end-expiratory oxygen, and end-expiratory carbon dioxide.

HR and BP were measured using a biometric monitor (WEP4204; Nhon Kohden, Tokyo, Japan). A pulse oximeter (3100WristOx; Nonin, Minnesota, USA) was used for transcutaneous oxygen saturation ( $SpO_2$ ) and pulse rate.

### **Participants' characteristics**

The data on age, gender, body mass index (BMI), name of the disease at admission, medications, FIM scores, MASA scores, biochemical findings (serum albumin [Alb], C-reactive protein [CRP], blood urea nitrogen, estimated glomerular filtration rate, and creatinine) were obtained from medical records.

### **Statistical analysis**

Average values obtained from 3 to 6 minutes (180–360 seconds) in each of the three experimental conditions (defined as the steady-state period) were used for analysis. For  $\dot{V}O_2$  and  $V_T$ ,  $\Delta\dot{V}O_2$

(Fowler) and  $\Delta V_T$  (Fowler) were calculated as the change from the supine position to the Fowler position, and  $\Delta \dot{V}O_2$  (80°) and  $\Delta V_T$  (80°) were calculated as the change from the supine position to the 80° position.

The Kolmogorov–Smirnov test was used to analyze the normality of each variable. Multiple comparisons were performed between the three positions of supine, Fowler, and 80° using Bonferroni's correction. Pearson's product-moment correlation coefficient was used to examine the relationship between metabolic and ventilatory parameters ( $\dot{V}O_2$  and  $V_T$ ). In the post-hoc power analysis, all variables had a power greater than 0.8. For statistical analysis, EZR (version 1.32) was used for comparisons between positions, and GPower 3.1 software was used for power analysis. The results for continuous variables are presented as mean  $\pm$  standard deviation, and the results for nominal variables are presented as number of persons (%). Two-tailed  $P < 0.05$  was considered statistically significance.

## Results

Of the 20 participants who met the inclusion criteria and participated in the experiment, 19 were included in the analysis after excluding one participant who spoke frequently during the experiment. The demographic data of the analyzed participants are shown in Table 1. The mean age of the participants was  $83.2 \pm 8.4$  years, and the mean BMI was  $17.3 \pm 3.3$  kg/m<sup>2</sup>. Of the 19 participants, 12 (63.2%) were male. Moreover, 15 (78.9%) participants had aspiration pneumonitis and bacterial pneumonia diseases. The mean Alb level was  $2.8 \pm 0.9$  g/dL, which was lower than the reference value (4.1–5.1 g/dL), and the mean CRP level was  $2.1 \pm 2.4$  mg/dL, which was higher than the reference value ( $\leq 0.30$  mg/dL).

The measured data obtained under each condition and the results of multiple comparisons are shown in Table 2. In the post-hoc power analysis, all variables demonstrated a power of  $>0.8$ . There were no significant differences in HR, and SpO<sub>2</sub> did not change among the three positions.  $\dot{V}O_2$  increased by 10.8 mL/min from the supine position ( $114.6 \pm 31.6$  mL/min) to the Fowler position ( $125.4 \pm 27.1$  mL/min) and decreased by 1.4 mL/min from the Fowler position to the 80° position ( $124 \pm 22.9$  mL/min). No significant difference in  $\dot{V}O_2/W$  was observed among the three positions. The R decreased with an increase in the reclining angle from  $0.94 \pm 0.05$  in the supine position to  $0.92 \pm 0.05$  in the 80° position, showing a significantly lower value in the 80° position than in the supine position ( $P = .01$ ).

$V_T$  significantly increased from the supine position ( $398.4 \pm 111.2$  mL) to the Fowler position



( $426.9 \pm 106.8$  mL) ( $P = .037$ ) and showed a decreasing trend from the Fowler position to the  $80^\circ$  position ( $416.8 \pm 92.5$  mL). The RR and minute ventilation rate tended to increase in both the Fowler and  $80^\circ$  positions, whereas  $T_i$  and  $T_{tot}$  tended to decrease with postural change.  $V_T/T_i$  was significantly higher in the supine and Fowler positions than in the  $80^\circ$  position ( $P = .02$ ).

We analyzed the relationship between the changes in  $V_T$  and  $\dot{V}O_2$  by changing the posture from supine to Fowler and  $80^\circ$ . Fig. 2A presents a scatter plot of  $\Delta\dot{V}O_2$  (Fowler) and  $\Delta V_T$  (Fowler) calculated as the change from the supine position to the Fowler position. Fig. 2B presents a scatter plot of  $\Delta\dot{V}O_2$  ( $80^\circ$ ) and  $\Delta V_T$  ( $80^\circ$ ) calculated as the change from the supine position to the  $80^\circ$  position. A significant positive correlation was found between  $\Delta\dot{V}O_2$  (Fowler) and  $\Delta V_T$  (Fowler) ( $r = 0.864$ , 95% Confidence interval [CI] [0.675–0.947],  $P < 0.001$ ). Similarly, there was a significant positive correlation between  $\Delta\dot{V}O_2$  ( $80^\circ$ ) and  $\Delta V_T$  ( $80^\circ$ ) ( $r = 0.781$ , 95% CI [0.507–0.912],  $P < 0.001$ ).

## Discussion

The purpose of the present study was to clarify the effects of postural change from the supine to reclining position on metabolic and ventilatory parameters in bedridden older patients. The following three findings were obtained in this study: 1) for bedridden older patients, the change in  $\dot{V}O_2$  driven by changing the posture from the supine position to the Fowler position was as small as 10.8 mL/min, which corresponds to an extremely low-intensity physical activity, 2) the ventilatory volume slightly increased from the supine position to the Fowler position, and 3) although the increase in  $\Delta\dot{V}O_2$  by changing posture was small, an increase in  $\Delta V_T$  was found, and there was a significant positive correlation with the increase in between  $\Delta\dot{V}O_2$  (Fowler) and  $\Delta V_T$  (Fowler), and between  $\Delta\dot{V}O_2$  ( $80^\circ$ ) and  $\Delta V_T$  ( $80^\circ$ ). These results indicated that elevating the trunk from the supine position in a bed enhances oxygen uptake as well as the volume of single ventilation.

Wheelchair seating is a very low-impact physical activity for bedridden older patients as well as for healthy people. In a study that set 70-degree tilt-up sitting for healthy participants,  $\dot{V}O_2$  increased by 10 mL/min from the supine position.<sup>10</sup> In a study with acute stroke patients, there was a 10 mL/min decrease in  $\dot{V}O_2$  with a 30-degree head-up from the supine position, a 10 mL/min increase in the sitting position, and a 140 mL/min increase in the standing position, indicating that the physical load is high in the standing position.<sup>13</sup> In the present study, the change in  $\dot{V}O_2$  was found to be 10.8 mL/min, an increase similar to that in healthy people, when sitting upright during the acute phase of stroke. This increase was small and was not considered as a clinically meaningful change.

$\dot{V}O_2$  increases with increasing exercise load, and its determinants are cardiac output, HR, and arteriovenous oxygenation rate.<sup>14</sup> In the reclining wheelchair posture used in the present study, the increase in  $\dot{V}O_2$ , an index of metabolic rate, may have been small because the reclining wheelchair posture does not require the activity of antigravity muscles and the oxygen demand of skeletal muscles is not increased. Furthermore, the wheelchair seating positions used in this experiment (the Fowler position and 80° position) did not cause significant changes in the participants' circulatory dynamics, as the changes in HR and SpO<sub>2</sub> were minimal. Therefore, for older bedridden patients, sitting in a wheelchair is indicated as a very low-intensity, safe physical activity demonstrating no remarkable changes in vital signs.

$V_T$  in the bedridden older patients was maximal in the Fowler position and differed from that in normal people. During postural change from the supine position to the Fowler position,  $V_T$  increased by 28.5 mL ( $P = 0.037$ ) and the  $V_T/Ti$  increased by 26.5 mL/s ( $P = 0.02$ ). Both values were significantly higher than the supine position values. This is consistent with the findings in a previous study that showed an increase in lung volume at 45° and 60° bed angles in patients with ARDS.<sup>15</sup> The  $V_T/Ti$  (mL/s) is also recognized as an indicator of ventilatory drive,<sup>16-18</sup> and it increases with increasing exercise load.<sup>19</sup> Furthermore, the correlation analysis showed a significant positive correlation between  $\Delta\dot{V}O_2$  and  $\Delta V_T$ , suggesting that postural change to the reclining sitting position can increase  $V_T$ , a ventilation index, and promote ventilatory drive, even though it is a low-intensity activity for bedridden older patients. However, in normal people, it has been reported that ventilation increases as the backrest tilt increases from the supine position.<sup>10,20</sup> In the present study,  $V_T$  was maximal in the Fowler position and was mildly decreased in the 80° position, indicating that  $V_T$  in bedridden older patients is different from that in normal people.

Ventilation did not increase with an increase in the reclining angle. The reason for this result may be the background specific to bedridden older patients. In general, chest wall motion in the sitting posture is associated with less gravitational compression of the thorax, higher thoracic compliance, mechanically favorable intercostal muscle length and contraction, and lower resistance to diaphragmatic contraction than in the supine position, resulting in increased ventilation compared to that in the supine and Fowler positions.<sup>21</sup> In contrast, in older people, age-related structural changes, such as round back and decreased mobility of the thorax, are thought to affect respiratory function.<sup>22</sup> In addition, respiratory function is adapted to a narrow range of physical activity states, and it stabilizes at a low level in bedridden cases.<sup>23</sup> Therefore, among the bedridden older participants in the present study, anterior tilt and a circular back posture with an increasing reclining angle made

antigravity extension of the trunk difficult, and limited thoracic movement in the direction of gravity<sup>24</sup> may have caused the decrease in ventilation volume in the 80° position. Further evaluation of the setting of the reclined sitting angle for bedridden older patients in clinical situations is needed.

### **Limitations**

This study has several limitations. First, since this was a single-center study and the number of participants was not large, we need to be cautious in generalizing the results of this study. Second, the degree of round back was not measured. It is necessary to clarify in a future study how much dorsal circularity affects the ventilation rate in the 80° position. Third, the bedridden older participants in this study were assumed to be in a stable condition after recovery from a condition that required hospitalization. Therefore, metabolic and ventilatory indices were evaluated after recovery. However, we could not exclude the influence of the causative disease at the time of admission. In addition, many of the participants had a poor nutritional condition. This may have decreased the muscle mass of the whole body, including respiratory muscles, and may have affected the results of metabolic and ventilatory indices. In order to further analyze postural and ventilatory changes in bedridden older patients, it is necessary to recruit participants who live at home or in facilities, have confirmed clinical stability, and have a good nutritional condition.

### **Implication**

The results of this study indicate that appropriate reclining sitting positions for bedridden older patients can promote increased ventilation in these patients, which may lead to the generation of effective coughing and may help protect these patients from pneumonia.

### **Conclusions**

Among bedridden older patients, sitting in a wheelchair is a safe and very low-intensity physical activity.  $V_T$  was maximal in the Fowler position, and the ventilatory volume did not increase with an increase in the reclining angle. These findings suggest that appropriate reclining postures in clinical situations can promote an increase in the ventilatory rate in bedridden older patients.

### **Acknowledgements**

We are indebted to patients who agreed to participate and cooperate in this study. The authors would like to thank the members of the Department of Rehabilitation, Kofu Kyoritsu Hospital, including Nana Muranaka, Ryota Sakai, Genki Komatsu, and Taiki Miura, for their cooperation in the acquisition of data. Finally, the authors would like to thank the Kobe University Graduate School of Health Sciences, including Kaoru Hanaie, Takumi Yamaguchi, Shigehumi Murakami, Kentaro Iwata, Yusuke Iwata, Ken Umehara, Keita Ohashi, Haruko Suzuki, Kohei Otake, Hiroki Mizusawa, Masaaki Kobayashi, for their constructive comments on this paper. We would like to thank Enago for English language editing.

### **Author Contributions**

YY prepared the manuscript and reviewed the literature. YMi (Yuji Mitani), AY, KY, YO (Yutaro Oki), YO (Yukari Oki), YK, and AI conceived and designed the study. YY and YMa (Yasumichi Maejima) collected the data. YY and AY analyzed the data. YY, AY, KM, KY, YO (Yutaro Oki), YO (Yukari Oki), and AI drafted and critically revised the manuscript for important intellectual content. All authors read and approved the final manuscript.

## References

1. Teramoto S, Fukuchi Y, Sasaki H, et al. High incidence of aspiration pneumonia in community- and hospital-acquired pneumonia in hospitalized patients: A multicenter, prospective study in Japan. *J Am Geriatr Soc* 2008;56(3):577–579.
2. Mitani Y, Oki Y, Fujimoto Y, et al. Relationship between functional independence measure and geriatric nutritional risk index in pneumonia patients in long-term nursing care facilities. *Geriatr Gerontol Int* 2017;17(10):1617–1622.
3. The Japanese Respiratory Society. The JRS guidelines for the management of pneumonia in adults. 2017.
4. Momosaki R, Yasunaga H, Matsui H, Horiguchi H, Fushimi K, Abo M. Effect of early rehabilitation by physical therapists on in-hospital mortality after aspiration pneumonia in the elderly. *Arch Phys Med Rehabil* 2015;96(2):205–209.
5. Mitani Y, Oki Y, Fujimoto Y, et al. Relationship between the Functional Independence Measure and Mann Assessment of Swallowing Ability in hospitalized patients with pneumonia. *Geriatr Gerontol Int* 2018;18(12):1620–1624.
6. Langmore SE, Skarupski KA, Park PS, Fries BE. Predictors of aspiration pneumonia in nursing home residents. *Dysphagia* 2002;17(4):298–307.
7. Miura T, Ishikawa Y. Respiratory physiotherapy for infants under respiratory management. (in Japanese). *ICU & CCU* 2017;41(3):187–192.
8. Dean E, Ross J. Discordance between cardiopulmonary physiology and physical therapy. Toward a rational basis for practice. *Chest* 1992;101(6):1694–1698.
9. Yoshizaki H, Yoshida A, Hayashi F, Fukuda Y. Effect of posture change on control of ventilation. *Jpn J Physiol* 1998;48(4):267–273.
10. Chang AT, Boots RJ, Brown MG, Paratz JD, Hodges PW. Ventilatory changes following head-up tilt and standing in healthy subjects. *Eur J Appl Physiol* 2005;95(5–6):409–417.
11. Morino A, Shida M, Tanaka M, et al. Comparison of changes in tidal volume associated with expiratory rib cage compression and expiratory abdominal compression in patients on prolonged mechanical ventilation. *J Phys Ther Sci* 2015;27(7):2253–2256.
12. Iki H, Sawa S, Teranishi T, Tomita M, Nishii K, Yamada K. Effect of exercise therapy on cytokine secretion in the saliva of bedridden patients. *J Phys Ther Sci* 2016;28(10):2871–2876.
13. Nakazawa A, Satake M, Kimoto Y, et al. Cardiorespiratory responses and metabolism during basic position in patients with acute stroke. *J Jpn Phys Ther Assoc* 2017;44(6):415–425.

14. Wasserman K. Diagnosing cardiovascular and lung pathophysiology from exercise gas exchange. *Chest* 1997;112(4):1091–1101.
15. Dellamonica J, Lerolle N, Sargentini C, et al. Effect of different seated positions on lung volume and oxygenation in acute respiratory distress syndrome. *Intensive Care Med* 2013;39(6):1121–1127.
16. Milic-Emili J, Grunstein MM. Drive and timing components of ventilation. *Chest* 1976;70(1); Supplement:131–133.
17. Boggs DF, Tenney SM. Scaling respiratory pattern and respiratory ‘drive’ *Respir Physiol* 1984;58(3):245–251.
18. Laveneziana P, Galarducci A, Binazzi B, Stendardi L, Duranti R, Scano G. Inhaled furosemide does not alleviate respiratory effort during flow-limited exercise in healthy subjects. *Pulm Pharmacol Ther* 2008;21(1):196–200.
19. Walls CE, Laine CM, Kidder IJ, Bailey EF. Human hypoglossal motor unit activities in exercise. *J Physiol* 2013;591(14):3579–3590.
20. Romei M, Mauro AL, D’Angelo MG, et al. Effects of gender and posture on thoraco-abdominal kinematics during quiet breathing in healthy adults. *Respir Physiol Neurobiol* 2010;172(3):184–191.
21. Sonpeayung R, Tantisuwat A, Klinphon T, Thaveeratitham P. Which body position is the best for chest wall motion in healthy adults? A meta-analysis. *Respir Care* 2018;63(11):1439–1451.
22. Bunki H, Shibata N, Yorimoto A. Functional characteristics and measurement significance of respiratory muscle strength in the elderly. *JAMHTS* 2013;40(4):451–456.
23. Truong AD, Fan E, Brower RG, Needham DM. Bench-to-bedside review: Mobilizing patients in the intensive care unit – From pathophysiology to clinical trials. *Crit Care* 2009;13(4):216–216.
24. Takashima S, Nozoe M, Mase K, Kouyama Y, Matsushita K, Ando H. Effects of posture on chest-wall configuration and motion during tidal breathing in normal men. *J Phys Ther Sci* 2017;29(1):29–34.

Table 1. Participants' characteristics<sup>1</sup>

Characteristics	All (n = 19)
Age, y	83.2 ± 8.4
Male (n, %)	12 (63.2)
BMI, kg/m <sup>2</sup>	17.3 ± 3.3
Cause of hospitalization	
-Pyelonephritis (n, %)	3 (15.8)
-Aspiration pneumonia (n, %)	10 (52.6)
-Bacterial pneumonia (n, %)	5 (26.3)
-Post-drug-induced bradycardia (n, %)	1 (5.3)
Alb, g/dl	2.8 ± 0.9
BUN, mg/dl	19.2 ± 8.4
Cr, mg/dl	0.7 ± 0.5
CRP, mg/dl	2.1 ± 2.4
eGFR, mlmin <sup>-1</sup> 1.73m <sup>-2</sup>	98 ± 49.9

Data presented as mean ± SD, or n (%).

<sup>1</sup>BMI, body mass index; alb, albumin; BUN, blood urea nitrogen; Cr, creatinine; CRP, C-reactive protein; eGFR, estimated glomerular filtration rate

Table 2. Changes in measurement parameters due to posture changes

	Supine	Fowler	80°
HR, bpm	76.0 ± 14.4	75.1 ± 15.0	76.5 ± 15.6
SpO <sub>2</sub> , %	95.5 ± 2.6	95.2 ± 2.1	94.7 ± 2.5
$\dot{V}O_2/W$ , ml/min/kg	2.9 ± 0.8	3.2 ± 0.5	3.2 ± 0.7
$\dot{V}O_2$ , ml/min	114.6 ± 31.6	125.4 ± 27.1	124.0 ± 22.9
$\dot{V}CO_2$ , ml/min	107.6 ± 31.6	116.6 ± 27.9	114.3 ± 24.9
R	0.94 ± 0.1	0.93 ± 0.1	0.92 ± 0.1 †
V <sub>T</sub> , ml	398.4 ± 112.2	426.9 ± 106.8 *	416.8 ± 92.5
VE, L/min	6.5 ± 1.4	7.1 ± 0.6	7.2 ± 1.5
RR, bpm	16.8 ± 3.0	17.0 ± 3.3	17.7 ± 3.8
Ti, s	1.4 ± 0.3	1.4 ± 0.3	1.4 ± 0.4
Te, s	2.4 ± 0.5	2.3 ± 0.5	2.3 ± 0.7
Ttot, s	3.8 ± 0.7	3.7 ± 0.7	3.7 ± 0.3
VT/Ti, ml/s	281.1 ± 54.1	307.6 ± 59.2 *	314.8 ± 75.7

The results for continuous variables are presented as mean ± standard deviation.

\* Bonferroni's correction was used to compare between the supine and Fowler position

† Bonferroni's correction was used to compare between the supine and 80° position

HR, heart rate; SpO<sub>2</sub>, transcutaneous oxygen saturation;  $\dot{V}O_2/W$ , oxygen uptake divided by weight;  $\dot{V}O_2$ , carbon dioxide output;  $\dot{V}CO_2$ , oxygen uptake; R, gas exchange ratio; V<sub>T</sub>, tidal volume; VE, minute volume; RR, respiratory rate; Ti, inspiratory time; Te, expiratory time; Ttot, total respiratory time; VT/Ti, mean inspiratory flow






Fig. 1. Three postures adopted by the subjects. From top to bottom: supine at rest (supine), 40° reclined and 20° tilt-up (Fowler), and 80° reclined wheelchair (80°). Measurements were taken for 6 minutes in each posture.




Fig. 2. Scatter plots of the change in oxygen uptake ( $\dot{V}O_2$ ) and tidal volume ( $V_T$ ) caused by the change from the supine position. A) from supine to Fowler and B) from supine to 80°. The black lines represent regression lines. Pearson's product-moment correlation coefficient was used to examine the relationship between metabolic and ventilatory parameters ( $\dot{V}O_2$  and  $V_T$ ).