

PDF issue: 2025-07-06

# Tooth Loss as a Predictor of Long-Term Care Requirements in the Elderly: A Study in Kobe City, Japan

Kakei, Yasumasa ; Kagimura, Tatsuo ; Yamamoto, Yasuji ; Osaki, Tohmi ; Kajita, Hiroyuki ; Kojima, Shinsuke ; Kowa, Hisatomo ; Kawabata, Miyuk…

(Citation) Cureus,15(12):e49851

(Issue Date) 2023-12-02

(Resource Type) journal article

# (Version)

Version of Record

# (Rights)

© Copyright 2023 Kakei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

# (URL)

https://hdl.handle.net/20.500.14094/0100485967



Review began 09/18/2023 Review ended 11/22/2023 Published 12/02/2023

#### © Copyright 2023

Kakei et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

# Tooth Loss as a Predictor of Long-Term Care Requirements in the Elderly: A Study in Kobe City, Japan

Yasumasa Kakei <sup>1</sup>, Tatsuo Kagimura <sup>2</sup>, Yasuji Yamamoto <sup>3</sup>, Tohmi Osaki <sup>4</sup>, Hiroyuki Kajita <sup>4</sup>, Shinsuke Kojima <sup>2</sup>, Hisatomo Kowa <sup>5</sup>, Miyuki Kawabata <sup>6</sup>, Takumi Hasegawa <sup>1</sup>, Masaya Akashi <sup>1</sup>, Yoji Nagai <sup>6</sup>

1. Department of Oral and Maxillofacial Surgery, Kobe University Hospital, Kobe, JPN 2. Translational Research Centre for Medical Innovation, Foundation for Biomedical Research and Innovation, Kobe, JPN 3. Department of Biosignal Pathophysiology, Kobe University Graduate School of Medicine, Kobe, JPN 4. Faculty of Rehabilitation, Kobe Gakuin University, Kobe, JPN 5. Department of Rehabilitation Science, Kobe University Graduate School of Health Sciences, Kobe, JPN 6. Department of Clinical Research Facilitation, Kyoto University Hospital, Kyoto, JPN

Corresponding author: Yasumasa Kakei, ykakei@med.kobe-u.ac.jp

### Abstract

Introduction: The Kobe project, which utilizes prospective data from the national health insurance system, focuses on early detection and preventive strategies through the Frail Kenshin health check-up program. Previous research has underscored the correlation between tooth loss and the decline in physical and cognitive functions. In this study, using Kobe project data, we examined the link between remaining teeth and long-term care needs in individuals aged 64-65 years, with primary and secondary objectives involving various health parameters and quality of life.

Methods: We analyzed baseline data from a prospective study conducted alongside the Frail Check program for generally healthy individuals aged 64-65 years to examine the relationship between the number of remaining teeth and various health indicators. This study focused on citizens aged 64-65 years to identify those at risk of needing long-term care by the age of 65 years.

Results: Data from 1,530 participants were obtained, excluding eight individuals for specific reasons. At the end of the follow-up period, 41 (2.7%) individuals required support and 15 (1.0%) needed long-term care alone. The data revealed a significant association between the number of remaining teeth and the need for long-term care or support, as demonstrated by the Cochran-Armitage trend test (p<0.001).

Although trends were noted for nutrition and total Cognitive Functional Instrument Self scores, they did not reach statistical significance. Additionally, a decrease in the number of remaining teeth was significantly associated with worse European Quality of Life Five Dimensions (EQ-5D-5L) visual analog scale scores, mobility, and regular activities (p<0.001).

Conclusion: Tooth loss indicates the potential long-term care needs of older adults. Monitoring oral health is crucial for addressing care requirements.

Categories: Geriatrics

Keywords: body mass index, frailty, long-term care need, quality of life, tooth loss

#### Introduction

Physical and cognitive decline necessitates the provision of long-term care and support services for community-dwelling older adults. As Japan currently has the largest population of super-aged individuals worldwide [1], care for this population places a serious burden on the healthcare system and has wide-ranging socioeconomic implications [2,3].

Various population-based studies have evaluated methods for the prevention and early detection of physical and cognitive decline in community-dwelling older adults in Japan. One such initiative is the Kobe project, which includes the first large prospective study to evaluate the predictive ability of neuropsychological assessment tools for long-term care certification in older adults [4]. One of the studies in this project used the data collected by local government municipal offices from older adults enrolled in the Kokumin Kenko Hoken National Health Insurance system in Japan; the insurance covers all Japanese citizens and long-term residents. In this context, Frail Kenshin is a health check-up program conducted by Kobe City that includes physically accessible components in addition to self-assessed questions related to frailty. It evaluates motor and oral functions, nutrition, forgetfulness, mental and dental health, chewing ability, swallowing ability, grip strength, height, weight, body mass index (BMI), and other health-related parameters.

Numerous epidemiological studies have reported an association between tooth loss and physical and

#### How to cite this article

Kakei Y, Kagimura T, Yamamoto Y, et al. (December 02, 2023) Tooth Loss as a Predictor of Long-Term Care Requirements in the Elderly: A Study in Kobe City, Japan. Cureus 15(12): e49851. DOI 10.7759/cureus.49851

cognitive decline in older adults [5-8]. Reports suggest that in addition to a reduction in the number of food items that can be consumed owing to a decrease in the number of chewing points, a decrease in bite strength leads to a decline in motor skills, including grip strength and walking function, which are also associated with a decline in cognitive function [9-11]. A study using nationwide data from Korea also showed an association between the number of remaining teeth and health-related quality of life (QOL) [12].

In view of these findings, we analyzed data obtained from a prospective study under the Kobe project. The study evaluated the relationship between data obtained using the Cognitive Functional Instrument (CFI), European Quality of Life Five Dimensions (EQ-5D-5L), and other frailty-related parameters and explored the risk of future long-term care and support needs within three years in older adults aged 64-65 years in the assessment year.

The primary objective of this study was to assess the association between the number of teeth and the need for long-term care. The secondary objectives were to assess the association between the number of teeth and chewing points, grip strength, motor functions such as walking, and QOL.

This article was previously posted to the Research Square preprint server on 27 July 2022.

## **Materials And Methods**

#### **Ethical issues**

All participants were treated in accordance with the principles of the Declaration of Helsinki developed by the World Medical Association in 1964 and modified in October 2013 at the World Medical Association Fortaleza General Assembly (Brazil) [13]. All research conformed to the "Law on Protection of Personal Information" (May 30, 2003; Law No. 57) [14]. This study was approved by the Ethics Review Committee of the WHO (ERC.0002899) and Kobe University (Approval No. 170018). The need for informed consent was waived as this was a post hoc analysis of the data obtained from a prospective study.

#### Study design and setting

This study analyzed baseline data from a prospective study conducted in conjunction with the Frail Check program for Kobe citizens aged 64-65 years, initiated by the municipal office of Kobe City. The Frail Check program, which aims to assess general health and frailty-related parameters, incorporates dental assessments in its evaluation process.

Participants aged 64-65 years were selected because they represented the minimum age of the elderly in Japan. Focusing on this age group, we aimed to identify individuals at risk of requiring long-term care at 65 years of age. Early intervention is crucial for preventing or minimizing the need for long-term care, particularly in individuals with fewer remaining teeth who are at higher risk.

Baseline data were obtained between August 2017 and March 2018. The program was conducted repeatedly at multiple venues within the community. Numerous subjects attended each occasion where various general health and frailty-related parameters, including dental assessments, were individually evaluated. Following the assessments, the subjects received their results along with medical and non-medical recommendations.

Long-term care insurance data up to March 31, 2020, were obtained from the administrative records of Kobe City, allowing for a comprehensive analysis of the association between the number of teeth and the need for long-term care.

By incorporating dental assessments within the Frail Check program, specific relationships between the number of teeth and the need for long-term care were identified, and secondary objectives were explored, such as the association between the number of teeth and variables including the number of chewing points, grip strength, motor functions (e.g., walking), and QOL.

#### **Subjects**

The subjects included citizens of Kobe, Japan, aged between 64 and 65 years who attended the Frail Check program in Kobe City. Hyogo Prefecture, where Kobe City is located, is representative of the national average with respect to the number of elderly people needing long-term care. Citizens who were already certified for long-term care at the time of the Frail Check data collection were excluded.

#### Variables evaluated

Data regarding the Frail Check test, CFI Self questionnaire, and EQ-5D-5L assessments were collected during the Frail Check program conducted in Kobe. The Frail Check sheet devised by Kobe City examines the following data: age, sex, daily living activity/forgetfulness, exercise function, nutritional status, oral and dental and cardiac health, calf circumference, results of the finger-ring (Yubi-wakka) test (using fingers of both hands to make a ring around one's calf), chewing and grip strength, swallowing, and stand-up tests.

As described by Nagai et al. [4], details pertaining to the number of remaining teeth were obtained from the Frail Check data for this analysis; the mastication score and number of foods that could be chewed were also assessed; the mastication score and the number of food items that could be chewed were also assessed. The remaining teeth were categorized into five groups: 0, 1-4, 5-9, 10-19, and >20. These categories were retained for this analysis.

#### Tools used in the prospective study

CFI Self Questionnaire Japanese Version

The Japanese version of the CFI Self questionnaire was used to evaluate cognitive function-related ADL (activities of daily living) in the prospective study. The questionnaire consists of 14 items, where "yes" = 1 point, "probable" = 0.5 points, and "no" = 0 points; the total of the points for all items constitutes the individual scores.

EQ-5D-5L for Evaluating Health-Related QOL

EQ-5D-5L was used for measuring health-related QOL; with this tool, individual QOL is defined as a single value between 0 and 1 with upper and decreased limits of "complete health = 1" and "death = 0," respectively, based on answers to five questions (five levels).

#### Information regarding long-term care certification

Information on subjects certified as needing long-term care within three years was obtained from the Long-Term Care Certification database of Kobe City. In accordance with the Long-Term Care Insurance Law standard, the Kobe City Long-Term Care Need Certification criteria evaluated the following: level of longterm care needed, scored between 1 and 5 based on the assessment of care requirements, and level of assistance needed, scored as 1 or 2.

#### Sample size

The Frail Check data for 1530 subjects from Kobe City were obtained for this study. Among the eight individuals excluded, five had moved out of Kobe City; therefore, the results of the Kobe City-approved certification for the need for long-term care or support were unknown. Two cases were excluded because of duplication of Frail Check results, and one patient was already certified as requiring long-term care at the time of Frail Check screening.

#### Statistical analysis

We performed a trend test for the number of teeth for each evaluation indicator. Dichotomous variables, such as the presence or absence of care needs, were evaluated using the Cochran-Armitage trend test, which determines the proportion of individuals requiring care depending on the number of remaining teeth. Continuous variables and ordinal data were evaluated using the Jonckheere-Terpstra test. For this analysis, p-values <0.001 were considered statistically significant. SAS version 9.4 (SAS Institute Inc., Cary, NC) and R (Version 3.6; R Core Team 2020) were used for the analyses.

#### **Results**

The data of 1508 older adults were analyzed in this study; there were 957 (63.5%) and 546 (36.2%) female and male participants, respectively. At the end of the follow-up period, 41 (2.7%) and 15 (1.0%) individuals were certified as needing support or long-term care and long-term care alone, respectively.

The Cochran-Armitage trend test demonstrated a significant increase in the need for long-term care or support, with a progressive decline in the number of remaining teeth (p<0.001). Certification for long-term care needs also followed a similar trend (Table 1).

			Number of Teeth (n)											
	Total	Fotal		0		1-4		5-9		10-19			Cochran-Armitage trend test	
	n	%	n	%	n	%	n	%	n	%	n	%		
Total	1,508	100	17	100	18	100	39	100	176	100	1,258	100		
Gender														
Female	957	63.5	6	35.3	7	38.9	21	53.8	91	51.7	832	66.1		
Male	546	36.2	11	64.7	11	61.1	18	46.2	85	48.3	421	33.5		
Information not provided	5	0.3	0	0	0	0	0	0	0	0	5	0.4		
Long-term care or support	need													
Yes	41	2.7	4	23.5	0	0	5	12.8	9	5.1	23	1.8	p<0.001	
No	1,467	97.3	13	76.5	18	100	34	87.2	167	94.9	1,235	98.2		
Long-term care need														
Yes	15	1	1	5.9	0	0	2	5.1	5	2.8	7	0.6	p=0.004	
No	1,493	99	16	94.1	18	100	37	94.9	171	97.2	1,251	99.4		

TABLE 1: Association between long-term care need or long-term care and support need and the number of remaining teeth, as indicated by the Frail Check data

Evaluation of the Frail Check data demonstrated that a decrease in the number of remaining teeth was associated with a decline in general frailty, including motor and oral functions, prevention of seclusion, memory, mental health, mastication scores, total number of food items that could be chewed, and BMI (p<0.001) (Figure 1). In addition, a reduction in the number of remaining teeth was associated with decreased calf and standing test scores during the check-up (p<0.001). Although similar trends were noted for the nutrition and total CFI Self scores, these associations were not statistically significant (Tables 2, 3). Analysis of the EQ-5D-5L utility index data demonstrated that a reduction in the number of remaining teeth was associated with a significant worsening of EQ-5D-5L visual analog scale scores, degree of mobility, and regular activities (p<0.001) (Table 4).





			Number of teeth (n)			Jonckheere- Terpstra test		
		Total	0	4- Jan	9- May	19- Oct	>20	
	Ν	1,508	17	18	39	176	1,258	
FC questionnaire Comprehensive score	Mean	2.66	5.82	3.56	4	3.4	2.45	p<0.001
	Total         0         Jan         May         Oct         >20           N         1,508         17         18         39         176         1,258           Mean         2.66         5.82         3.56         4         3.4         2.45           SD         2.28         4.59         2.33         2.62         2.4         2.13           Mean         1,508         17         18         39         176         1,258           Mean         2.62         4.59         2.33         2.62         2.4         2.13           Mean         1,508         17         18         39         176         1,258           Mean         0.74         1.71         0.83         1.15         1         0.68           SD         1.01         1.72         1.04         1.01         1.12         0.96           Mean         0.35         0.35         0.33         0.38         0.32         0.35           SD         0.52         0.49         0.49         0.49         0.49         0.48         0.53           Mean         0.62         0.76         1.06         1.03         0.9         0.56 <td></td>							
	Ν	1,508	17	18	39	176	1,258	
FC questionnaire Motor function score	Mean	0.74	1.71	0.83	1.15	1	0.68	p<0.001
	SD	1.01	1.72	1.04	1.01	1.12	0.96	
	Ν	1,508	17	18	39	176	1,258	
FC questionnaire Nutrition	Mean	0.35	0.35	0.33	0.38	0.32	0.35	p=0.863
	SD	0.52	0.49	0.49	0.49	0.48	0.53	
	Ν	1,508	17	18	39	176	1,258	p<0.001
	Mean	0.62	0.76	1.06	1.03	0.9	0.56	
FC questionnaire Oral function score		Numb	Number of teeth (n)					Jonckheere- Terpstra test
		Total	0	4- Jan	9- May	19- Oct	>20	
	Ν	1,508	17	18	39	176	1,258	
FC questionnaire Prevention of withdrawal score	Mean	0.15	0.41	0.33	0.26	0.21	0.13	p<0.001
	SD	0.38	0.62	0.49	0.44	0.42	0.36	

	Ν	1,508	17	18	39	176	1,258	
FC questionnaire Forgetfulness score	Mean	0.36	1.18	0.5	0.54	0.45	0.33	p<0.001
	SD	0.61	1.13	0.71	0.68	0.66	0.58	
	Ν	1,508	17	18	39	176	1,258	
FC questionnaire Mental health score	Mean	0.64	2.24	1.22	1.05	1.03	0.55	p<0.001
	SD	1.2	2.05	1.63	1.54	1.48	1.09	
	Image<	39	176	1,258				
FC questionnaire Chewing test score	Mean	48.22	38.18	41.5	43.54	46.34	48.86	p<0.001
	SD	4.44	14.85	8.28	6.27	5.04	3.41	
	Ν	1,506	17	18	39	176	1,256	
	Mean	24.14	19.09	20.75	21.77	23.17	24.47	p<0.001
Total number of foods that can be chewed (1 point: chewable 0.5 points: hard to chew)	Mean0.361.180.4SD0.611.130Man1.5081.71.2Man1.22.051Man1.5001.71.2Man4.4238.184Man4.4238.184Man2.441.4001.2Man2.441.4001.2Man2.4141.0001.2Man2.4141.0001.2Man2.4141.0001.2Man2.041.31.2Man2.041.31.2Man1.4011.41.2Man1.83.11.2Man1.83.11.2Man1.83.11.2Man1.83.11.2Man1.83.11.2Man1.41.31.2Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4Man1.41.41.4M	eth (n)	Jonckheere- Terpstra test					
		Total	0	4- Jan	9- May	19- Oct	>20	
	SD	2.04	7.43	4.14	3.13	2.52	1.4	
	Ν	1,451	16	16	39	171	1,209	p=0.007
Total CFI Self score	Mean	1.86	3.5	2.47	2.03	2.25	1.77	
	SD	1.88	3.49	2.57	1.94	2.22	1.77	
	Ν	1,507	17	18	39	176	1,257	p<0.001
FC questionnaire Calf circumference	Mean	35	36.1	35.3	35.9	35.8	34.9	
	SD	3.2	3.1	3.3	3.3	3.4	3.2	
	Ν	1,467	15	17	38	167	1,230	
FC questionnaire Standing up examination score	Mean	9.27	8.2	8.24	8.68	8.79	9.38	p<0.001
	SD	2.59	2.46	2.44	2.69	2.26	2.62	
	Ν	1,445	15	17	39	172	1,202	
FC questionnaire BMI	Mean	22.58	24.46	23.19	24.14	23.7	22.34	p<0.001
	SD	3.4	4.5	2.54	4.09	3.69	3.28	
	Ν	1,408	11	11	29	150	1,207	
	N	1,408 Numb	11 er of tee	11 eth (n)	29	150	1,207	Jonckheere- Terpstra test
FC questionnaire Mastication	N	1,408 Numb Total	11 er of tee 0	11 eth (n) 4- Jan	29 9- May	150 19- Oct	1,207 >20	Jonckheere- Terpstra test
FC questionnaire Mastication	N	1,408 Numb Total 2.35	11 er of tee 0 2.91	11 eth (n) 4- Jan 2.82	29 9- May 2.62	150 <b>19-</b> Oct 2.5	1,207 > <b>20</b> 2.31	Jonckheere- Terpstra test
FC questionnaire Mastication	N Mean SD	1,408 Numb Total 2.35 0.99	11 er of tee 0 2.91 1.14	11 eth (n) 4- Jan 2.82 1.08	29 <b>9-</b> <b>May</b> 2.62 1.29	150 <b>19-</b> <b>Oct</b> 2.5 1.06	1,207 >20 2.31 0.97	Jonckheere- Terpstra test
FC questionnaire Mastication	N Mean SD N	1,408 Numb Total 2.35 0.99 1,485	11 er of tee 0 2.91 1.14 17	11 eth (n) 4- Jan 2.82 1.08 17	29 <b>9-</b> <b>May</b> 2.62 1.29 39	150 <b>19-</b> <b>Oct</b> 2.5 1.06 172	1,207 > <b>20</b> 2.31 0.97 1,240	Jonckheere- Terpstra test
FC questionnaire Mastication	N Mean SD N Mean	1,408 Numb Total 2.35 0.99 1,485 27.8	11 er of tee 0 2.91 1.14 17 27.9	11 eth (n) 4- Jan 2.82 1.08 17 32.5	29 <b>9-</b> <b>May</b> 2.62 1.29 39 28.5	150 <b>19-</b> <b>Oct</b> 2.5 1.06 172 28.9	1,207 <b>&gt;20</b> 2.31 0.97 1,240 27.5	Jonckheere- Terpstra test p=0.004 p=0.019
FC questionnaire Mastication	N Mean SD N Mean SD	1,408 Numb Total 2.35 0.99 1,485 27.8 8.6	11 er of tee 0 2.91 1.14 17 27.9 9.1	11 eth (n) Jan 2.82 1.08 17 32.5 9.7	29 <b>9- May</b> 2.62 1.29 39 28.5 8.9	150 <b>19-</b> <b>0ct</b> 2.5 1.06 1.72 28.9 9.3	1,207 <b>&gt;20</b> 2.31 0.97 1,240 27.5 8.4	p=0.019
FC questionnaire Mastication	N Mean SD N Mean SD N SD	1,408 Numb 2.35 0.99 1,485 27.8 8.6 1,470	11 er of tee 0 2.91 1.14 17 27.9 9.1 17	11 eth (n) 4- Jan 2.82 1.08 17 32.5 9.7 17	29 <b>9-</b> <b>May</b> 2.62 1.29 39 28.5 8.9 39	150 <b>19-</b> <b>Oct</b> 2.5 1.06 172 28.9 9.3 171	1,207 <b>&gt;20</b> 2.31 0.97 1,240 27.5 8.4 1,226	Jonckheere- Terpstra test p=0.004 p=0.019

# TABLE 2: Data pertaining to items in the Frail Check questionnaire that demonstrated association with the number of remaining teeth

FC, Frail Check; CFI, cognitive functional instrument; BMI, body mass index; SD, standard deviation.

			Number of teeth (n)										lonakhaara
	Total		0		104	4	509	9	10□19		>20		-Terpstra
	n	%	n	%	n	%	n	%	n	%	n	%	test
Total	1,508	100	17	100	18	100	39	100	176	100	1,258	100	
FC questionnaire/Grip strength assessment													p=0.029
Weak	22	1.5	2	11.8	0	0.0	2	5.1	4	2.3	14	1.1	
Somewhat weak	176	11.7	3	17.6	1	5.6	6	15.4	24	13.6	142	11.3	
Normal	230	15.3	2	11.8	4	22.2	7	17.9	27	15.3	190	15.1	
Somewhat strong	789	52.3	9	52.9	9	50.0	18	46.2	90	51.1	663	52.7	
Strong	274	18.2	1	5.9	3	16.7	6	15.4	28	15.9	236	18.8	
Not performed	17	1.1	0	0.0	1	5.6	0	0.0	3	1.7	13	1.0	
FC questionnaire/Standing up assessment													p<0.001
Slow	12	0.8	1	5.9	0	0.0	0	0.0	1	0.6	10	0.8	
Somewhat slow	349	23.1	4	23.5	7	38.9	16	41.0	45	25.6	277	22.0	
Normal	272	18.0	2	11.8	3	16.7	6	15.4	34	19.3	227	18.0	
Somewhat fast	591	39.2	7	41.2	5	27.8	12	30.8	71	40.3	496	39.4	
Fast	243	16.1	1	5.9	2	11.1	4	10.3	16	9.1	220	17.5	
Not performed	41	2.7	2	11.8	1	5.6	1	2.6	9	5.1	28	2.2	
FC questionnaire Finger-ring frailty test													p=0.014
A gap is present	187	12.4	1	5.9	1	5.6	4	10.3	17	9.7	164	13.0	
An exact circle	657	43.6	6	35.3	7	38.9	13	33.3	76	43.2	555	44.1	
Cannot make a complete circle due to the thickness of muscles	660	43.8	10	58.8	10	55.6	22	56.4	83	47.2	535	42.5	
Not performed	4	0.3	0	0.0	0	0.0	0	0.0	0	0.0	4	0.3	
FC questionnaire Mastication													p=0.004
Green	17	1.1	1	5.9	0	0.0	2	5.1	4	2.3	10	0.8	
Yellow-green	157	10.4	2	11.8	4	22.2	6	15.4	24	13.6	121	9.6	
Pink	448	29.7	4	23.5	2	11.1	8	20.5	44	25.0	390	31.0	
Dark cerise	461	30.6	3	17.6	4	22.2	5	12.8	49	27.8	400	31.8	
Red	325	21.6	1	5.9	1	5.6	8	20.5	29	16.5	286	22.7	
Not performed	100	6.6	6	35.3	7	38.9	10	25.6	26	14.8	51	4.1	
FC questionnaire Swallowing													p=0.462
Less than 2 times	27	1.8	0	0.0	1	5.6	0	0.0	2	1.1	24	1.9	
More than 3 times	1,463	97.0	17	100.0	17	94.4	39	100.0	169	96.0	1,221	97.1	
Not performed	18	1.2	0	0.0	0	0.0	0	0.0	5	2.8	13	1.0	

# TABLE 3: Data pertaining to assessed items in the Frail Check questionnaire that demonstrated an association with the number of remaining teeth

FC, Frail Check.

		Tatal	Numb	er of te	eth (n)	lanakhaana Tamatra taat		
		Iotai	0	1-4	5-9	10-19	>20	Jonckneere-Terpstra test
EQ-5D utility index	Ν	1,414	14	15	34	162	1,189	
	Mean	0.92	0.81	0.91	0.87	0.90	0.92	p=0.002
	SD	0.10	0.25	0.12	0.17	0.12	0.10	
EQ-5D VAS	Ν	1,416	16	15	35	164	1,186	
	Mean	80.4	72.9	77.0	75.9	77.4	81.1	p<0.001
	SD	13.5	21.5	13.5	19.4	13.7	13.0	
EQ5D 1 - Degree of mobility	Ν	1,430	15	15	36	164	1,200	
	Mean	1.16	1.53	1.20	1.36	1.28	1.13	p<0.001
	SD	0.50	1.06	0.56	0.80	0.67	0.44	
EQ5D 2 - Personal care	Ν	1,432	16	15	35	164	1,202	
	Mean	1.03	1.50	1.00	1.11	1.03	1.02	p=0.044
	SD	0.24	1.10	0.00	0.53	0.20	0.19	
	Ν	1,434	16	16	36	164	1,202	
EQ5D 3 - Regular activities	Mean	1.08	1.31	1.44	1.25	1.12	1.06	p<0.001
	SD	0.35	0.70	1.09	0.65	0.43	0.29	
	Ν	1,430	15	15	36	165	1,199	
EQ5D 4 - Pain/discomfort	Mean	1.54	1.93	1.53	1.75	1.69	1.51	p=0.004
	SD	0.71	1.03	0.74	0.97	0.85	0.66	
	Ν	1,424	16	16	35	162	1,195	
EQ5D 5 - Anxiety/melancholy	Mean	1.18	1.63	1.31	1.29	1.21	1.17	p=0.010
	SD	0.46	1.02	0.48	0.52	0.50	0.44	

# TABLE 4: Data pertaining to items in the EQ-5D questionnaire that demonstrated association with the number of remaining teeth

EQ-5D, European Quality of Life Five Dimension; VAS, visual analog scale; SD, standard deviation.

## **Discussion**

The present study used baseline data from a prospective study performed under the Kobe project to explore new strategies for reducing the social burden of dementia in Japan [4]. The results from this cohort indicated a significant increase in long-term care or support needs, with a reduction in the number of remaining teeth among healthy older adults in Kobe City; a similar trend was also observed for long-term care needs.

The findings also suggested that a reduction in the number of remaining teeth was associated with general frailty, including motor and oral functions, prevention of seclusion, memory, mental health, mastication scores, and the total number of food items that could be chewed. BMI was also significantly associated with the number of remaining teeth, indicating the impact of poor nutrition on declining function.

Previous studies demonstrated an association between tooth loss and functional disabilities in older adults. A large prospective study from Australia, which used the EQ-5D-5L to evaluate the extent of disability, demonstrated an independent effect of self-reported oral health on responses to the questionnaire, indicating the impact of oral health problems on general health. The authors noted that oral symptoms were related to compliance with dietary guidelines, which in turn were associated with health status [9]. In our cohort, an association was observed between the number of remaining teeth and oral function, mastication scores, and the number of food items that could be chewed. These findings reinforce the need for nutritional screening and intervention in this population.

Numerous studies have reported an association between tooth loss and functional disability in older adults. A large prospective cohort study performed among community-dwelling adults in Japan, who were aged at least 70 years, showed a significantly increased risk of functional disability in those with fewer remaining teeth [10]. The authors concluded that regular dental care may reduce the risk of functional disability in older adults with missing teeth. Another prospective study performed under the same project investigated the association between maximum occlusal force, an objective predictor of chewing ability, and incident functional disability in Japanese adults aged at least 70 years. The findings demonstrated that a decreased maximum occlusal force conferred an increased risk of functional disability even after adjusting for possible confounding factors; this effect was observed irrespective of the number of remaining teeth. Nevertheless, these findings indicate the importance of maintaining oral function; they also demonstrate the association between the state of oral and general health in the elderly [11].

A recent prospective cohort study in Japan that evaluated the association between the number of teeth and functional disability in community-dwelling older adults found that having fewer than 20 teeth was an independent risk factor for functional disability; this association persisted after propensity score matching [5]. The authors reported that dementia, stroke, frailty, and other systemic conditions were the major causes of long-term care certification, which was the primary endpoint of this study. In this context, changes in nutrition and malnutrition have been reported as risk factors for the development of dementia, stroke, and frailty [15-17]. The number of teeth, therefore, plays a vital role in preventing conditions that require long-term care.

In the context of QOL, a study using data from the Korean National Health and Nutrition Examination Survey conducted over two years between 2010 and 2012 employed the EQ-5D-5L tool to evaluate the relationship between the number of remaining teeth and health-related QOL. The findings from this large dataset indicated that subjects who had more teeth had increased QOL scores; high QOL was particularly associated with certain components of the EQ-5D-5L, including mobility, self-care, and daily living [12]. Our findings also indicate a definite association between the number of teeth and motor function.

Cognitive impairment is a major contributor to the nursing care needs of older adults. In view of the likelihood of tooth loss in this population, a prospective cohort study from Japan that extended over four years evaluated the association between tooth loss and cognitive impairment in older adults living in the community. The subjects in this study were of a similar age group as that of our cohort. However, the authors used Mini-Mental State Examination scores to determine subjects' cognitive function. These findings suggest that tooth loss was a predictor of cognitive decline in this cohort and that both tooth loss and cognitive impairment chronically accumulated over time [6]. In our cohort, significant associations were observed between the number of remaining teeth, memory loss, and mental health status.

The major limitation of the present study is the sample size. Although the number of subjects was adequate for a reliable estimation of association, the level of significance of the association with individual items on the Frail Check and EQ-5D-5L questionnaires and Frail Check tests might have been affected. In addition, our cohort comprised individuals aged 64-65 years, which is a relatively young age group for assessing long-term care needs.

Further large-scale prospective studies with a longitudinal design are needed to address these limitations, including adults from older age groups. Despite these limitations, the findings of the present analysis indicate that tooth loss has a significant impact on the long-term care and support needs of community-dwelling older adults.

## Conclusions

This study found a significant association between the number of remaining teeth and the need for longterm care or support in a group of community-dwelling older adults aged 64-65 years. These findings emphasize the importance of maintaining oral health and highlight the potential impact of tooth loss on overall health and functional abilities. Further research with larger cohorts and older age groups is warranted to validate these findings and better understand the implications of tooth loss in long-term care.

# **Additional Information**

#### Disclosures

**Human subjects:** Consent was obtained or waived by all participants in this study. Ethics Review Committee of the World Health Organization and that of Kobe University issued approval ERC.0002899 and 170018, respectively. All participants in this study were treated in accordance with the principles of the Declaration of Helsinki, developed by the World Medical Association in 1964 and modified in October 2013 at the World Medical Association Fortaleza General Assembly (Brazil). All research conformed to the "Law on the Protection of Personal Information (May 30, 2003; Law No. 57). **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info**: Statistical analysis supported by the World Health Organization Centre for Health Development (WHO Kobe Centre – WKC: K18001). **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

#### Acknowledgements

The authors would like to thank Editage and Cactus Communications for editorial support in the form of medical writing, assembling tables, creating high-resolution images based on detailed directions, collating author comments, copyediting, fact-checking, and referencing as well as journal submission support. The authors have authorized the submission of this manuscript through Editage.

### **References**

- 1. World Health Organization. World Health Statistics 2021: Monitoring health for the SDGs . (2021). Accessed: July 31, 2023: https://apps.who.int/iris/bitstream/handle/10665/342703/9789240027053-eng.pdf.
- 2. World Health Organization. Global health and aging. (2021). Accessed: July 31, 2023: https://www.who.int/news-room/fact-sheets/detail/ageing-and-health.
- Akiyama N, Shiroiwa T, Fukuda T, Murashima S, Hayashida K: Healthcare costs for the elderly in Japan: Analysis of medical care and long-term care claim records. PLoS One. 2018, 13:e0190392. 10.1371/journal.pone.0190392
- Nagai Y, Kojima S, Kowa H, et al.: Kobe project for the exploration of newer strategies to reduce the social burden of dementia: A study protocol of cohort and intervention studies. BMJ Open. 2021, 11:e050948. 10.1136/bmjopen-2021-050948
- Komiyama T, Ohi T, Tomata Y, Tanji F, Tsuji I, Watanabe M, Hattori Y: Dental status is associated with incident functional disability in community-dwelling older Japanese: A prospective cohort study using propensity score matching. J Epidemiol. 2020, 30:84-90. 10.2188/jea.JE20180203
- Saito S, Ohi T, Murakami T, et al.: Association between tooth loss and cognitive impairment in communitydwelling older Japanese adults: A 4-year prospective cohort study from the Ohasama study. BMC Oral Health. 2018, 18:142. 10.1186/s12903-018-0602-7
- Tsakos G, Watt RG, Rouxel PL, de Oliveira C, Demakakos P: Tooth loss associated with physical and cognitive decline in older adults. J Am Geriatr Soc. 2015, 63:91-9. 10.1111/jgs.13190
- Yamamoto T, Kondo K, Hirai H, Nakade M, Aida J, Hirata Y: Association between self-reported dental health status and onset of dementia: A 4-year prospective cohort study of older Japanese adults from the Aichi Gerontological Evaluation Study (AGES) Project. Psychosom Med. 2012, 74:241-8. 10.1097/PSY.0b013e318246dffb
- 9. Brennan DS, Singh KA: Dietary, self-reported oral health and socio-demographic predictors of general health status among older adults. J Nutr Health Aging. 2012, 16:437-41. 10.1007/s12603-012-0006-3
- Komiyama T, Ohi T, Miyoshi Y, et al.: Association between tooth loss, receipt of dental care, and functional disability in an elderly Japanese population: The Tsurugaya project. J Am Geriatr Soc. 2016, 64:2495-502. 10.1111/jgs.14390
- 11. Ohi T, Komiyama T, Miyoshi Y, et al.: Maximum occlusal force and incident functional disability in older adults: The Tsurugaya project. JDR Clin Trans Res. 2018, 3:195-202. 10.1177/2380084418761329
- Park HE, Song HY, Han K, Cho KH, Kim YH: Number of remaining teeth and health-related quality of life: The Korean National Health and Nutrition Examination Survey 2010-2012. Health Qual Life Outcomes. 2019, 17:5. 10.1186/s12955-019-1078-0
- World Medical Association: World Medical Association Declaration of Helsinki: Ethical principles for medical research involving human subjects. JAMA. 2013, 310:2191-4. 10.1001/jama.2013.281053
- 14. Act on the Protection of Personal Information (Partly Unenforced) . (2003). Accessed: August 26, 2023: https://www.japaneselawtranslation.go.jp/en/laws/view/4241.
- Bonnefoy M, Berrut G, Lesourd B, et al.: Frailty and nutrition: Searching for evidence. J Nutr Health Aging. 2015, 19:250-7. 10.1007/s12603-014-0568-3
- 16. Hankey GJ: Nutrition and the risk of stroke. Lancet Neurol. 2012, 11:66-81. 10.1016/S1474-4422(11)70265-4
- Ozawa M, Ninomiya T, Ohara T, et al.: Self-reported dietary intake of potassium, calcium, and magnesium and risk of dementia in the Japanese: The Hisayama Study. J Am Geriatr Soc. 2012, 60:1515-20. 10.1111/j.1532-5415.2012.04061.x