



# Segmental mandibulectomy for mandibular osteoradionecrosis in an older adult with underweight status

Kusumoto, Junya ; Hayase, Yuka ; Susukida, Yuriko ; Sato, Takumi ; Sakakibara, Akiko ; Akashi, Masaya

---

**(Citation)**

Journal of Oral and Maxillofacial Surgery, Medicine, and Pathology, 36(1):58-63

**(Issue Date)**

2024-01

**(Resource Type)**

journal article

**(Version)**

Accepted Manuscript

**(Rights)**

© 2023Asian AOMS, ASOMP, JSOP, JSOMS, JSOM, and JAMI. Published by Elsevier Ltd. This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International license.

**(URL)**

<https://hdl.handle.net/20.500.14094/0100490886>



## **Case report**

### **Segmental mandibulectomy for mandibular osteoradionecrosis in an older adult with underweight status**

Junya Kusumoto <sup>a\*</sup>, Yuka Hayase <sup>a</sup>, Yuriko Susukida <sup>a,b</sup>, Takumi Sato <sup>a</sup>, Akiko Sakakibara <sup>a,c</sup>, Masaya Akashi <sup>a</sup>

<sup>a</sup> Department of Oral and Maxillofacial Surgery, Kobe University Graduate School of Medicine, 7-5-2, Kusunoki-cho Chuo-ku, Kobe, 650-0017, Japan

<sup>b</sup> Department of Oral and Maxillofacial Surgery, Kakogawa Central City Hospital, 439, Hon-machi, Kakogawa-cho, Kakogawa, 675-8611, Japan

<sup>c</sup> Department of Oral and Maxillofacial Surgery, Mitsubishi Kobe Hospital, 6-1-34, Wadamiyadori, Hyogo-ku, Kobe, 652-0863, Japan

## **Acknowledgments**

We would like to thank Editage ([www.editage.com](http://www.editage.com)) for English language editing.

## **Funding**

This research did not receive any specific grants from funding agencies in the public, commercial, or not-for-profit sectors.

## **Declaration of conflicts of interest**

The authors declare no conflicts of interests.

**Data availability**

No data was used for the research described in the article.

**\*Correspondence and reprint requests should be addressed to:**

Junya Kusumoto

Department of Oral and Maxillofacial Surgery, Kobe University Graduate School of Medicine

7-5-2, Kusunoki-cho Chuo-ku, Kobe 650-0017, Japan

Telephone: +81-78-382-6213

FAX: +81-78-382-6229

E-mail: [chivalry\\_2727@people.kobe-u.ac.jp](mailto:chivalry_2727@people.kobe-u.ac.jp)

# **Segmental mandibulectomy for mandibular osteoradionecrosis in an older adult with underweight status**

## **Abstract**

Refractory osteomyelitis can lead to recurrent cellulitis and pathological fractures in patients with advanced mandibular osteoradionecrosis (ORN), the optimal treatment for which remains to be established. In addition, difficulties in intake due to trismus and severe pain caused by mandibular ORN may lead to malnutrition and underweight. Therefore, refeeding syndrome (RFS) should be considered when highly invasive surgical procedures are performed. In this report, we discuss the case of an 80-year-old woman with underweight status who had undergone radiotherapy for oropharyngeal malignant melanoma, following which she developed severe pain and a pathological fracture in the right mandible. Given her advanced age and cognitive decline, decision-making regarding the treatment plan was difficult. After repeated consultations with the patient and her family members, the treatment priorities were established as pain control, infection control, and increased mouth opening. A segmental mandibulectomy was performed. After nutritional intake was initiated postoperatively, a marked decrease in serum potassium and phosphorus levels was observed. Because there were no symptoms suggestive of RFS, imminent RFS was considered. Appropriate management in accordance with the National Institute for Health and Care Excellence guidelines prevented progression to RFS. Overall, surgery was successful in achieving the initial treatment objectives, and the patient exhibited a general improvement in quality of life. Careful perioperative management for RFS prevention should be considered necessary when performing surgical procedures for advanced mandibular ORN, especially in older adults.

25    **Keywords:** hypophosphatemia, osteoradionecrosis, refeeding syndrome, segmental  
26    mandibulectomy, underweight

27

## 1. Introduction<sup>1</sup>

Refractory osteomyelitis, which occurs in patients with osteoradionecrosis (ORN) of the jaw and medication-related osteonecrosis of the jaw, can lead to recurrent cellulitis and pathological fractures in advanced stages. In addition, advanced refractory osteomyelitis can lead to trismus and severe pain, feeding problems, chronic malnutrition, and weight loss (Fig. 1). Because refractory osteomyelitis is most common in older adults, decision-making regarding treatment plans is often difficult, especially since the optimal treatment strategy has not been established. Complications after radiation therapy include taste disorders, decreased saliva production, and trismus, which increase the risk of eating disorders [1]. As both irradiation and malnutrition are detrimental to wound healing [2, 3], reconstructive surgery with good blood flow is preferable [4].

Refeeding syndrome (RFS) is a potentially fatal condition that can develop in patients with chronic malnutrition when rapid nutritional supplementation is administered [5]. Although a few reports have discussed cases of RFS in the head and neck region [6, 7], to our knowledge, none have focused on patients with refractory osteomyelitis.

In this report, we discuss a case of ORN in an older patient with a pathological fracture who experienced a good outcome following segmental resection despite difficulties in decision-making.

---

<sup>1</sup> Abbreviations: ORN: osteoradionecrosis; RFS: refeeding syndrome; AS: aortic stenosis; BMI: body mass index; ADL: activities of daily living; NICE: National Institute for Health and Care Excellence; NST: nutritional support team; PAD: peripheral artery disease

## 2. Case report

An 80-year-old woman with a history of severe aortic stenosis (AS), hypertension, osteoporosis, dyslipidemia, and coronary artery stenosis presented to our department for the first time with pain in the right mandibular molar region. Eight years before the first visit to our department, she underwent radiotherapy (conventional, 66 Gy/33 Fr.) for malignant melanomas of the oropharynx. Following the diagnosis of ORN, sequestrectomy and saucerization were performed 8 months after the initial visit. Thereafter, although wound dehiscence and mild pain were confirmed, the extent of bone exposure was small, and there were no signs of acute inflammation. Therefore, conservative treatment was recommended. However, 1 year and 4 months after the initial visit, the patient presented with severe pain in the right mandible.

### 2.1 Physical findings

The patient's height, weight, and body mass index (BMI) were 143 cm, 29 kg, and 14.2 kg/m<sup>2</sup>, respectively, and she exhibited independence in activities of daily living (ADL). No abnormal findings were observed in the eyelids or ocular conjunctiva. Chest examination revealed no heart murmurs or abnormal breath sounds. Abdominal examination results were normal, and edema of the extremities was not observed. Although she exhibited mild impaired swallowing function, there were no obvious findings suggestive of aspiration. Cognitive function was moderately impaired (Mini-Mental State Examination: 17/30, disorientation, short-term memory impairment).

### 2.2 Echocardiographic findings

Echocardiographic examination revealed an ejection fraction of 59.5% and severe AS. No pulmonary congestion was observed.

### ***2.3 Blood test findings***

Blood tests revealed mild anemia, as well as low serum levels of sodium and chloride. In addition, the lymphocyte count, serum albumin, and total protein levels were low. The serum potassium and phosphorus levels were within the normal ranges (Table 1).

### ***2.4 Local findings***

Local findings included erythema, swelling, severe pain in the right mandible, and paresthesia in the area innervated by the right inferior alveolar nerve. In addition, trismus (approximately 0.5 finger breadth) and thrush (scarring) at the corners of the mouth made it difficult for the patient to ingest food. Pain assessment revealed allodynia. Bone exposure was observed in the mandibular right first and second molar equivalent areas (Fig. 2A).

### ***2.5 Imaging findings***

Panoramic X-ray and computed tomography findings revealed bone destruction from the right lower quadrant to the right ramus of the mandible and a pathological fracture at the right mandibular angle (Fig. 2B, C, D).

### ***2.6 List of concerns***

The following concerns were raised: 1) pathological fracture of the right mandible with ORN, 2) severe pain, 3) trismus, 4) difficulty in dietary intake, 5) underweight and malnutrition, 6) severe AS, 7) advanced age and cognitive decline, and 8) history of radiation therapy.

### ***2.7 Treatment strategy***

Advanced age, severe AS, underweight, and malnutrition raised concerns regarding the patient's ability to tolerate surgery. In such cases, our institutional policy typically includes segmental



mandibulectomy alone, without reconstructive surgery. Additionally, cognitive decline was observed, making it difficult to obtain informed consent. After repeated consultations with the patient and her family members, the treatment priorities were pain control, infection control, and increased mouth opening. Surgical treatment, nerve destruction, and conservative treatment (periodic irrigation for bone exposure area and administration of antibiotics) were presented as treatment plans, and the characteristics of each were repeatedly explained (Table 2). Finally, a surgical treatment plan was developed.

### ***2.7.1 Surgical treatment***

Sulbactam/ampicillin was chosen for perioperative antimicrobial treatment and as prophylaxis for infectious endocarditis. Segmental mandibulectomy was performed under general anesthesia (range: mandibular right first premolar to the central ramus of the mandible) (Fig. 3A, B). Good hemorrhaging was observed at the amputation edges. For mucosal defects, suture manipulation from the oral cavity was difficult because of inadequate mouth opening (Fig. 3C). Therefore, the wound was closed completely using the Gambee suture technique for intestinal anastomosis via a cervical approach (Fig. 3D). The watertight wound closure could be performed in the oral cavity, and the wound edge was everted on the oral side. The operative time was 3.5 hours, and the hemorrhage volume was small.

### ***2.8 Postoperative course***

The patient was intubated and admitted to the intensive care unit postoperatively. She was extubated on the first postoperative day and was confirmed to have a good respiratory condition. She did not eat on the day of surgery or the first postoperative day. Given her severe AS, dehydration and excessive infusion were carefully managed. Pre-hospitalization interviews revealed a daily intake of 700–800 kcal, and the same level of caloric intake was planned

postoperatively. Accordingly, 40 kcal was administered via infusion, and 600 kcal of enteral nutrition was started on the second and third postoperative days. Blood tests on the third postoperative day revealed a serum potassium level of 3.0 mmol/L and phosphorus level of 2.0 mg/dL (Table 3), representing significant decreases when compared with preoperative levels; thus, RFS was suspected. In accordance with the National Institute for Health and Care Excellence (NICE) guidelines [8], caloric intake was reduced to approximately 10 kcal/kg/day (300 kcal), and vitamin B<sub>1</sub> supplementation was initiated. Thereafter, caloric intake was increased by 150 kcal every few days with the confirmation of the nutritional support team (NST). Oral intake became possible on the 8th postoperative day, and the final caloric intake was increased to 1,200 kcal without complication (Fig. 4). There were no findings suggestive of RFS, such as arrhythmia or signs of heart failure.

The wound showed no dehiscence or obvious signs of infection (Fig. 5A, B), and the patient was discharged on postoperative day 26. At 12 months postoperatively, the pain had largely resolved, and the infection was under control (Fig. 5C, D). There was no wound dehiscence or bone exposure in the oral cavity, and the wound was completely healed. She exhibited a slight improvement in mouth opening (approximately 1.5 finger breadths) and an increase in body weight (32 kg). She mainly ingested food processed in a blender, with oral enteral nutrition as a supplement. Although rightward deviation of the mandible was observed, the initial treatment objectives were generally achieved overall, and the patient's quality of life improved. She maintained independence in ADL but developed peripheral artery disease (PAD) in her lower limbs, which made walking difficult. Thirteen months after the mandibulectomy, she was admitted to our cardiology department to treat the PAD. Blood tests at 22 days, 5 months, and 13 months postoperatively are also shown in Table 3.

### 3. Discussion

Segmental mandibulectomy was performed in an older patient with underweight status and mandibular ORN. Decision-making regarding the treatment plan was difficult because of the patient's cognitive decline. However, after clarifying the treatment priorities, the characteristics of each treatment method were repeatedly explained and discussed with the patient and her family members, and the treatment decision was made in accordance with the patient's wishes and general condition [9]. Furthermore, the patient was considered at high risk for RFS because of severe emaciation. Therefore, perioperative management was performed to avoid RFS.

RFS is a metabolic disorder caused by the rapid resumption of nutritional administration in patients with malnutrition. RFS is characterized by hypokalemia, hypophosphatemia, and hypomagnesemia, which are associated with the development of neurological symptoms and heart failure. Severe cases can lead to death, necessitating extreme caution [10]. In the NICE guidelines [8], patients at high risk of RFS are defined as follows: 1) one or more of the following: i) BMI less than 16 kg/m<sup>2</sup>; ii) unintentional weight loss of 15% or more in the past 3 to 6 months; iii) minimal or no nutritional intake for more than 10 days; iv) hypokalemia, hypophosphatemia, or hypomagnesemia before refeeding; 2) two or more of the following: i) BMI less than 18.5 kg/m<sup>2</sup>, ii) unintentional weight loss of more than 10% in the past 3 to 6 months, iii) very little or no nutritional intake for more than 5 days; iv) history of alcohol dependence or use of insulin, anticancer drugs, antacids, or diuretics.

As the patient's BMI was 14.2 kg/m<sup>2</sup>, she was thus considered at high risk for RFS. Nevertheless, she had the same caloric intake before and after admission (700–800 kcal). Since there was almost no fasting period and her caloric requirement was approximately 1200 kcal, we believed it was fine to start the initial postoperative nutritional intake at the same level as before surgery while taking RFS into consideration. However, hypokalemia and hypophosphatemia were observed immediately after the patient was started on enteral feeding, and RFS was suspected. At this point, there were no clinical symptoms suggestive of RFS;

therefore, the condition was considered imminent RFS (i.e., one step before RFS) [11]. Thus, even though the patient did not have acute nutritional intake, we believed that early intervention could avoid progression to full RFS. In this case, imminent RFS may have occurred due to chronically biased nutrition from the preoperative period and decreased self-regulation due to old age. Previous reports have also suggested that RFS can occur even after a short period of fasting [12]. In cases such as the present case, where the risk of RFS is high, the intervention of the NST seems necessary prior to surgery and hospitalization.

Populations potentially at high risk of RFS may have decreased intake (dysphagia, anorexia nervosa, mental health disorders, and chronic alcoholism), decreased nutrient absorption (inflammatory bowel disease), or increased metabolic demand (in cancer and surgery) [10, 13, 14]. The incidence rate of RFS among malnourished individuals, mostly including geriatric patients and those with underlying diseases, is estimated to range from 0% to 38% [15]. Most reports of RFS in the head and neck region focused on patients with cancer [16, 17]. Some patients scheduled for oral surgery are at risk of developing RFS due to advanced oral cancer and severe infections that make it difficult to ingest food. In one previous report, the authors noted that surgical treatment with management for RFS prevented the onset of RFS in a patient with severe odontogenic infection [18]. Patients with refractory osteomyelitis who are also candidates for highly invasive surgery are at a higher risk of developing RFS. It is assumed that the risk of RFS is further increased in older patients and those with cognitive decline, as in the present case (Fig. 1). The recognition of RFS in surgery is considered low [7, 18]. RFS should always be considered in older adults undergoing surgical treatment for refractory osteomyelitis, and great care should be taken in perioperative management. In this case, it was difficult to suture the mucosal defect via an oral approach due to severe trismus. As precise suturing was necessary given her post-irradiation status, we adopted the Gambee suture method, which involves inverting the cervical approach into the oral cavity [19]. This allowed

us to close the intraoral wound from the neck, and wound healing occurred without subsequent wound dehiscence.

#### **4. Conclusion**

Advanced refractory osteomyelitis tends to lead to poor nutritional status in older adults due to difficulties in food intake. When surgery is more invasive, careful attention should be paid to perioperative management, including nutritional management for RFS.

#### **Consent for publication**

The patient and her family provided written informed consent for the publication of this report.

#### **Acknowledgements**

None.

## References

- [1] Trotti A. Toxicity in head and neck cancer: a review of trends and issues. *Int J Radiat Oncol Biol Phys.* 2000;47:1–12. [https://doi.org/10.1016/s0360-3016\(99\)00558-1](https://doi.org/10.1016/s0360-3016(99)00558-1).
- [2] Li S, Zhang J, Zheng H, Wang X, Liu Z, Sun T. Prognostic role of serum albumin, total lymphocyte count, and mini nutritional assessment on outcomes after geriatric hip fracture surgery: a meta-analysis and systematic review. *J Arthroplasty.* 2019;34:1287–96. <https://doi.org/10.1016/j.arth.2019.02.003>.
- [3] Zhang Y, Ideguchi H, Aoyagi H, Yamashiro K, Yamamoto T, Nishibori M et al. Malnutrition delayed wound healing after tooth extraction by HMGB1-related prolonged inflammation. *Int Immunopharmacol.* 2021;96:107772. <https://doi.org/10.1016/j.intimp.2021.107772>.
- [4] Lee M, Chin RY, Eslick GD, Sritharan N, Paramaesvaran S. Outcomes of microvascular free flap reconstruction for mandibular osteoradionecrosis: a systematic review. *J Craniomaxillofac Surg.* 2015;43:2026–33. <https://doi.org/10.1016/j.jcms.2015.03.006>.
- [5] Friedli N, Stanga Z, Sobotka L, Culkin A, Kondrup J, Laviano A et al. Revising the refeeding syndrome: results of a systematic review. 2017;35:151–60.
- [6] Mehanna H, Nankivell PC, Moledina J, Travis J. Refeeding syndrome-awareness, prevention and management. *Head Neck Oncol.* 2009;1:4. <https://doi.org/10.1186/1758-3284-1-4>.
- [7] Kaderbay A, Atallah I, Fontaine E, Chobert-Bakouline M, Schmitt S, Mitariu P et al. Malnutrition and refeeding syndrome prevention in head and neck cancer patients: from theory to clinical application. *Eur Arch Otorhinolaryngol.* 2018;275:1049–58. <https://doi.org/10.1007/s00405-018-4935-2>.
- [8] Nutrition support for adults: oral nutrition support, enteral tube feeding and parenteral nutrition. NICE, <https://www.nice.org.uk/guidance/cg32/chapter/introduction>; 2006.

- 236 [9] Bhatt J, Walton H, Stoner CR, Scior K, Charlesworth G. The nature of decision-making in  
237 people living with dementia: a systematic review. *Aging Ment Health*. 2020;24:363–73.  
238 <https://doi.org/10.1080/13607863.2018.1544212>.
- 239 [10] Mehanna HM, Moledina J, Travis J. Refeeding syndrome: what it is, and how to prevent  
240 and treat it. *BMJ*. 2008;336:1495–8. <https://doi.org/10.1136/bmj.a301>.
- 241 [11] Friedli N, Stanga Z, Culkin A, Crook M, Laviano A, Sobotka L et al. Management and  
242 prevention of refeeding syndrome in medical inpatients: an evidence-based and consensus-  
243 supported algorithm. *Nutrition*. 2018;47:13–20. <https://doi.org/10.1016/j.nut.2017.09.007>.
- 244 [12] Rinninella E, D’Angelo M, Borriello R, Galasso T, Cintoni M, Raoul P et al. Incidence  
245 and impact of refeeding syndrome in an internal medicine and gastroenterology ward of an  
246 Italian tertiary referral center: a prospective cohort study. *Nutrients*. 2022;14:1343.  
247 <https://doi.org/10.3390/nu14071343>.
- 248 [13] Ahmed S, Travis J, Nehanna H. Re-feeding syndrome in head and neck – Prevention and  
249 management. *Oral Oncol*. 2011;47:792–6.  
250 <https://doi.org/10.1016/j.oraloncology.2010.06.009>.
- 251 [14] da Silva JSV, Seres DS, Sabino K, Adams SC, Berdahl GJ, Citty SW et al. ASPEN  
252 consensus recommendations for refeeding syndrome. *Nutr Clin Pract*. 2020;35:178–95.  
253 <https://doi.org/10.1002/ncp.10474>.
- 254 [15] Cioffi I, Ponzo V, Pellegrini M, Evangelista A, Bioletto F, Ciccone G et al. The incidence  
255 of the refeeding syndrome. A systematic review and meta-analyses of literature. *Clin Nutr*.  
256 2021;40:3688–701. <https://doi.org/10.1016/j.clnu.2021.04.023>.
- 257 [16] Windpessl M, Mayrbaeurl B, Baldinger C, Tiefenthaller G, Prischl FC, Wallner M et al.  
258 Refeeding syndrome in oncology: report of four cases. *World J Oncol*. 2017;8:25–9.  
259 <https://doi.org/10.14740/wjon1007w>.
- 260 [17] Bhootra K, Bhootra AR, Desai K, Bhootra RK, R MS. Wernike’s encephalopathy as a

261 part of refeeding syndrome. J Assoc Physicians India. 2020;68:80–2.

262 [18] Amemiya T, Fukuoka A, Horiuchi T, Shigematsu H, Sato K, Hamada Y. A case of severe

263 odontogenic infection in which refeeding syndrome could be avoided by intervention of the

264 nutritional support team. Jpn J Oral Maxillofac Surg. 2020;66:399–405. (in Japanese).

265 <https://doi.org/10.5794/jjoms.66.399>.

266 [19] Gambee LP. A single-layer open intestinal anastomosis applicable to the small as well as

267 the large intestine. West J Surg Obstet Gynecol. 1951;59:1–5.

268



## FIGURE LEGENDS

### **Figure 1. Development of malnutrition and underweight due to refractory osteomyelitis and associated factors**

As refractory osteomyelitis progresses, difficulty in dietary intake predisposes patients to trismus and severe pain. Radiation therapy and medication treatment—coupled with the conditions that necessitate them, advanced age, and cognitive decline—can also lead to malnutrition and underweight.

### **Figure 2. Facial and imaging findings during pain enhancement**

A) Redness and swelling were observed in the right mandibular angle. B) Panoramic X-ray revealed a fracture in the right mandibular angle. C) Three-dimensional reconstruction computed tomography (CT) image showing bone destruction and fracture in the right mandibular angle. D) CT image showing bone destruction from the right lower quadrant to the ramus of the mandible.

### **Figure 3. Intraoperative findings and suture schema**

A) Status of the mandible, including the fractured area. The fracture area was identified, and the resection area was set from the right lower fourth to the middle of the mandibular ramus. B) After segmental mandibulectomy, good hemorrhaging was observed from the resection edge. C) Confirmation of connection to the oral cavity at the mucosal defect. D) Scheme at the time of suturing, which was performed using the Gambee suture method.

### **Figure 4. Changes in preoperative and postoperative caloric intake and electrolyte balance**

Enteral feeding was started 2–3 days postoperatively. Serum potassium and phosphorus levels


decreased after the start of enteral nutrition, and serum phosphorus decreased by approximately 39% (2.0 mg/dL) compared with the preoperative level (3.3 mg/dL). The serum magnesium levels remained within the reference values. Oral intake was initiated on postoperative day 8.



Pre-op, pre-operation

Bar chart: total calorie intake (kcal); blue line: serum potassium (mmol/dL); red line: serum phosphorus (mg/dL); black line: serum magnesium (mg/dL).





### **Figure 5. Postoperative wound healing and panoramic X-ray findings**


A) Wound findings on postoperative day 11. The wound was mildly erythematous, but no wound dehiscence or fistulas were observed. B) Panoramic X-ray findings on postoperative day 20. C) Wound findings at 12 months postoperatively. No evidence of infection or fistula was observed. D) Panoramic X-ray findings 12 months after surgery. No abnormal resorption of the bony transection was observed.
















1 **Table 1. Blood test findings on admission**

Variables	
<i>Complete blood count</i>	
White blood cell ( $\times 10^3/\mu\text{L}$ )	5.3
Neutrophil (%)	69.2
Lymphocyte (%)	16.9
Red blood cell ( $\times 10^6/\mu\text{L}$ )	3.5
Hemoglobin (g/dL)	10.6
Hematocrit (%)	31.9
Platelet ( $\times 10^4/\mu\text{L}$ )	36.8
<i>Coagulation examination</i>	
Activated partial thromboplastin time (s)	34.8
Prothrombin time (%)	88.2
PT-INR	1.08
<i>Biochemical examination</i>	
C-reactive protein (mg/dL)	0.18
Aspartate aminotransferase (U/L)	21
Alanine aminotransferase (U/L)	10
$\gamma$ -glutamyl transpeptidase (IU/L)	19
Creatine kinase (U/L)	43
Cholinesterase (U/L)	221
Lactate dehydrogenase (U/L)	196
Sodium (mmol/L)	130
Potassium (mmol/L)	4.4



Chlorine (mmol/L)	96
Calcium (mmol/L)	9.0
Phosphorus (mg/dL)	3.6
Blood urea nitrogen (mg/dL)	11.5
Uretic acid (mg/dL)	3.2
Creatinine (mg/dL)	0.64
eGFR (mL/min/1.73m <sup>2</sup> )	66.4
Total protein (g/dL)	6.1
Albumin (g/dL)	3.3
Total cholesterol (mg/dL)	264
Triglycerol (mg/dL)	107
Glucose (mg/dL)	79
Brain natriuretic peptide (pg/mL)	99.6

- 
- 2 PT-INR, prothrombin time international normalized ratio; eGFR, estimated glomerular
- 3 filtration rate

1 **Table 2. Effectiveness and limitations of each treatment method**

	Surgical treatment	Nerve destruction	Conservative treatment
Pain	I	I or U	U or E
Trismus	U or I	U	U or E
Swallowing function	U or E	U	U
Facial deformity	E	U	U
Infectious control	I or E	U or E	U or E
QOL after treatment	I or E	I or U	U or E
Risk of perioperative complication	Yes	Yes	No

2 QOL, quality of life

3 I, improved; U, unchanged; E, exaggerated

1 **Table 3. Postoperative blood test findings**

Variables	3 days	22 days	5 months	13 months
<i>Complete blood count</i>				
White blood cell ( $\times 10^3/\mu\text{L}$ )	14.6	6.4	6.4	5.8
Neutrophil (%)	90.4	75.6	72.2	
Lymphocyte (%)	3.5	11.9	15.3	
Red blood cell ( $\times 10^6/\mu\text{L}$ )	3.7	2.8	3.6	3.3
Hemoglobin (g/dL)	11.1	8.6	11.0	10.3
Hematocrit (%)	33.6	26.1	33.3	31.3
Platelet ( $\times 10^4/\mu\text{L}$ )	35.3	51.0	29.1	26.5
<i>Biochemical examination</i>				
C-reactive protein (mg/dL)	12.17	0.73	0.14	0.06
Aspartate aminotransferase (U/L)	29	36	31	25
Alanine aminotransferase (U/L)	12	29	17	14
$\gamma$ -glutamyl transpeptidase (IU/L)	17	29	14	11
Creatine kinase (U/L)	364	27	52	46
Cholinesterase (U/L)	201	202	260	
Sodium (mmol/L)	132	127	135	137
Potassium (mmol/L)	3.0	4.6	4.3	4.1
Chlorine (mmol/L)	95	90	98	103
Calcium (mmol/L)	9.1		9.8	9.8
Phosphorus (mg/dL)	2.0		3.8	3.8
Blood urea nitrogen (mg/dL)	5.9	14.3	17.5	19.9
Uretic acid (mg/dL)	2.0	3.9	4.9	4.8

Creatinine (mg/dL)	0.50	0.57	0.83	0.84
eGFR (mL/min/1.73m <sup>2</sup> )	87.0	75.4	49.8	49.2
Total protein (g/dL)	6.3	6.1	6.9	6.5
Albumin (g/dL)	3.2	2.9	4.0	3.3

---

2 eGFR, estimated glomerular filtration rate