

PDF issue: 2025-05-16

# Late intrathecal retraction of a lumboperitoneal shunt

Kim, Young Ju ; Fujita, Atsushi ; Maeyama, Masahiro ; Hori, Tatsuo ; Tanaka, Kazuhiro ; Sasayama, Takashi

(Citation) Surgical Neurology International,14(417):1-4

(Issue Date) 2023-12-08

(Resource Type) journal article

(Version) Version of Record

(Rights) ©2023 Published by Scientific Scholar on behalf of Surgical Neurology International Creative Commons Attribution-Non Commercial-Share Alike 4.0 License

(URL) https://hdl.handle.net/20.500.14094/0100491600



www.surgicalneurologyint.com

### Surgical Neurology International

Editor-in-Chief: Nancy E. Epstein, MD, Professor of Clinical Neurosurgery, School of Medicine, State U. of NY at Stony Brook.

SNI: Spine



Editor

Nancy E. Epstein, MD Professor of Clinical Neurosurgery, School of Medicine, State U. of NY at Stony Brook

## Late intrathecal retraction of a lumboperitoneal shunt

Young Ju Kim, Atsushi Fujita, Masahiro Maeyama, Tatsuo Hori, Kazuhiro Tanaka, Takashi Sasayama

Department of Neurosurgery, Kobe University Graduate School of Medicine, Kobe, Japan.

E-mail: \*Young Ju Kim - yk1023102368@gmail.com; Atsushi Fujita - afujita@med.kobe-u.ac.jp; Masahiro Maeyama - m1982@med.kobe-u.ac.jp; Tatsuo Hori - sprewell.hori@hotmail.co.jp; Kazuhiro Tanaka - kazutana@med.kobe-u.ac.jp; Takashi Sasayama - takasasa@med.kobe-u.ac.jp



Case Report

\***Corresponding author:** Young Ju Kim, Department of Neurosurgery, Kobe University Graduate School of Medicine, Kobe, Japan.

ScientificScholar<sup>®</sup>

Publisher of Scientific Journals

Knowledge is power

yk1023102368@gmail.com

Received: 05 September 2023 Accepted: 10 November 2023 Published: 08 December 2023

DOI 10.25259/SNI\_742\_2023

Quick Response Code:



### ABSTRACT

**Background:** Lumboperitoneal (LP) shunt placement is a good option for treating elderly patients with communicating normal pressure hydrocephalus (NPH) who are also on antiplatelet therapy following endovascular treatment of unruptured bilateral internal carotid artery aneurysms. Here, in an 80-year-old male with an LP shunt, the catheter was "pinched" between adjacent spinous processes, resulting in laceration of the catheter and intrathecal catheter migration.

**Case Description:** An 80-year-old male was treated with a LP shunt for NPH 1 year after undergoing endovascular treatment of unruptured bilateral internal carotid artery aneurysms. The lumbar catheter was placed at the L2–3 level. Six months later, when he clinically deteriorated, the follow-up computed tomography showed recurrent ventricular enlargement. Further, studies additionally confirmed intrathecal migration of the lumbar catheter, warranting secondary ventriculoperitoneal shunt placement.

**Conclusion:** Patients with LP shunts may develop lumbar catheter lacerations secondary to a "pinching" effect from adjacent spinous processes, resulting in intrathecal catheter migration.

Keywords: Catheter rupture, Idiopathic normal pressure hydrocephalus, Intrathecal migration, Lumboperitoneal shunt, Ventricular shunt

### INTRODUCTION

Lumboperitoneal (LP) shunts are good alternatives to ventriculoperitoneal (VP) shunts for patients requiring antiplatelet therapy. Here, one year following endovascular treatment of unruptured bilateral intracranial aneurysms, an 80-year-old patient developed idiopathic normal pressure hydrocephalus (NPH). He initially underwent placement of an LP shunt to avoid the increased hemorrhagic risks of a VP shunt. However, when the catheter became "pinched"/lacerated by the adjacent spinous processes and migrated intrathecally, he ultimately required a VP shunt.

### **CASE REPORT**

An 80-year-old male underwent coiling of bilateral carotid artery aneurysms [Figures 1a-f]. One year later, he developed progressive gait disturbance and cognitive dysfunction consistent with NPH. This was confirmed on brain computed tomography (CT), that showed ventricular enlargement with periventricular lucency. To reduce the risk of dual antiplatelet therapy (i.e., aspirin and clopidogrel) and intracranial hemorrhage associated with placing a VP shunt, the patient received an LP shunt (i.e., at the CT-documented "wider" L2/L3 level) [Figures 2a-c]. The shunt included a programmable valve with plastic connectors and an antisiphon device

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. ©2023 Published by Scientific Scholar on behalf of Surgical Neurology International



**Figure 1:** Angiography after flow diverter placement with coiling of bilateral carotid artery aneurysms. (a-c) Left internal carotid artery. (d-f) Right internal carotid artery. (a and d) Three-dimensional rotational angiography. (b and e) Cone-beam computed tomography. (c and f) Angiography at the 6-month follow-up.



**Figure 2:** Perioperative lumbar images when lumboperitoneal (LP) shunt placed. (a) computed tomography showed no scoliosis, hypertrophy of spinous processes, and/or marked calcification of interspinous ligament. (b) X-ray after LP shunt placement. (c) showed a shunt catheter inserted cephalad through the L2–L3 level.

(pressure setting 4, 110 mmH<sub>2</sub>O; outer diameter of catheter, 1.65 mm).

and within five days, the patient was clinically improved (i.e., Mini-Mental State Examination score improved from 19/30 to 28/30).

#### Postoperative course

Postoperatively, his gait/cognitive function improved, and the ventricles decreased in size. However, six months later, symptoms recurred, and the CT once again showed enlarged ventricles/ periventricular lucency, plus L2 intrathecal migration of the lumbar catheter [Figures 3a-d and 4a-d]. A VP shunt was placed,

### DISCUSSION

#### Incidence of LP shunt intrathecal retraction/migration

An 80-year-old male developed a 6-month delayed rupture of an LP shunt catheter that was originally placed at the L2/L3 level (i.e., migrated intrathecally). Catheter



**Figure 3:** Brain computed tomography (CT). (a) Preoperative-flow diverters. (b) CT Image after recurrent worsening of normal pressure hydrocephalus symptoms/signs following bilateral flow diverter placement; enlarged lateral ventricles/periventricular lucency-anterior horns. (c) Image when symptoms improved after lumboperitoneal shunt. (d) Six months later, the patient deteriorated.



**Figure 4:** Lumbar spine images after shunt retraction. (a) Computed tomography (CT) showed that the shunt catheter retracted/migrated intrathecally. (b) X-ray at the site of catheter retraction. The arrow indicates the location. (c) Sagittal bone window CT scan. (d) Coronal CT of spinous processes.

rupture is described in <1% of patients following LP shunt surgery.<sup>[4]</sup> One study found a 13% incidence of proximal LP shunt migration due to failure of fixation.<sup>[1,3]</sup> Other studies demonstrate the following additional mechanisms for LP shunt catheter migration/failure: increased intra-abdominal pressure, changes in cerebrospinal/respiratory pressures, degenerative changes in the intervertebral disc with reduction of an interspinous space, vertebral deformity, and others.<sup>[5]</sup>

### Choice to initially place an LP shunt, followed secondarily by a VP shunt

We chose to place an LP shunt rather than a VP shunt as our patient was on dual antiplatelet therapy following endovascular treatment of unruptured bilateral internal carotid artery aneurysms. Hudson *et al.* reported that such patients are at increased risk for intracranial hemorrhages following VP shunt placement.<sup>[2]</sup> We placed the proximal LP shunt at the widest L2/L3 level (3.1 mm). Still, the catheter presumably became "lacerated/pinched" by the adjacent spinous processes and migrated intrathecally, warranting secondary placement of a VP shunt. Obtaining a preoperative CT to document a shortened distance between lumbar spinous processes may have convinced us to place a VP rather than an LP shunt primarily. In addition, paramedian rather than midline placement of the proximal lumbar end of the LP shunt catheters may limit/avoid such catheter ruptures.<sup>[6]</sup>

### CONCLUSION

Multiple factors go into choosing an LP rather than a VP shunt for treating elderly patients with NPH. Obtaining preoperative CT studies to document sufficient interspinous space/distance may avoid future lumbar proximal catheter lacerations and intrathecal retraction.

### Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

### Financial support and sponsorship

Nil.

### **Conflicts of interest**

There are no conflicts of interest.

### Use of artificial intelligence (AI)-assisted technology for manuscript preparation

The authors confirm that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

### REFERENCES

- 1. AlBakry A, Taha MM, Al Menshawy HA. Cranial migration of lumboperitoneal shunt: A case report and review of literature. Surg Neurol Int 2019;10:124.
- 2. Hudson JS, Nagahama Y, Nakagawa D, Starke RM, Dlouhy BJ, Torner JC, *et al.* Hemorrhage associated with ventriculoperitoneal shunt placement in aneurysmal subarachnoid hemorrhage patients on a regimen of dual

antiplatelet therapy: A retrospective analysis. J Neurosurg 2018;129:916-21.

- 3. Jusue-Torres I, Lu J, Robison J, Hoffberger J, Wemmer J, Sanyal A, *et al.* Diagnostic assessment of adult hydrocephalus log compared to standard normal pressure hydrocephalus diagnostic tools. Fluids Barriers CNS 2015;12:O44.
- 4. Kazui H, Miyajima M, Mori E, Ishikawa M, SINPHONI-2 Investigators. Lumboperitoneal shunt surgery for idiopathic normal pressure hydrocephalus (SINPHONI-2): An openlabel randomised trial. Lancet Neurol 2015;14:585-94.
- 5. Kikuchi A, Maegawa T, Arai N, Yamazaki K, Arai N, Chernov M, *et al.* Broken lumboperitoneal shunt with intra-thecal migration of the lumbar catheter in a patient with idiopathic normal pressure hydrocephalus: Case report. Curr Pract Neurosurg 2021;31:e12-8.
- Tucker A, Kajimoto Y, Ohmura T, Ikeda N, Furuse M, Nonoguchi N, *et al.* Fluoroscopic-guided paramedian approach for lumbar catheter placement in cerebrospinal fluid shunting: Assessment of safety and accuracy. Oper Neurosurg (Hagerstown) 2019;16:471-7.

How to cite this article: Kim YJ, Fujita A, Maeyama M, Hori T, Tanaka K, Sasayama T. Late intrathecal retraction of a lumboperitoneal shunt. Surg Neurol Int 2023;14:417. doi: 10.2559/SNI\_742\_2023

### Disclaimer

The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Journal or its management. The information contained in this article should not be considered to be medical advice; patients should consult their own physicians for advice as to their specific medical needs.