



Circularly polarized gravitational waves in Chern-Simons gravity originating from an axion domain wall

Kanno, Sugumi
Soda, Jiro
Taniguchi, Akira

(Citation)

Physical Review D, 108(8):083525

(Issue Date)

2023-10-20

(Resource Type)

journal article

(Version)

Version of Record

(Rights)

Creative Commons Attribution 4.0 International license

(URL)

<https://hdl.handle.net/20.500.14094/0100491850>



Circularly polarized gravitational waves in Chern-Simons gravity originating from an axion domain wall

Sugumi Kanno¹, Jiro Soda^{2,3} and Akira Taniguchi¹

¹*Department of Physics, Kyushu University, Fukuoka 819-0395, Japan*

²*Department of Physics, Kobe University, Kobe 657-8501, Japan*

³*International Center for Quantum-field Measurement Systems for Studies of the Universe and Particles (QUP), KEK, Tsukuba 305-0801, Japan*



(Received 18 April 2023; accepted 28 September 2023; published 20 October 2023)

We study a scattering problem of gravitational waves (GWs) by an axion domain wall in Chern-Simons (CS) gravity. We find that a circular polarization of GWs is produced after passing through the domain wall. It turns out that the circular polarization is sizable if the frequency of the GW is comparable to a critical value determined by the characteristic CS length scale and the energy scale of the axion domain wall. Thus, observations of the circular polarization could give a stringent constraint on the characteristic CS length scale or could be a new avenue to search for axions.

DOI: [10.1103/PhysRevD.108.083525](https://doi.org/10.1103/PhysRevD.108.083525)

I. INTRODUCTION

Gravitational waves (GWs) are a useful probe of new physics. In string theory, axions [1–4] or axionlike particles (ALPs) [5] are ubiquitous. We call the ALPs axion hereafter. Since the axion is pseudoscalar, it typically couples with GWs through Chern-Simons (CS) term [6,7]. If the axion has an expectation value, the CS term often induces circular polarization of GWs [8]. For example, axion inflation in CS gravity can produce circularly polarized primordial GWs [9], and the circular polarization is shown to be enhanced in the presence of the Gauss-Bonnet term [10], violation of the null-energy condition [11], or bouncing universe [12]. The axion is also a candidate for dark matter [13–15]. Studies of circular polarization of GWs propagating in homogeneous axion dark matter are initiated in [16,17] but it was pointed out that the circular polarization is not large enough by using LIGO data or in a realistic halo profile by subsequent studies [18–20]. In this paper, we investigate circular polarization of GWs passing through a domain wall which consists of the axion. In this case, the axion becomes inhomogeneous in the background in contrast to the cosmological homogeneous axion studied so far. We analyze a scattering problem and study how the axion domain wall induces the circular polarization of GWs.

The formation of domain walls in the early Universe is an important phenomenon [21]. When the Peccei-Quinn

(PQ) symmetry is spontaneously broken after inflation, domain walls are formed [22]. The energy density of the domain walls formed in such scenarios is of the order of $10^9 \sim 10^{12}$ GeV [23,24]. However, since their energy density decays too slowly relative to the energy density of the surrounding matter, they overclose the universe [25]. This is a notorious domain wall problem. On top of the domain wall problem, cosmic microwave background (CMB) observations give a stringent constraint on the surface energy density of domain walls $\sigma < (0.93 \text{ MeV})^3$ at the 95% confidence level for the standard Λ -CDM cosmology [26]. Thus, the domain walls formed via the spontaneous breaking of the PQ symmetry after inflation seem to be ruled out by the CMB observation.

However, several ideas to resolve the domain wall problem are proposed. For instance, one can simply consider preinflationary breakdown of the PQ symmetry. In [27], the authors consider a tilt of the potential or biased initial conditions. Interestingly, in the scenario [28,29], the domain wall does not have the domain wall problem because the energy density of the domain wall decreases faster than that of radiation.

Let us consider the domain walls from the perspective of CS gravity. The coupling constant between axions and CS gravity is a free parameter from the theoretical point of view. On the other hand, the strength of the coupling can be constrained by measurements of the frame dragging of objects orbiting the Earth [30] or by observations of double-binary pulsars [31]. The authors in [32] analyzed the measurements of frame-dragging effects around the Earth by Gravity Probe B in the case of massless axions and gave the upper bound of the characteristic CS length scale ℓ such as $\ell \lesssim 1 \text{ AU} \sim 10^8 \text{ km}$. We here point out that there is

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

no constraint on the strength of the coupling in the case of massive axions.

In this paper, we consider massive axions coupled to the CS gravity under the assumption that the axion domain wall with the surface energy density $\sigma \sim (1 \text{ MeV})^3$ exists within the cosmological horizon at present. We then evaluate the circular polarization of GWs when they pass through the domain wall. Then, we argue the observability of the circular polarization in this setup, and discuss the implications of circular polarization for the coupling constant between the axion and CS gravity. If the circular polarization is observed, it would be a new avenue to search for axion.

The paper is organized as follows. In Sec. II, we present the setup and derive the equation of motion (EOM) of GWs in CS gravity. In Sec. III, we solve the EOM of GWs passing through the domain wall. In Sec. IV, we discuss phenomenological implications. In particular, we argue how to constrain the coupling constant. The final section is devoted to conclusion. In the Appendix, the derivation of basic equations is shown. We work in the natural unit: $\hbar = c = 1$.

II. GWs PASSING THROUGH A DOMAIN WALL IN CS GRAVITY

In this section, we consider GWs propagating in the background of an axion domain wall. We first present a background solution of the axion domain wall and then consider the GWs in the background.

A. Axion domain wall background

The action of the axion field ϕ is given by

$$S = - \int d^4x \sqrt{-g} \left[\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + V(\phi) \right], \quad (2.1)$$

where $V(\phi)$ is a double-well potential with two minima located at $\phi = \pm\eta$ such as

$$V(\phi) = \frac{\lambda}{4} (\phi^2 - \eta^2)^2, \quad (2.2)$$

where λ is a coupling constant and the axion mass is $\sqrt{\lambda}\eta$. Note that the action is parity even because of the quadratic form of the ϕ even if the ϕ is pseudoscalar. We assume that the domain wall is static and planar. Then, without loss of generality, the planar domain wall is assumed to be in the (x, y) -plane and it is orthogonal to the z -axis. In this case, the variation of the action (2.1) with respect to ϕ gives

$$\frac{d^2 \phi(z)}{dz^2} = \lambda (\phi^2(z) - \eta^2) \phi(z). \quad (2.3)$$

If we impose the boundary condition

$$\phi(\pm\infty) = \pm\eta, \quad (2.4)$$

the solution is given by [21]

$$\phi(z) = \eta \tanh \left(\sqrt{\frac{\lambda}{2}} \eta z \right). \quad (2.5)$$

By using this solution, the surface energy density is calculated as

$$\sigma = \int dz \phi'^2(z) \sim \sqrt{\lambda} \eta^3. \quad (2.6)$$

In the next subsection, we consider the situation in which GWs propagate in the z -direction and cross the domain wall. In our setup, we ignore the backreaction of the domain wall to the spacetime.

B. Gravitational waves in Chern-Simons gravity

Let us consider GWs in the Minkowski space expressed by the tensor mode perturbation in the three-dimensional metric,

$$ds^2 = -dt^2 + (\delta_{ij} + h_{ij}) dx^i dx^j, \quad (2.7)$$

where δ_{ij} are Kronecker delta. The metric perturbation h_{ij} satisfies the transverse traceless gauge conditions $h^i_{j,i} = h^i_i = 0$. The indices (i, j) run from 1 to 3, and $(1, 2, 3) = (x, y, z)$.

The action we consider is Chern-Simons gravity expressed by

$$S = \frac{M_p^2}{2} \int d^4x \sqrt{-g} R + \frac{M_p \ell^2}{8} \int d^4x \sqrt{-g} \phi R \tilde{R}, \quad (2.8)$$

where $M_p^2 = 1/(8\pi G)$, ℓ is a length scale characterizing the coupling strength between the axion field and Chern-Simons gravity which refers to as the characteristic CS length scale in the following and $R\tilde{R} = 1/2 \epsilon^{\rho\sigma\alpha\beta} R^{\mu\nu}_{\alpha\beta} R_{\nu\mu\rho\sigma}$. Here, the four-dimensional Levi-Civita tensor $\epsilon^{\rho\sigma\alpha\beta}$ is defined by $\epsilon^{0123} = -1/\sqrt{-g}$. Note that the action is parity invariant because the parity-odd Chern-Simons gravity $R\tilde{R}$ couples with the pseudoscalar ϕ . Substituting the metric Eq. (2.7) into the action Eq. (2.8), we obtain the action of quadratic form of h_{ij} ,

$$\begin{aligned} {}^{(2)}S = & \frac{M_p^2}{8} \int d^4x \left[\dot{h}^{ij} \dot{h}_{ij} - h^{ij,k} h_{ij,k} \right. \\ & \left. - \frac{\ell^2}{M_p} \epsilon^{izk} \partial_z \phi \{ \ddot{h}^m_i h_{km} + \dot{h}_{im,\ell} (h^\ell_k{}^{,m} - h^m_k{}^\ell) \} \right], \end{aligned} \quad (2.9)$$

where a dot denotes the derivative with respect to the time and ϵ^{ijk} is a three-dimensional Levi-Civita symbol. Note that ϕ depends only on z . The variation of the action with respect to h_{ij} gives the equation of motion for GWs in CS gravity,

$$\begin{aligned}\square h^{ij} &= \frac{\ell^2}{2M_p} \varepsilon^{zik} [\phi' \square + \phi'' \partial_z] \dot{h}^j_k \\ &+ \frac{\ell^2}{2M_p} \varepsilon^{zjk} [\phi' \square + \phi'' \partial_z] \dot{h}^i_k,\end{aligned}\quad (2.10)$$

where a prime denotes the derivative with respect to z . Since we assumed that GWs propagate along the z -axis, the wave vector is expressed as $k^\mu = (\omega, 0, 0, \omega)$ in the asymptotic region. We introduce the polarization tensors for the right-handed and left-handed circularly polarized modes as

$$e_{ij}^{(R)} = \begin{pmatrix} 1 & i & 0 \\ i & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad e_{ij}^{(L)} = \begin{pmatrix} 1 & -i & 0 \\ -i & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (2.11)$$

Note that we have relations

$$\varepsilon^{zpk} e_{ik}^{(R)} = i e_{ip}^{(R)}, \quad \varepsilon^{zpk} e_{ik}^{(L)} = -i e_{ip}^{(L)}. \quad (2.12)$$

Using these polarization tensors, the GWs can be decomposed into the amplitude and the polarization such as

$$h_{ij}(t, z) = h_R(t, z) e_{ij}^{(R)} + h_L(t, z) e_{ij}^{(L)}, \quad (2.13)$$

where $h_R(t, z)$ and $h_L(t, z)$ are the amplitude of the right-handed and left-handed modes respectively. Substituting Eq. (2.13) into Eq. (2.10), we obtain

$$\square h_{R/L}(t, z) = \pm i \frac{\ell^2}{M_p} [\phi' \square + \phi'' \partial_z] \dot{h}_{R/L}(t, z). \quad (2.14)$$

The $\pm$ correspond to the right-handed and the left-handed mode, respectively. By using the time translation symmetry in Eq. (2.14), we can write them by the Fourier mode $h_{R/L}(t, z) = H_{R/L}(z) e^{i\omega t}$. Then Eq. (2.14) is written as

$$\begin{aligned}\left(1 \pm \frac{\omega \ell^2}{M_p} \phi'\right) H''_{R/L} \pm \frac{\omega \ell^2}{M_p} \phi'' H'_{R/L} \\ + \omega^2 \left(1 \pm \frac{\omega \ell^2}{M_p} \phi'\right) H_{R/L} = 0.\end{aligned}\quad (2.15)$$

Apparently, right-handed and left-handed modes obey different equations. Thus, the circular polarization is expected to be produced after passing through the domain wall. Curiously, the above equations have been derived in the context of condensed matter physics [33].

III. CIRCULAR POLARIZATION INDUCED BY A DOMAIN WALL

In this section, we derive the effective potential for the GWs. Then we give a formula of reflection coefficients, transmission coefficients, and circular polarization.

A. Effective potential for GWs

If we change the variables in Eq. (2.15) such as

$$F(z) \equiv 1 \pm \frac{\omega \ell^2}{M_p} \phi'(z), \quad H_{R/L}(z) \equiv \frac{f_{R/L}(z)}{\sqrt{F(z)}}, \quad (3.1)$$

we obtain the canonical form of Eq. (2.15) given by

$$\left[-\frac{d^2}{dz^2} + V_{\text{eff}}(z)\right] f_{R/L} = \omega^2 f_{R/L}, \quad (3.2)$$

where we can read off the effective potential $V_{\text{eff}}(z)$ of the form

$$V_{\text{eff}} = -\frac{1}{4} \left(\frac{F'}{F}\right)^2 + \frac{1}{2} \frac{F''}{F}, \quad (3.3)$$

where $\phi(z)$ in $F(z)$ is given by (2.5). The effective potential for right-handed and left-handed circular polarizations is depicted in Fig. 1 where the value of the potential vanishes to the far left (region:I) and far right (region:II). Thus, there is no distinction between $f_{R/L}(z)$ and $H_{R/L}(z)$ in these regions because $F(z) = 1$.

B. Reflection and transmission coefficients

We now consider an incident wave from the region I which is scattered by the potential according to Eq. (3.2). Since the potential vanishes in regions I and II, we can express the solution of Eq. (3.2) as

$$f_{R/L}(z) = \begin{cases} \frac{e^{i\omega z}}{\sqrt{2\omega}} + \mathcal{R}_{R/L} \frac{e^{-i\omega z}}{\sqrt{2\omega}} & (\text{region:I}) \\ \mathcal{T}_{R/L} \frac{e^{i\omega z}}{\sqrt{2\omega}} & (\text{region:II}) \end{cases}, \quad (3.4)$$

where $\mathcal{R}_{R/L}$ and $\mathcal{T}_{R/L}$ are the reflection and transmission coefficients. In order to obtain the transmission coefficient




FIG. 1. The effective potentials $V_{\text{eff}}(z)$ are depicted for right-handed and left-handed circular polarization modes. The parameters are set to be $\eta = 0.9$, $\lambda = 2$, and $\omega \ell^2 / M_p = 1$. We see the potential vanishes in the asymptotic region I and II.

$\mathcal{T}_{R/L}$, we need to solve Eq. (3.2) from the region $z < 0$ numerically. However, the reflection coefficient $\mathcal{R}_{R/L}$ in $z < 0$ is unknown *a priori*, so we consider two auxiliary problems. One is to consider an incident wave $e^{i\omega z}/\sqrt{2\omega}$ from the far left in region I and a solution of Eq. (3.2) in region II. The other is to consider a wave $e^{-i\omega z}/\sqrt{2\omega}$ that moves backward in z in region I and a solution of Eq. (3.2) in region II. That is,

$$f_{R/L}^{(+)}(z) = \begin{cases} \frac{e^{i\omega z}}{\sqrt{2\omega}} & (\text{region: I}) \\ A_{R/L}^{(+)} \frac{e^{i\omega z}}{\sqrt{2\omega}} + B_{R/L}^{(+)} \frac{e^{-i\omega z}}{\sqrt{2\omega}} & (\text{region: II}) \end{cases}, \quad (3.5)$$

$$f_{R/L}^{(-)}(z) = \begin{cases} \frac{e^{-i\omega z}}{\sqrt{2\omega}} & (\text{region: I}) \\ A_{R/L}^{(-)} \frac{e^{i\omega z}}{\sqrt{2\omega}} + B_{R/L}^{(-)} \frac{e^{-i\omega z}}{\sqrt{2\omega}} & (\text{region: II}) \end{cases}. \quad (3.6)$$

We can construct the solution $f_{R/L}$ of the original problem Eq. (3.4) as a superposition of the solutions $f_{R/L}^{(+)}$ and $f_{R/L}^{(-)}$. If we impose the condition that no wave from the far right by multiplying it by $-B_{R/L}^{(+)} / B_{R/L}^{(-)}$, we can consider an incoming wave $e^{i\omega z}/\sqrt{2\omega}$ and a wave that moves backwards toward the far left $-(B_{R/L}^{(+)} / B_{R/L}^{(-)})e^{-i\omega z}/\sqrt{2\omega}$. Then the reflection and transmission coefficients in Eq. (3.4) can be obtained as

$$\mathcal{R}_{R/L} = -\frac{B_{R/L}^{(+)}}{B_{R/L}^{(-)}}, \quad \mathcal{T}_{R/L} = A_{R/L}^{(+)} - \frac{B_{R/L}^{(+)}}{B_{R/L}^{(-)}} A_{R/L}^{(-)}. \quad (3.7)$$

Defining the mode function as $v(z) = e^{-i\omega z}/\sqrt{2\omega}$, we can deduce $A_{R/L}^{(\pm)}$ and $B_{R/L}^{(\pm)}$ as follows:

$$A_{R/L}^{(\pm)} = -i[v(z)f'_{R/L}^{(\pm)}(z) - f_{R/L}^{(\pm)}(z)v'(z)], \quad (3.8)$$

$$B_{R/L}^{(\pm)} = i[v^*(z)f'_{R/L}^{(\pm)}(z) - f_{R/L}^{(\pm)}(z)v'^*(z)], \quad (3.9)$$

where an asterisk denotes the complex conjugate. The degree of the circular polarization is defined as

$$\Pi \equiv \frac{|\mathcal{T}_R|^2 - |\mathcal{T}_L|^2}{|\mathcal{T}_R|^2 + |\mathcal{T}_L|^2}. \quad (3.10)$$

IV. IMPLICATIONS FOR COSMOLOGY

We are now in a position to discuss implications for cosmology. First, we need to clarify the condition that a sizable circular polarization of GWs can be produced. From Eq. (2.15), we see that the CS correction to the Einstein gravity comes in the form of $\omega\ell^2\phi'/M_p$. Since the CS gravity has to be small relative to the Einstein gravity, the CS correction has to satisfy




FIG. 2. The plot of the degree of the circular polarization as a function of frequency normalized by the critical frequency. Here, $M_p = 10^{18}$ GeV, $\ell = 10^8$ km, $\eta = 1$ MeV, and $\lambda = 2$, thereby $\omega_c = 10^{-21}$ eV are used. The CS gravity is valid as long as $\omega/\omega_c < 1$. As ω/ω_c approaches the 1, the degree of circular polarization increases.

$$\frac{\omega\ell^2}{M_p}\phi' = \frac{\ell^2}{M_p}\omega\eta^2\sqrt{\frac{\lambda}{2}}\text{sech}^2\left(\sqrt{\frac{\lambda}{2}}\eta z\right) \leq \frac{\ell^2}{M_p}\omega\eta^2\sqrt{\frac{\lambda}{2}} < 1, \quad (4.1)$$

where we used Eq. (2.5). The critical frequency ω_c is obtained from the last inequality

$$\omega_c = \frac{M_p}{\ell^2\eta^2}\sqrt{\frac{2}{\lambda}}. \quad (4.2)$$

In Fig. 2, we plotted the degree of the circular polarization Π for the parameters $\ell = 10^8$ km, $\eta = 1$ MeV, and $\lambda = 2$. We see that a sizable circular polarization can be produced. In particular, it becomes maximum when the frequency of GWs satisfies $\omega = \omega_c$, that is, when the CS correction gets comparable to Einstein gravity.

The domain wall surface energy density is constrained as $\sigma \lesssim (\text{MeV})^3$ by the CMB power spectrum [26]. Then we find $\eta \sim \text{MeV}$ at most by using Eq. (2.6). In [28], the authors consider a model where the domain wall tension decreases with the expansion of the Universe and satisfies this constraint. If we assume the coupling constant $\sqrt{\lambda} \sim \mathcal{O}(1)$ and use $M_p = 10^{18}$ GeV, the constraint of Eq. (4.1) is written as

$$\omega\ell^2 < 10^{15} \text{ eV}^{-1}. \quad (4.3)$$

Since the characteristic CS scale ℓ is free from constraints by observations in our model of massive axion, let us take $\ell \sim 10^8$ km just for reference. Then Eq. (4.3) gives $\omega_c \sim 10^{-21}$ eV ~ 1 μHz . We should observe circular polarization around $\omega_c \sim 1$ μHz . If the circular polarization

was not observed, it would mean that ℓ has to be smaller than 10^8 km. Repeating this procedure by decreasing ℓ , we can get more stringent constraints on ℓ . Note that decreasing ℓ corresponds to increasing ω_c . If the circular polarization of GWs was observed at a frequency, say $\omega_c = 10$ GHz, it would imply the existence of domain walls and the characteristic CS scale is around 1 km. Then, it will be important to probe such high frequency GWs [34–40].

Although we supposed that GWs are scattered by the domain walls at present, the scattering could have occurred in the early Universe. If so, the GWs having passed through the domain wall should be red shifted at present. Then we need to take into account that the frequency ω of the GWs becomes higher in the past. Thus, we would be able to obtain a stronger constraint on ℓ using Eq. (4.3). It would be also worth studying circular polarization of primordial GWs from the point of view of searching axion. Future space observations [41] would be desired for this purpose.

V. CONCLUSION

In this paper, we studied the circular polarization of GWs in a massive axion domain wall background. We first derived the EOM for the GWs in CS gravity. We solved the EOM numerically and evaluated the degree of the circular polarization. We found that the degree of the circular polarization depends on the frequency of the GW and increases as the characteristic CS length scale ℓ becomes larger. When the effect of the CS correction becomes significant compared to Einstein gravity, the degree of the circular polarization becomes maximum. Depending on the value of the coupling strength of CS gravity ℓ , we may be able to observe the circular polarization that arose out of the axion domain walls. If it happened, observation of the circular polarization would provide us the information about CS gravity and a new avenue to search for axions.

If the circular polarization of GWs is found in a large solid angle of the sky, the origin of the circular polarization could be a condensation of the massive axion that consists of the domain wall. To be consistent with CMB observations, we need to assume that the energy scale of the domain wall η is less than 1 MeV. We argued that we can obtain new constraints on the characteristic CS length scale in CS gravity that consists of massive axion by searching for the circular polarization of the high frequency GWs. Indeed, it would turn out that the massive axion exists and the CS gravity with $\ell \sim 1$ km holds if the circular polarization of GWs around 10 GHz was observed.

ACKNOWLEDGMENTS

S.K. was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grant No. JP22K03621. J.S. was in part supported by JSPS KAKENHI Grants No. JP17H02894, No. JP17K18778, No. JP20H01902, and No. JP22H01220.

APPENDIX A: QUADRATIC ACTION OF GWs IN CS GRAVITY

We derive the quadratic action of CS gravity (2.9). Up to second order in h , the metric and its inverse can be expressed as

$$g_{ij} = \delta_{ij} + h_{ij}, \quad (\text{A1})$$

$$g^{ij} = \delta^{ij} - h^{ij} + h^{il}h_{il}. \quad (\text{A2})$$

The second order perturbations of Chern-Simons term is given by

$$\begin{aligned} [\sqrt{-g}\phi R\tilde{R}]^{(2)} &= \frac{1}{2}\sqrt{-g}\phi\epsilon^{\rho\sigma\alpha\beta}[R^{\mu\nu}{}_{\alpha\beta}R_{\nu\mu\rho\sigma}]^{(2)} \\ &= 2\phi\epsilon^{ijk}([2R^0{}_{i0m}R^{0m}{}_{jk}]^{(2)} + [R^0{}_{ilm}R^{lm}{}_{jk}]^{(2)}), \end{aligned} \quad (\text{A3})$$

where the superscript (2) denotes the second order perturbation. Since the background is Minkowski spacetime, in the transverse traceless gauge, we obtain

$$\epsilon^{ijk}[2R^0{}_{i0m}R^{0m}{}_{jk}]^{(2)} = \epsilon^{ijk}\dot{h}^m{}_{k,j}\ddot{h}_{im}, \quad (\text{A4})$$

$$\epsilon^{ijk}[R^0{}_{ilm}R^{lm}{}_{jk}]^{(2)} = \epsilon^{ijk}\dot{h}_{im,l}(h^l{}_{k,m,j} - h^m{}_{k,l,j}). \quad (\text{A5})$$

It is straightforward to show the following identity

$$\begin{aligned} 2\epsilon^{ijk}(\dot{h}^m{}_{k,j}\ddot{h}_{im} + \dot{h}_{im,l}h^l{}_{k,m,j} - \dot{h}_{im,l}h^m{}_{k,l,j}) \\ = \epsilon^{ijk}\partial_t(\dot{h}^m{}_{k,j}\dot{h}_{im} + h_{im,l}h^l{}_{k,m,j} - h_{im,l}h^m{}_{k,l,j}) \\ + \epsilon^{ijk}\partial_j(\ddot{h}^m{}_i\dot{h}_{km} + \dot{h}_{im,l}h^l{}_{k,m} - \dot{h}_{im,l}h^m{}_{k,l}). \end{aligned} \quad (\text{A6})$$

The reason that the action is expressed in terms of total derivatives is that the CS term is a topological term. In other words, the CS term appears when there is a coupling field ϕ . Thus, the quadratic action reads

$$\begin{aligned} S_{CS}^{(2)} &= \frac{M_p\ell^2}{8} \int d^4x [\sqrt{-g}\phi R\tilde{R}]^{(2)} \\ &= \frac{M_p\ell^2}{8} \int d^4x [\phi\epsilon^{ijk}\partial_t \\ &\quad \times (\dot{h}^m{}_{k,j}\dot{h}_{im} + h_{im,l}h^l{}_{k,m,j} - h_{im,l}h^m{}_{k,l,j}) \\ &\quad + \phi\epsilon^{ijk}\partial_j(\ddot{h}^m{}_i\dot{h}_{km} + \dot{h}_{im,l}h^l{}_{k,m} - \dot{h}_{im,l}h^m{}_{k,l})] \\ &= -\frac{M_p\ell^2}{8} \int d^4x (\partial_z\phi)\epsilon^{izk} \\ &\quad \times (\ddot{h}^m{}_i\dot{h}_{km} + \dot{h}_{im,l}h^l{}_{k,m} - \dot{h}_{im,l}h^m{}_{k,l}), \end{aligned} \quad (\text{A7})$$

where we used the fact ϕ depends only on z .

APPENDIX B: EQUATION FOR GWs IN CS GRAVITY

Now, we derive EOM from the quadratic action. Taking the variation of the action (2.9) with respect the metric perturbation h_{ij} , we obtain

$$\square h^{ij} = -\frac{\ell^2}{2M_p} \varepsilon^{izk} (\partial_z \phi \square \dot{h}_k^j + \partial_z^2 \phi \partial_z \dot{h}_k^j) + (i \leftrightarrow j) \quad (\text{B1})$$

or

$$\square h^{ij} = \frac{\ell^2}{2M_p} \varepsilon^{zik} (\phi' \square + \phi'' \partial_z) \dot{h}_k^j + (i \leftrightarrow j), \quad (\text{B2})$$

where $\square$ denotes the d'Alembert operator, and prime denotes the derivative with respect to z .

APPENDIX C: EFFECTIVE POTENTIAL

We derive the effective potential (3.3). Substituting $H(z)_{R/L} = x(z)f(z)_{R/L}$ into the Eq. (2.15), we obtain,

$$f'' + \left(2\frac{x'}{x} + \frac{F'}{F}\right)f' + \left(\frac{x''}{x} + \frac{F'}{F}\frac{x'}{x} + \omega^2\right)f = 0, \quad (\text{C1})$$

where $F \equiv 1 \pm \omega \ell^2 \phi' / M_p$ and the indices R/L are omitted. In order to transform (C1) into the form of Schrödinger-type equation, the coefficients of f' must vanish, hence we have

$$\frac{x'}{x} = -\frac{1}{2} \frac{F'}{F}. \quad (\text{C2})$$

Without loss of generality, this can be solved as $x = 1/\sqrt{F}$. Substituting (C2) into (C1), we obtain

$$\left[-\frac{\partial^2}{\partial z^2} - \frac{1}{4} \left(\frac{F'}{F}\right)^2 + \frac{1}{2} \frac{F''}{F}\right]f = \omega^2 f. \quad (\text{C3})$$

-
- [1] Jihn E. Kim, Weak interaction singlet and strong CP invariance, *Phys. Rev. Lett.* **43**, 103 (1979).
 - [2] Mikhail A. Shifman, A.I. Vainshtein, and Valentin I. Zakharov, Can confinement ensure natural CP invariance of strong interactions?, *Nucl. Phys.* **B166**, 493 (1980).
 - [3] A.R. Zhitnitsky, On possible suppression of the axion hadron interactions. (In Russian), *Sov. J. Nucl. Phys.* **31**, 260 (1980).
 - [4] Michael Dine, Willy Fischler, and Mark Srednicki, A simple solution to the strong CP problem with a harmless axion, *Phys. Lett.* **104B**, 199 (1981).
 - [5] Asimina Arvanitaki, Savas Dimopoulos, Sergei Dubovsky, Nemanja Kaloper, and John March-Russell, String axiverse, *Phys. Rev. D* **81**, 123530 (2010).
 - [6] Bruce A. Campbell, M. J. Duncan, Nemanja Kaloper, and Keith A. Olive, Gravitational dynamics with Lorentz Chern-Simons terms, *Nucl. Phys.* **B351**, 778 (1991).
 - [7] R. Jackiw and S. Y. Pi, Chern-Simons modification of general relativity, *Phys. Rev. D* **68**, 104012 (2003).
 - [8] Arthur Lue, Li-Min Wang, and Marc Kamionkowski, Cosmological signature of new parity violating interactions, *Phys. Rev. Lett.* **83**, 1506 (1999).
 - [9] Masaki Satoh and Jiro Soda, Higher curvature corrections to primordial fluctuations in slow-roll inflation, *J. Cosmol. Astropart. Phys.* **09** (2008) 019.
 - [10] Masaki Satoh, Sugumi Kanno, and Jiro Soda, Circular polarization of primordial gravitational waves in string-inspired inflationary cosmology, *Phys. Rev. D* **77**, 023526 (2008).
 - [11] Yong Cai, Generating enhanced parity-violating gravitational waves during inflation with violation of the null energy condition, *Phys. Rev. D* **107**, 063512 (2023).
 - [12] Mian Zhu and Yong Cai, Parity-violation in bouncing cosmology, *J. High Energy Phys.* **04** (2023) 095.
 - [13] John Preskill, Mark B. Wise, and Frank Wilczek, Cosmology of the invisible axion, *Phys. Lett.* **120B**, 127 (1983).
 - [14] L. F. Abbott and P. Sikivie, A cosmological bound on the invisible axion, *Phys. Lett.* **120B**, 133 (1983).
 - [15] Michael Dine and Willy Fischler, The not so harmless axion, *Phys. Lett.* **120B**, 137 (1983).
 - [16] Daisuke Yoshida and Jiro Soda, Exploring the string axiverse and parity violation in gravity with gravitational waves, *Int. J. Mod. Phys. D* **27**, 1850096 (2018).
 - [17] Chong-Sun Chu, Jiro Soda, and Daisuke Yoshida, Gravitational waves in axion dark matter, *Universe* **6**, 89 (2020).
 - [18] Sunghoon Jung, TaeHun Kim, Jiro Soda, and Yuko Urakawa, Constraining the gravitational coupling of axion dark matter at LIGO, *Phys. Rev. D* **102**, 055013 (2020).
 - [19] Tomohiro Fujita, Ippei Obata, Takahiro Tanaka, and Kei Yamada, Resonant gravitational waves in dynamical Chern-Simonsaxion gravity, *Classical Quantum Gravity* **38**, 045010 (2021).
 - [20] Takuya Tsutsui and Atsushi Nishizawa, Observational constraint on axion dark matter in a realistic halo profile with gravitational waves, *Phys. Rev. D* **107**, 103516 (2023).
 - [21] Alexander Vilenkin and E. Paul S. Shellard, *Cosmic Strings and Other Topological Defects* (Cambridge University Press, Cambridge, England, 1994).
 - [22] P. Sikivie, Of axions, domain walls and the early universe, *Phys. Rev. Lett.* **48**, 1156 (1982).
 - [23] Takashi Hiramatsu, Masahiro Kawasaki, Ken'ichi Saikawa, and Toyokazu Sekiguchi, Production of dark matter axions from collapse of stringwall systems, *Phys. Rev. D* **85**, 105020 (2012); **86**, 089902(E) (2012).
 - [24] Peter W. Graham, Igor G. Irastorza, Steven K. Lamoreaux, Axel Lindner, and Karl A. van Bibber, Experimental searches for the axion and axion-like particles, *Annu. Rev. Nucl. Part. Sci.* **65**, 485 (2015).

- [25] Ya. B. Zeldovich, I. Yu. Kobzarev, and L. B. Okun, Cosmological consequences of the spontaneous breakdown of discrete symmetry, *Zh. Eksp. Teor. Fiz.* **67**, 3 (1974).
- [26] A. Lazanu, C. J. A. P. Martins, and E. P. S. Shellard, Contribution of domain wall networks to the CMB power spectrum, *Phys. Lett. B* **747**, 426 (2015).
- [27] Sebastian E. Larsson, Subir Sarkar, and Peter L. White, Evading the cosmological domain wall problem, *Phys. Rev. D* **55**, 5129 (1997).
- [28] E. Babichev, D. Gorbunov, S. Ramazanov, and A. Vikman, Gravitational shine of dark domain walls, *J. Cosmol. Astropart. Phys.* **04** (2022) 028.
- [29] E. Babichev *et al.*, NANOGrav spectral index $\gamma = 3$ from melting domain walls, [arXiv:2307.04582](https://arxiv.org/abs/2307.04582).
- [30] Tristan L. Smith, Adrienne L. Erickcek, Robert R. Caldwell, and Marc Kamionkowski, The effects of Chern-Simons gravity on bodies orbiting the earth, *Phys. Rev. D* **77**, 024015 (2008).
- [31] Nicolas Yunes and David N. Spergel, Double binary pulsar test of dynamical Chern-Simons modified gravity, *Phys. Rev. D* **80**, 042004 (2009).
- [32] Yacine Ali-Haïmoud and Yanbei Chen, Slowly-rotating stars and black holes in dynamical Chern-Simons gravity, *Phys. Rev. D* **84**, 124033 (2011).
- [33] A. Martín-Ruiz and L. F. Urrutia, Gravitational waves propagation in nondynamical Chern-Simons gravity, *Int. J. Mod. Phys. D* **26**, 1750148 (2017).
- [34] Nancy Aggarwal *et al.*, Challenges and opportunities of gravitational-wave searches at MHz to GHz frequencies, *Living Rev. Relativity* **24**, 4 (2021).
- [35] Asuka Ito, Tomonori Ikeda, Kentaro Miuchi, and Jiro Soda, Probing GHz gravitational waves with graviton-magnon resonance, *Eur. Phys. J. C* **80**, 179 (2020).
- [36] Asuka Ito and Jiro Soda, A formalism for magnon gravitational wave detectors, *Eur. Phys. J. C* **80**, 545 (2020).
- [37] Asuka Ito and Jiro Soda, Exploring high frequency gravitational waves with magnons, *Eur. Phys. J. C* **83**, 766 (2023).
- [38] Aldo Ejlli, D. Ejlli, A. M. Cruise, G. Pisano, and H. Grote, Upper limits on the amplitude of ultra-high-frequency gravitational waves from graviton to photon conversion, *Eur. Phys. J. C* **79**, 1032 (2019).
- [39] Valerie Domcke, Camilo Garcia-Cely, and Nicholas L. Rodd, Novel search for high-frequency gravitational waves with low-mass axion haloscopes, *Phys. Rev. Lett.* **129**, 041101 (2022).
- [40] Asher Berlin, Diego Blas, Raffaele Tito D'Agnolo, Sebastian A. R. Ellis, Roni Harnik, Yonatan Kahn, and Jan Schütte-Engel, Detecting high-frequency gravitational waves with microwave cavities, *Phys. Rev. D* **105**, 116011 (2022).
- [41] Pau Amaro-Seoane *et al.*, eLISA/NGO: Astrophysics and cosmology in the gravitational-wave millihertz regime, *GW Notes* **6**, 4 (2013).