Kobe University Repository : Kernel

PDF issue: 2026-11-16

A Common Fixed Point Theorem

Iséki, Kiyoshi

(Citation)
Mathematics Seminar Notes,2:50-51

(Issue Date)
1974

(Resource Type)
journal article

(Version)
Version of Record

(JaLCDOI)
https://doi.org/10.24546/0100491903

(URL)
https://hdl.handle.net/20.500.14094/0100491903

KOBE UNIVERSITY

MATHEMATICS SEMINAR NOTES
Vol. 2 (1974) February 1974

A Common Fixed Point Theorem

By Kiyoshi ISÉKI

In this note, we shall prove a fixed point theorem which is a generalization of M. G. Maia's theorem [1].

Theorem. *Let X be a metric space with two metrics d and δ. If X satisfies the following conditions*:

(1) $d(x, y) \leq \delta(x, y)$ *for every* x, y *in* X,

(2) X *is complete with respect to* d,

(3) *two mappings* $f, g: X \to X$ *are continuous with respect to the metric* d, *and*

$$\delta(f(x), g(y)) \leq \alpha\delta(x, y) + \beta[\delta(x, f(x)) + \delta(y, g(y))] + \gamma[\delta(x, g(y)) + \delta(y, f(x))]$$

for every x, y *in* X, *where* α, β, γ *are non-negative and* $\alpha + 2\beta + 2\gamma < 1$, *then* f, g *have a unique common fixed point.*

Proof. Let x_0 be a point in X, put

$$x_1 = f(x_0),\ x_2 = g(x_1),\ \cdots,\ x_{2n} = g(x_{2n-1}),\ x_{2n+1} = f(x_{2n}),\ \cdots.$$

Then

$$\delta(x_n, x_{n+1}) \leq \left(\frac{\alpha + \beta + \gamma}{1 - \beta - \gamma}\right)^n \delta(x_0, x_1).$$

(see I. Rus [2], p.20). Therefore, by $d \leq \delta$,

$$d(x_n, x_{n+1}) \leq \left(\frac{\alpha + \beta + \gamma}{1 - \beta - \gamma}\right)^n \delta(x_0, x_1).$$

This shows that the sequence $\{x_n\}$ is a Cauchy sequence with respect to d. Since X is complete with respect to d, $\{x_n\}$ has a limit point y_0 in X, i.e. $x_n \xrightarrow{d} y_0$. Hence, by the continuity of f with respect to the metric d,

$$y_0 = \lim_{n\to\infty} x_{2n+1} = \lim_{n\to\infty} f(x_{2n}) = f(\lim_{n\to\infty} x_{2n}) = f(y_0).$$

Similary we have $y_0 = g(y_0)$. Therefore y_0 is a common fixed point of f and g.

Let z_0 be a common fixed point of f, g, then

$$\begin{aligned}\delta(y_0, z_0) &= \delta(f(y_0), g(z_0)) \\ &\leq \alpha\delta(y_0, z_0) + 2\gamma\delta(y_0, z_0),\end{aligned}$$

which implies $y_0 = z_0$. Hence f, g have a unique common fixed point in X, which completes the proof.

Remark. In Theorem, let $f(x) = g(x)$, $\beta = \gamma = 0$. then we have a theorem by M. G. Maia [1].

References

[1] M. G. Maia; Un'osservazione sulle contrazioni metriche, Rend. Sem. Mat. Padova, 40 (1968), 139-143.

[2] I. A. Rus: Teoria punctului fix, II, Cluj, 1973.