



Oscillation and Nonoscillation Theorems for Second Order Ordinary Differential Equations

Coffman, C. V.

Wong, J. S. W.

(Citation)

Funkcialaj Ekvacioj, 15(2):119-130

(Issue Date)

1972-08

(Resource Type)

journal article

(Version)

Version of Record

(JaLCD0I)

<https://doi.org/10.24546/0100498872>

(URL)

<https://hdl.handle.net/20.500.14094/0100498872>



Oscillation and Nonoscillation Theorems for Second Order Ordinary Differential Equations

By C. V. COFFMAN* and J. S. W. WONG

(Carnegie-Mellon University & The University of Iowa)

1.

Consider the following second order nonlinear equation

$$(1) \quad y'' + q(x) \operatorname{sgn} y |y|^\alpha = 0, \quad x > 0,$$

where $q(x)$ is continuous and nonnegative for $x > 0$ and α is a positive constant. Equation (1) may be conveniently classified as superlinear or sublinear according to whether the constant α is greater than or less than 1. We are here interested in the oscillatory behavior of solutions of (1), and in particular, in the extension of results concerning equation (1) to the more general equation

$$(2) \quad y'' + yF(y^2, x) = 0,$$

where $yF(y^2, x)$ is continuous for $x > 0$ and $|y| < \infty$, and $F(t, x)$ is nonnegative for $t \geq 0$ and $x > 0$. Accordingly, we say that equation (2) is *superlinear* if $F(t, x)$ satisfies

$$(3) \quad F(t_1, x) \leq F(t_2, x), \quad t_1 \leq t_2,$$

for all x , and it is *sublinear* if $F(t, x)$ satisfies

$$(4) \quad F(t_1, x) \geq F(t_2, x), \quad t_1 \leq t_2,$$

for all x .

Results on the oscillatory behavior of solutions of (2) are of two types, namely, (i) sufficient conditions for all solutions to be oscillatory and for the converse, the existence of a nonoscillatory solution, and (ii) sufficient conditions for all solutions to be nonoscillatory and for the converse, the existence of an oscillatory solution. Here a solution always means a nontrivial solution and it is called *oscillatory* if it has arbitrarily large zeros. On the other hand, a solution is called *nonoscillatory* if it is not oscillatory, i. e. if it is of one sign for all large t . In this paper, we consider both the superlinear and sublinear equations with regard to necessary and sufficient conditions for oscillation and sufficient conditions for nonoscillation. For results concerning sufficient conditions for the existence of an oscillatory solution, we refer the reader to our

* The research of this author was supported by NSF grant GP 12182.

earlier work [3] and [4]. The results in this paper differ from those of [3] and [4] in that we consider only integral conditions rather than monotonicity conditions concerning $F(t, x)$. Although we treat both the superlinear and sublinear equations here, most of the new results are concerned with the sublinear case. Our approach to this problem follows from the study of duality between superlinear and sublinear equations initiated in our latest work [4]. The main results presented below may be considered as genuine extensions of results for equation (1) to the more general equation (2), but our emphasis will be on the duality exhibited between solutions of superlinear and sublinear equations. Such an investigation has not been made even for the simpler equation (1).

The integral conditions of concern to equation (1) are

$$(5) \quad \int^{\infty} xq(x)dx = \infty,$$

and

$$(6) \quad \int^{\infty} x^{\alpha}q(x)dx = \infty.$$

As far as necessary and sufficient conditions for the oscillation of all solutions of (1) are concerned, we have the following two interesting results:

Theorem A. (Atkinson [1]) *Let $\alpha > 1$. All solutions of (1) are oscillatory if and only if (5) holds.*

Theorem B. (Belohorec [2]) *Let $0 < \alpha < 1$. All solutions of (1) are oscillatory if and only if (6) holds.*

Implicit in the proofs of Theorems A and B are the following alternative results:

Theorem A₁. *Let $\alpha > 1$. Equation (1) has a bounded nonoscillatory solution if and only if*

$$(7) \quad \int^{\infty} xq(x)dx < \infty.$$

Theorem B₁. *Let $0 < \alpha < 1$. Equation (1) has an unbounded asymptotically linear solution if and only if*

$$(8) \quad \int^{\infty} x^{\alpha}q(x)dx < \infty.$$

Here by an asymptotically linear solution $y(x)$, we mean a solution for which there exist constants a, b not both zero such that

$$(9) \quad \lim_{x \rightarrow \infty} \frac{y(x)}{ax + b} = 1.$$

Theorem A₁ follows from a more general result of Nehari [9] and Theorem

B_1 will follow from Theorem 2 of this paper. Combining Theorems A and A_1 , we obtain

Theorem A_2 . *Let $\alpha > 1$. The following statements are equivalent :*

- (i) *Equation (1) has a bounded asymptotically linear solution,*
- (ii) *Equation (1) has a nonoscillatory solution,*
- (iii) $\int^{\infty} xq(x)dx < \infty$.

On the other hand, Theorems B and B_1 give.

Theorem B_2 . *Let $0 < \alpha < 1$. The following statements are equivalent :*

- (i) *Equation (1) has an unbounded asymptotically linear solution,*
- (ii) *Equation (1) has a nonoscillatory solution,*
- (iii) $\int^{\infty} x^{\alpha}q(x)dx < \infty$.

Thus from the oscillation and nonoscillation point of view, the following properties may be considered as duals to one another :

(a) the superlinear equation has an asymptotically constant solution, (b) the sublinear equation has an unbounded asymptotically linear solution. Similarly, the integral conditions (7) and (8) become dual to each other. Upon examining Theorem B_1 , we find that the conclusion stated is in fact true even for $\alpha > 1$, a result due to Nehari [9] who proved it for the more general equation (2). Now further manifestation of the duality just described led us to conjecture that Theorem A_1 remains valid for $0 < \alpha < 1$, which is indeed the case (Corollary 1).

The search for sufficient conditions for nonoscillation of (1) offers further evidence of the usefulness of the concept of duality between results for superlinear and sublinear equations. Let us consider the following result :

Theorem C. (Atkinson [1]) *Let $\alpha > 1$ and let $q(x)$ be nonincreasing. Then (8) is sufficient for all solutions of (1) to be nonoscillatory.*

The dual role played by conditions (7) and (8) necessitates the following :

Theorem D. (Heidel [6]) *Let $0 < \alpha < 1$ and let $q(x)$ be nonincreasing. Then (7) is sufficient for all solutions of (1) to be nonoscillatory.*

Further evidence of this duality between properties of solutions of superlinear and sublinear equations may be found in [4].

The extensions of the above mentioned results to the more general equation (2) are the main results of this paper. Although the techniques involved in such extensions are in general rather intricate, the results are quite easy to describe. The generalization of the integral conditions (7) and (8) are

$$(10) \quad \int^{\infty} xF(c^2, x)dx < \infty, \quad c > 0,$$

and

$$(11) \quad \int^{\infty} xF(c^2x^2, x)dx < \infty, \quad c > 0.$$

Theorem A₁ has been generalized by Nehari [9] to equation (2) with the condition that (10) hold for some $c > 0$ replacing condition (7). In [9], Nehari also has introduced the following stronger notion of superlinearity :

$$(12) \quad t_2^{-\epsilon} F(t_2, x) \geq t_1^{-\epsilon} F(t_1, x), \quad t_2 \geq t_1, \quad \epsilon > 0.$$

Under this stronger assumption, we have established the generalization of Theorem A₂ to equation (2) in [4] as follows :

Theorem E. *Let $F(t, x)$ satisfy, for some $\epsilon > 0$, condition (12). The following statements are equivalent :*

- (i) *Equation (2) has a bounded asymptotically linear solution,*
- (ii) *Equation (2) has a nonoscillatory solution,*
- (iii) *For some $c > 0$, (10) holds.*

We introduce, as a dual condition to (12), the "stronger" sublinearity condition : for some $\epsilon > 0$, $F(t, x)$ satisfies

$$(13) \quad t_2^{\epsilon} F(t_2, x) \leq t_1^{\epsilon} F(t_1, x), \quad t_2 \leq t_1.$$

The corresponding result for equation (1) when $F(t, x)$ satisfies the stronger sublinearity condition (13) is given as Theorem 2 in the next section. The condition that $q(x)$ be nonincreasing takes the form

$$(14) \quad F(t, x_1) \geq F(t, x_2), \quad x_1 \leq x_2$$

for all $t > 0$. Theorem C has been generalized to equation (2) again by Nehari [9]. The desired generalization of Theorem D is given below as Theorem 3. The main results concerning oscillation and nonoscillation of solutions of (2) may be summarized in the table given on page 123.

We may consider further improvements of Theorems C and D wherein we relax the monotonicity condition on $q(x)$. In particular, it can be shown that the condition

$$(15) \quad \int^{\infty} \frac{q'_+(x)}{q(x)} dx < \infty,$$

where $q'_+(x) = \max(0, q'(x))$, is sufficient for the validity of Theorems C and D. These results are also available for the more general equation (2).

Portions of Theorems 1 and 2 overlapped with some results of a recent paper by S. Belohorec, Monotone and Oscillatory Solutions of a Class of Non-linear Differential Equations, *Matematicky Casopis*, **19** (1969), 169-187. In particular, the sufficiency part of Theorem 1 and the first statement of Theorem

$F(t, x)$	(10) $\int^{\infty} xF(c^2, x) dx < \infty$	(11) $\int^{\infty} xF(cx^2, x) dx < \infty$
(3)	Necessary and sufficient for existence of a bounded nonoscillatory solution. (Nehari [9])	Necessary and sufficient for existence of an unbounded nonoscillatory solution. (Nehari [9])
(4)	Necessary and sufficient for existence of a bounded nonoscillatory solution (Theorem 1)	Necessary and sufficient for existence of an unbounded asymptotically linear solution (Theorem 2)
(12)	Necessary and sufficient for existence of a nonoscillatory solution (Coffman and Wong [4])	(11) \Rightarrow (10)
(13)	(10) \Rightarrow (11)	Necessary and sufficient for existence of a nonoscillatory solution (Theorem 2)
(3), (14)		Sufficient for nonoscillation (Nehari [9])
(4), (14)	Sufficient for nonoscillation (Theorem 3)	

2 are included in Theorems 1 and 3 of Belohorec's paper respectively. However, the results here are presented in order to emphasize the duality relationships between solutions of superlinear and sublinear equations and our proofs for the overlapping parts differ significantly from that of Belohorec's. We are indebted to Professor J. W. Heidel for bringing this matter to our attention.

The referee has kindly pointed out that the proof of Theorem 1 for the sublinear case resembles that of Theorem 1.1 for the superlinear case as given in the paper, "On the Condition for the Oscillation and Nonoscillation of Solutions of Nonlinear second order Differential Equations", *Differential Equations*, 2 (1966), 814-821, by D. V. Izyumova.

2.

In this section, we state and prove three theorems concerning the sublinear equation (2) and which constitute the main results of this paper. The first is the sublinear analogue of a well known result of Nehari [9, Theorem I].

Theorem 1. *Let $F(t, x)$ satisfy (4). Then equation (2) has a bounded*

nonoscillatory solution if and only if for some $c > 0$, (10) holds.

Proof. Let $y(x)$ be a bounded nonoscillatory solution of (2) and suppose that for $x > x_0 \geq 0$,

$$(16) \quad \frac{c}{2} < y(x) < c.$$

Integrating (2) from 0 to x twice, we obtain

$$(17) \quad y(x) = y(0) + xy'(x) + \int_0^x sy(s)F(y^2(s), s)ds.$$

It is easy to see that $y'(x)$ is positive and non-increasing for $x \geq x_0$, thus

$$(18) \quad y(x) - y(x_0) = \int_{x_0}^x y'(s)ds \geq (x - x_0)y'(x).$$

Since $y(x)$ is nondecreasing and bounded it follows from (18) that $xy'(x)$ remains bounded as x tends to infinity, which in turn implies by (17) that

$$(19) \quad \int_0^\infty sy(s)F(y^2(s), s)ds < \infty.$$

Using condition (4) and (16), we obtain from (19)

$$\frac{c}{2} \int_{x_0}^\infty sF(c^2, s)ds \leq \int_{x_0}^\infty sy(s)F(y^2(s), s)ds < \infty,$$

thus proving the necessity of the condition.

To prove sufficiency, we suppose that condition (10) holds for some positive constant c , and we construct a solution $y(x)$ of (2) which satisfies

$$\lim_{x \rightarrow \infty} y(x) = c',$$

where c' is to be chosen later. Let $c' > c$ and define $y_n(x)$ inductively by

$$(20) \quad \begin{cases} y_0(x) = c' \\ y_n(x) = c' - \int_x^\infty (s-x)y_{n-1}(s)F(y_{n-1}^2(s), s)ds \end{cases}$$

If x_1 is chosen so that

$$c' \int_{x_1}^\infty sF(c^2, s)ds < c' - c,$$

then using (4) and (20), we get for $x \geq x_1$,

$$y_1(x) \leq c',$$

and

$$y_1(x) \geq c' - c' \int_{x_1}^\infty sF(c^2, s)ds > c.$$

Thus, inductively we obtain for $x \geq x_1$ and $n=1, 2, \dots$,

$$y_n(x) \leq c',$$

and

$$y_n(x) \geq c.$$

Using the above, we find for $x \geq x_1$ and all $n=1, 2, \dots$ that

$$(21) \quad \begin{cases} y'_n(x) = \int_x^\infty y_{n-1}(s)F(y_{n-1}^2(s), s)ds \\ \leq c' \int_x^\infty F(c^2, s)ds \leq c' - c. \end{cases}$$

The above estimates for y_n and (21) imply that $\{y_n(x)\}$ forms a uniformly bounded and equicontinuous family, hence it follows from the Arzela-Ascoli theorem that there exists a subsequence $\{y_{n_k}(x)\}$, uniformly convergent on every compact subinterval of $[x_1, \infty)$. Now a standard argument, see for example [10; Theorem 3], yields a function $y(x)$ satisfying

$$y(x) = c' - \int_x^\infty (s-x)y(s)F(y^2(s), s)ds,$$

which is the desired bounded non-oscillatory solution of (2).

As a corollary to the above result, we obtain

Corollary 1. *Let $0 < \alpha < 1$. Equation (1) has a bounded nonoscillatory solution if and only if (7) holds.*

We next prove the desired extension of Theorem B₁.

Theorem 2. *Let F satisfy condition (4). Then equation (2) has an unbounded asymptotically linear solution if and only if (11) holds for some $c > 0$. If F satisfies the stronger condition (13), then (11) is also necessary for the existence of any nonoscillatory solution.*

Proof. We first show that (11) is sufficient for the existence of an unbounded asymptotically linear solution. Choose x_0 such that

$$(22) \quad \int_{x_0}^\infty xF(c^2x^2, x)dx < \frac{1}{4}.$$

Let $c' > c$, then $cx < c'(x-x_0)$ for $x > c'x_0/(c'-c) = x_1$. Thus, in view of (4),

$$\begin{aligned} & \int_{x_0}^\infty (x-x_0)F(c'^2(x-x_0)^2, x)dx \\ & \leq \int_{x_0}^{x_1} (x-x_0)F(c'^2(x-x_0)^2, x)dx + \int_{x_1}^\infty xF(c^2x^2, x)dx \end{aligned}$$

Since $x_1 = c'x_0/(c'-c)$ tends to x_0 as c' tends to infinity, it follows that

$$(23) \quad \int_{x_0}^{\infty} (x-x_0)F(c'^2(x-x_0)^2, x)dx < \frac{1}{4}$$

for sufficiently large c' . Let $y(x)$ be a solution of (2) satisfying

$$(24) \quad y(x_0)=0, \quad y'(x_0)=2c'$$

where x_0 and c' are chosen so that (23) holds. We claim that $y'(x) > c'$ for $x \geq x_0$. Suppose that there exists $x_2 > x_0$ such that $y'(x) > c'$ for $x_0 \leq x \leq x_2$, and $y'(x_2) = c'$. Integrating (2) and using the initial conditions (24), we have

$$(25) \quad y'(x) = 2c' - \int_{x_0}^x y(s)F(y^2(s), s)ds$$

for $x_0 \leq x \leq x_2$. From (25), we know that $y'(x) \leq 2c'$, and since $y(x_0) = 0$, it follows from $y'(x) > c'$ that

$$(26) \quad c'(x-x_0) \leq y(x) \leq 2c'(x-x_0), \quad x_0 \leq x \leq x_2.$$

Using (26) in (25), we obtain

$$(27) \quad y'(x) \geq 2c' \left[1 - \int_{x_0}^x (s-x_0)F(c'^2(s-x_0)^2, s)ds \right].$$

Hence (23) and (27) together imply $y'(x) \geq 3c'/2$, and in particular, $y'(x_2) \geq 3c'/2$ contradicting our assumption. Thus, for all $x \geq x_0$, we have

$$(28) \quad c' < y'(x) \leq 2c'.$$

Since $y(x) \geq 0$ on $[x_0, \infty)$, the integrand in (25) is nonnegative, therefore, $\lim_{x \rightarrow \infty} y'(x)$ exists, and in view of (28) it must be finite and positive. Clearly this implies that y is unbounded and asymptotically linear.

Next we suppose that there exists an unbounded asymptotically linear solution $y(x)$ of (2), i.e. a solution y such that

$$(29) \quad \lim_{x \rightarrow \infty} \frac{y(x)}{x} = c > 0.$$

From (29), it follows that there exists $x_0 > 0$ such that for $x \geq x_0$

$$(30) \quad \frac{c}{2} < \frac{y(x)}{x} < 2c.$$

Integrating (2), we have

$$(31) \quad y'(x) = y'(x_0) - \int_{x_0}^x y(s)F(y^2(s), s)ds.$$

We claim that $y'(x) \geq 0$ for all $x \geq x_0$. Suppose that there exists $x_1 > x_0$ such that $y'(x_1) < 0$. By (2) and (30). We have $y''(x) \leq 0$, thus $y'(x) \leq y'(x_1)$ and $y(x) \leq y(x_1) + y'(x_1)(x-x_1)$ for $x \geq x_1$. Letting $x \rightarrow \infty$, we obtain a contradiction

to (30). Now, $y'(x) \geq 0$ and (31) give

$$\int_{x_0}^{\infty} y(x)F(y^2(x), x)dx < \infty.$$

Using (30) in the above, we obtain

$$\frac{c}{2} \int_{x_0}^{\infty} xF(4c^2x^2, x)dx \leq \int_{x_0}^{\infty} y(x)F(y^2(x), x)dx < \infty$$

proving (11).

Finally, we assume $F(t, x)$ satisfies the stronger condition (13) and wish to show that (11) is also necessary for the existence of any nonoscillatory solution. Let y be a nonoscillatory solution of (2) and let $y(x) > 0$ for $x > x_0$. Observe that

$$(32) \quad \begin{aligned} y(x) &= y(x_0) + \int_{x_0}^x y'(s)ds \\ &\geq \int_{x_0}^x y'(s)ds \geq y'(x)(x - x_0). \end{aligned}$$

On the other hand, we have

$$(33) \quad y(x) \leq y(x_0) + y'(x_0)(x - x_0).$$

From (32) and (33), it follows that there exist constants c_1 and x_1 such that for $x \geq x_1 > x_0$

$$(34) \quad 0 < y(x) \leq c_1x.$$

Using (34), we obtain from the given hypothesis that

$$(35) \quad y^{2\epsilon}(x)F(y^2(x), x) \geq (c_1x)^{2\epsilon}F(c_1^2x^2, x)$$

for all $x \geq x_1$. Consider

$$(36) \quad \begin{aligned} -(y'^{2\epsilon}(x))' &= -2\epsilon(y'(x))^{2\epsilon-1}y''(x) \\ &= 2\epsilon(y'(x))^{2\epsilon-1}y(x)F(y^2(x), x). \end{aligned}$$

Using (35) in (36), we obtain

$$(37) \quad -(y'^{2\epsilon}(x))' \geq 2\epsilon(y'(x))^{2\epsilon-1}(y(x))^{1-2\epsilon}(c_1x)^{2\epsilon}F(c_1^2x^2, x).$$

Choose $x_1 \geq x_0$ so that $x - x_0 \geq x/2$ for all $x \geq x_1$. We may now use (32) to estimate (37) as follows

$$(38) \quad \begin{aligned} -(y'^{2\epsilon}(x))' &\geq 2\epsilon(y'(x))^{2\epsilon-1} \left(y'(x) \frac{x}{2} \right)^{1-2\epsilon} (c_1x)^{2\epsilon} F(c_1^2x^2, x) \\ &\geq (2c_1)^{2\epsilon} x F(c_1^2x^2, x). \end{aligned}$$

Integrating (38) from x_1 to X , we have

$$(39) \quad y'^{2\epsilon}(x_1) - y'^{2\epsilon}(X) \geq (2c_1)^{2\epsilon} \int_{x_1}^X xF(c_1^2x^2, x)dx.$$

Since $y'(x) > 0$ for all large x , (11) follows immediately from (39).

Notice that for the linear equation, Theorem A₁ and Corollary 1, taken together, yield the well known theorem of Bocher, [5, Corollary 9.1, p.380] as well as its converse. Restricting our attention to equation (1), we see in Theorem A₂ the equivalence of (i) and (ii) for $0 < \alpha < 1$ and in Theorem B₂ that of (i) and (ii) for $\alpha > 1$. This observation gives an explanation of why when $\alpha \neq 1$, we can find necessary and sufficient conditions for all solutions of (1) to be oscillatory. This being that, in either case, the existence of any nonoscillatory solution implies the existence of a nonoscillatory solution of a particular type for whose existence even in the linear case one can give necessary and sufficient conditions. Finally, Theorem E shows that such an equivalence remains valid for the more general equation (2) with $F(t, x)$ satisfying the strong superlinearity condition (12) and Theorem 2 shows that the dual statement is true for equation (2) with $F(t, x)$ satisfying (13). Such non-linearity conditions obviously exclude the linear equation in which case, when (7) holds, there exist both unbounded and bounded asymptotically linear (hence nonoscillatory) solutions. This perhaps offers an explanation of why for the linear equation there is no necessary and sufficient conditions for all solutions to be oscillatory. For an explanation of this from an entirely different point of view, see [8, Theorem I] and the remarks which follow.

We now wish to present a generalization of Theorem D to equation (2) under the sublinearity condition (4).

Theorem 3. *Let $F(t, x)$ satisfy (4) and (14). Suppose that for each $c > 0$, (10) holds, then all solutions are nonoscillatory.*

Proof. We note first that condition (4) and (14) imply left uniqueness of the zero solution of (2), (see [4], Appendix). In particular, a solution $y(x)$ satisfying nontrivial initial conditions at some point x_0 has only a finite number of zeros in any bounded interval $[x_0, x_1]$, $x_1 > x_0$. Indeed these facts follow readily from the proof below, but to keep our discussion simple, we shall rely on the results of [4].

Assume now that (2) has an oscillatory solution $y(x)$ and let $x_1 < x_2 < \dots$ denote the consecutive zeros of $y(x)$, then in view of the above observation,

$$(40) \quad \lim_{k \rightarrow \infty} x_k = \infty.$$

Let $\bar{x}_k \in [x_k, x_{k+1}]$ denote the point in that interval where y' vanishes. We wish to show that $|y(\bar{x}_k)| \leq |y(\bar{x}_{k+1})|$, for $k=1, 2, \dots$. For a given value of k we can assume $y(\bar{x}_k) > 0$, then $y(\bar{x}_{k+1}) < 0$, and $y'(x) < 0$ in $[\bar{x}_k, \bar{x}_{k+1}]$. We have then

$$(41) \quad \begin{aligned} (y'(x_{k+1}))^2 &= -2 \int_{\bar{x}_k}^{x_{k+1}} y' y F(y^2, x) dx \\ &= \int_0^{y^2(\bar{x}_k)} F(\lambda, x_1(\lambda)) d\lambda, \end{aligned}$$

similarly,

$$(42) \quad \begin{aligned} (y'(x_{k+1}))^2 &= 2 \int_{x_{k+1}}^{\bar{x}_{k+1}} y' y F(y^2, x) dx \\ &= \int_0^{y^2(\bar{x}_{k+1})} F(\lambda, x_2(\lambda)) d\lambda, \end{aligned}$$

where $x_1(\lambda)$ and $x_2(\lambda)$ are obtained by inverting respectively

$$y^2(x) = \lambda, \quad \bar{x}_k \leq x \leq x_{k+1},$$

and

$$y^2(x) = \lambda, \quad x_{k+1} \leq x \leq \bar{x}_{k+1}.$$

It follows from the definition of $x_1(\lambda)$ and $x_2(\lambda)$ that $x_2(\lambda) > x_{k+1}$, $0 < \lambda < y^2(\bar{x}_{k+1})$ and $x_1(\lambda) < x_{k+1}$, $0 < \lambda < y^2(\bar{x}_k)$. Thus, if we assume $y^2(\bar{x}_{k+1}) < y^2(\bar{x}_k)$, then from (14) we conclude that

$$\int_0^{y^2(\bar{x}_k)} F(\lambda, x_1(\lambda)) d\lambda > \int_0^{y^2(\bar{x}_{k+1})} F(\lambda, x_2(\lambda)) d\lambda,$$

but in view of (41) and (42) this yields a contradiction, thus $y^2(\bar{x}_{k+1}) \geq y^2(\bar{x}_k)$. Consider now the sequence $\{|y(\bar{x}_k)|\}$, which is non-decreasing and hence must tend to a positive limit, finite or infinite. In any case, there exists a constant $c > 0$, and a sequence of zeros of $y'(x)$, say $\{x_k\}$, such that

$$(43) \quad \liminf_{k \rightarrow \infty} y(x_k) = c > 0.$$

From Theorem 1 it follows that there exists a nonoscillatory $z(x)$ which tends monotonically to c from below. Hence there must exist for sufficiently large k points s_1 and s_2 in an interval $[x_k, x_{k+1}]$ such that $y(s_i) = z(s_i)$, $i = 1, 2$, $0 < z(x) < y(x)$ for $x \in (s_1, s_2)$, $y'(s_1) > z'(s_1)$ and $y'(s_2) < z'(s_2)$. Now consider the Wronskian of $y(x)$ and $z(x)$, defined by $w(y, z)(x) = y(x)z'(x) - z(x)y'(x)$. Using (2) and (4), we find, for $s_1 \leq x \leq s_2$,

$$(44) \quad \frac{d}{dx} w(y, z)(x) = y(x)z(x)(F(y^2(x), x) - F(z^2(x), x)) \leq 0.$$

Integrating (44) from s_1 and s_2 , we obtain

$$w(y, z)(s_2) - w(y, z)(s_1) \leq 0,$$

or

$$y(s_2)(z'(s_2) - y'(s_2)) - y(s_1)(z'(s_1) - y'(s_1)) \leq 0.$$

However, the left hand side of (44) is positive, which is impossible. This contradiction proves that all solutions of (1) are nonoscillatory.

Remark 1. It is clear that Theorem D follows from the result above. In fact, we have relaxed the positivity assumption on $q(x)$ to that of non-negativity.

Remark 2. The technique used in connection with the derivation of inequality (42) has been introduced in Moroney [7] in treating a similar problem arisen from a different context.

References

- [1] F. V. Atkinson, "On second order nonlinear oscillation", *Pacific J. Math.*, **5** (1955), 643-647.
- [2] S. Belohorec: "Oscilatoricke riesenia isteje nelinearnej diferencialnej rovnice druheho radu", *Matematicky Casopis*, **11** (1961), 250-255.
- [3] C. V. Coffman and J. S. W. Wong, "On a second order nonlinear oscillation problem", *Trans. Amer. Math. Soc.*, **147** (1970), 357-366.
- [4] C. V. Coffman and J. S. W. Wong, "Oscillation and nonoscillation of solutions of generalized Emden Fowler equations," *Trans. Amer. Math. Soc.*, **167** (1972), 399-434.
- [5] P. Hartman, *Ordinary differential equations*, New York, Wiley, 1964.
- [6] J. W. Heidel, "A nonoscillation theorem for a nonlinear second order differential equation", *Proc. Amer. Math. Soc.*, **22** (1969), 485-488.
- [7] R. M. Moroney, "Note on a theorem of Nehari", *Proc. Amer. Math. Soc.*, **13** (1962), 407-410.
- [8] Z. Nehari, "Oscillation Criteria for second order linear differential equations", *Trans. Amer. Math. Soc.*, **85** (1957), 428-445.
- [9] Z. Nehari, "On a class of nonlinear second order differential equations", *Trans. Amer. Math. Soc.*, **95** (1960), 101-123.
- [10] J. S. W. Wong, "On second order nonlinear oscillation", *Funkcialaj Ekvacioj*, **11** (1969), 207-234.

(Ricevita la 21-an de februaro, 1972)

(Reviziita la 16-an de majo, 1972)