



Perron' s Method for Viscosity Solutions Associated with Piecewise-Deterministic Processes

Lenhart, Suzanne M.

Yamada, Naoki

(Citation)

Funkcialaj Ekvacioj, 34(1):173-186

(Issue Date)

1991-04

(Resource Type)

journal article

(Version)

Version of Record

(JaLCD0I)

<https://doi.org/10.24546/0100499658>

(URL)

<https://hdl.handle.net/20.500.14094/0100499658>



Perron's Method for Viscosity Solutions Associated with Piecewise-Deterministic Processes

By

Suzanne M. LENHART¹ and Naoki YAMADA²

(University of Tennessee, U.S.A. and Kobe University, Japan)

0. Introduction

In this paper, we consider viscosity solutions [2, 4] of integro-differential equations of the following form:

$$(0.1) \quad F\left(x, u(x), \nabla u(x), \int_{\Omega} u(y)Q(dy, x)\right) = 0,$$

where ∇u is the gradient of u and $Q(\cdot, x)$ is a probability measure. The applications for (0.1) considered here are dynamic programming equations associated with the control of piecewise-deterministic (PD) processes.

A PD process follows deterministic dynamics between random jumps, and is determined by its three "local characteristics":

(i) a continuous vector field g , which determines the flow

$$\dot{x}_t = (g_1(x_t), \dots, g_n(x_t))$$

(ii) a jump rate $\lambda: \mathbf{R}^n \rightarrow \mathbf{R}_+$, and

(iii) a transition measure $Q: \bar{\Omega} \rightarrow P(\Omega)$, where $P(\Omega)$ is the set of probability measures on $\bar{\Omega}$.

The extended generator of x_t is

$$(0.2) \quad g(x) \cdot \nabla u + \lambda(x) \int_{\Omega} (u(y) - u(x))Q(dy, x);$$

thus such terms occur in (0.1). See Davis [5] for background and definition of PD processes. For detailed information about applications and control of PD processes see Davis [6], Soner [22], Vermes [23], Gugerli [9], Hordijk and van Duyn Schouten [11], Pliska [18], Rosberg, Variaga and Walrand [21], Pragarauskas [19, 20], Davis, Dempster, Sethi, and Vermes [7], Lenhart [14], Lenhart and Liao [15, 16], and Gatarek [8].

¹ Supported in part by NSF grant DMS-8906226 and University of Tennessee Science Alliance Research Award.

² This work was completed while the second author was visiting the Mathematics Department of the University of Tennessee, Knoxville.

In the case that the state space is Ω , a bounded domain in \mathbf{R}^n , the PD process jumps back into Ω upon hitting the boundary, which leads to the boundary condition to be coupled with equation (0.1),

$$(0.3) \quad u(x) = \int_{\Omega} u(y)Q(dy, x), \quad x \in \partial\Omega.$$

The main goal of this paper is to prove existence results for equation (0.1) with boundary condition (0.2), using Perron's method. Perron's method for viscosity solutions for Hamilton-Jacobi equations was introduced by Ishii [12]. The key idea is that the supremum of a class of subsolutions is a solution. The main advantage of this method is a weakening of assumptions, especially eliminating the "largeness of the discount factor" (zeroth order coefficient) assumption. The assumption was used by Lenhart [14] and Lenhart and Liao [15, 16] in *a priori* estimates necessary for existence of solutions to integro-differential equations for PD processes. With Perron's method, *a priori* $W^{1,\infty}$ estimates are no longer necessary for existence.

After proving a general existence result in section 1, we apply this result in section 2 to examples arising in the control of PD processes. We consider the integro-differential equation with operator (0.2) and the dynamic programming equation associated with optimal stopping control of PD processes. The system associated with the switching control problem for PD processes is treated by an iterative scheme using single equation results.

1. General existence result

In this section we consider the existence of viscosity solutions for equation (0.1) with boundary conditions (0.3). We use Perron's method, which is essentially that "the supremum of a set of subsolution is a solution".

Let Ω be a bounded open set in \mathbf{R}^n with smooth boundary. We make the following assumptions:

- (A.1) $F(x, u, p, r): \Omega \times \mathbf{R} \times \mathbf{R}^n \times \mathbf{R} \rightarrow \mathbf{R}$ is continuous.
- (A.2) If $u_0 \leq u_1$, then $F(x, u_0, p, r) \leq F(x, u_1, p, r)$ for any $(x, p, r) \in \Omega \times \mathbf{R}^n \times \mathbf{R}$.
- (A.3) If $r_0 \leq r_1$, then $F(x, u, p, r_0) \geq F(x, u, p, r_1)$ for any $(x, u, p) \in \Omega \times \mathbf{R} \times \mathbf{R}^n$.
- (A.4) $Q(\cdot, x)$ is a probability measure on Ω for $x \in \bar{\Omega}$, which satisfies

$$\left| \int_{\Omega} v(y)Q(dy, x) \right| \leq C\|v\|_{L^1(\Omega)} \quad \text{for all } v \in L^1(\Omega).$$

- (A.5) The function

$$x \rightarrow \int_{\Omega} v(y)Q(dy, x),$$

is continuous with respect to $x \in \bar{\Omega}$, uniformly for $v \in L^\infty(\Omega)$.

Notice if we fix $v \in L^\infty(\Omega)$, then the equation

$$F\left(x, u(x), \nabla u(x), \int_{\Omega} v(y)Q(dy, x)\right) = 0 \quad \text{in } \Omega$$

is a first order Hamilton-Jacobi equation.

We give some notation necessary to state the definition of viscosity solution. For bounded functions,

$$u^*(x) = \limsup_{r \rightarrow 0} \{u(y) \mid |x - y| < r\} \quad \text{upper semicontinuous envelope of } u$$

and

$$u_*(x) = \liminf_{r \rightarrow 0} \{u(y) \mid |x - y| < r\} \quad \text{lower semicontinuous envelope of } u.$$

Definition. Let u be a bounded function.

(i) u is a viscosity subsolution of (0.1) if

$$F\left(x, u^*(x), \nabla \phi(x), \int_{\Omega} u^*(y)Q(dy, x)\right) \leq 0$$

wherever $u^* - \phi$ attains its maximum for $\phi \in C^1(\Omega)$.

(ii) u is a viscosity supersolution of (0.1) if

$$F\left(x, u_*(x), \nabla \phi(x), \int_{\Omega} u_*(y)Q(dy, x)\right) \geq 0$$

wherever $u_* - \phi$ attains its minimum for $\phi \in C^1(\Omega)$.

(iii) u is a viscosity solution if u is a viscosity sub- and supersolution.

In the following, “(sub/super) solution” means “viscosity (sub/super) solution”.

Assume that there exists a supersolution W of (0.1) such that

$$(1.1) \quad W(x) \geq \int_{\Omega} W_*(y)Q(dy, x) \quad \text{on } \partial\Omega.$$

Define

$$\mathcal{S} = \left\{ v \mid v \text{ is a subsolution of (0.1) such that } v \leq W \text{ in } \Omega \text{ and } v(x) \leq \int_{\Omega} v^*(y)Q(dy, x) \text{ on } \partial\Omega \right\}.$$

Define $u_0(x) = \sup \{v(x) \mid v \in \mathcal{S}\}$.

Perron's method consists of the following two propositions:

Proposition 1.1. *Assume that \mathcal{S} is not empty and F satisfies (A.1)–(A.5), then $u_0 \in \mathcal{S}$.*

Proof. Suppose that $\phi \in C^1$, $u_0^* - \phi$ attains its maximum at $y_0 \in \Omega$. Without loss of generality, we assume

$$(u_0^* - \phi)(y_0) = 0, \quad u_0^* - \phi \leq 0 \quad \text{in } \Omega,$$

and

$$(u_0^* - \phi)(x) \leq -|x - y_0|^2 \quad \text{on } B(y_0, r) \text{ for some } r > 0.$$

Then there exists $x_n \in B(y_0, r)$ such that $x_n \rightarrow y_0$ and $(u_0 - \phi)(x_n) \nearrow 0$. Let $a_n = (u_0 - \phi)(x_n)$. For each n , there exists $u_n \in \mathcal{S}$ such that $(u_n^* - \phi)(x_n) > a_n - 1/n$ and $(u_n^* - \phi)(x) \leq -|x - y_0|^2$ in $\overline{B}(y_0, r)$. Let $y_n \in \overline{B}(y_0, r)$ be such that

$$(u_n^* - \phi)(y_n) = \max_{x \in \overline{B}(y_0, r)} (u_n^* - \phi)(x).$$

Since the chain of inequalities,

$$\begin{aligned} -|y_n - y_0|^2 &\geq (u_0^* - \phi)(y_n) \geq (u_n^* - \phi)(y_n) \\ &\geq (u_n^* - \phi)(x_n) \geq a_n - \frac{1}{n}, \end{aligned}$$

implies that $y_n \rightarrow y_0$, we have

$$\lim_{n \rightarrow \infty} u_n^*(y_n) = \lim_{n \rightarrow \infty} \phi(y_n) = \phi(y_0) = u_0(y_0).$$

Since u_n is a viscosity subsolution, we have

$$F\left(y_n, u_n^*(y_n), \nabla \phi(y_n), \int_{\Omega} u_n^*(y) Q(dy, y_n)\right) \leq 0.$$

Since $u_n \leq u_0$, we can let $n \rightarrow \infty$ to get

$$F\left(y_0, u_0^*(y_0), \nabla \phi(y_0), \int_{\Omega} u_0^*(y) Q(dy, y_0)\right) \leq 0.$$

Also, we have

$$u_0(x) \leq \int_{\Omega} u_0^*(y) Q(dy, x) \quad \text{on } \partial\Omega.$$

Thus $u_0 \in \mathcal{S}$. ■

Proposition 1.2. *Assume $\mathcal{S} \neq \emptyset$. If $v \in \mathcal{S}$ is not a supersolution, then there exists $w \in \mathcal{S}$ such that $v(y) < w(y)$ at some $y \in \Omega$.*

Proof. Suppose $v \in \mathcal{S}$ is not a supersolution. Then there exists $y_0 \in \Omega$, $\tilde{\phi} \in C^1$, and $\eta > 0$ such that $v_* - \tilde{\phi}$ attains its minimum at y_0 and

$$(1.2) \quad F\left(y_0, v_*(y_0), \nabla \tilde{\phi}(y_0), \int_{\Omega} v_*(y) Q(dy, y_0)\right) < -\eta.$$

We may assume $v_*(y_0) = \tilde{\phi}(y_0)$.

We claim that $v_*(y_0) < W_*(y_0)$, the supersolution. If not, $v_*(y_0) = W_*(y_0)$ and $W_* - \tilde{\phi}$ attains its minimum at y_0 . Since W is a supersolution and $v \leq W$,

$$F\left(y_0, v_*(y_0), \nabla \tilde{\phi}(y_0), \int_{\Omega} v_*(y) Q(dy, y_0)\right) \geq 0$$

which is a contradiction.

There exists $\delta_1 > 0$ such that

$$v_*(y_0) + \delta_1 < W_*(x) \quad \text{for } x \in B(y_0, \delta_1).$$

Using (1.2) and (A.1), there exists $\delta_2 > 0$ ($\delta_2 < \delta_1$) such that if x, u, p, \tilde{v} satisfies

$$\begin{aligned} |x - y_0| < \delta_2, \quad |(u - v_*)(y_0)| < \delta_2, \\ |p - \nabla \tilde{\phi}(y_0)| < \delta_2 \quad \text{and} \quad \int_{\Omega} |\tilde{v}(y) - v_*(y)| Q(dy, y_0) < \delta_2, \end{aligned}$$

then

$$F\left(x, u, p, \int_{\Omega} \tilde{v}(y) Q(dy, x)\right) \leq 0.$$

We can find $\delta > 0$ ($\delta < \min\{\delta_2/2, 1\}$) and $\phi \in C^1$ such that

$$\begin{aligned} \phi(y_0) &= \tilde{\phi}(y_0), \quad \nabla \phi(y_0) = \nabla \tilde{\phi}(y_0), \\ |\phi(x) - v_*(y_0)| &< \frac{\delta_2}{2} \quad \text{for } x \in B(y_0, 2\delta), \\ |\nabla \phi(y_0) - \nabla \phi(x)| &< \delta_2 \quad \text{for } x \in B(y_0, 2\delta), \\ \int_{\Omega} |\phi(y) - v_*(y)| Q(dy, y_0) &< \frac{\delta_2}{2}. \end{aligned}$$

We can also choose ϕ and δ to satisfy

$$u_*(x) - \phi(x) > |x - y_0|^2 \quad \text{for } x \in B(y_0, 2\delta).$$

These inequalities imply

$$|\phi(x) + \delta^2 - v_*(y_0)| < \delta_2$$

and

$$\int_{\Omega} |\phi(y) + \delta^2 - v_*(y)| Q(dy, y_0) < \delta_2$$

which give

$$F\left(x, \phi(x) + \delta^2, \nabla\phi(x), \int_{\Omega} (\phi(y) + \delta^2) Q(dy, x)\right) \leq 0 \quad \text{for } x \in B(y_0, 2\delta).$$

This means $\phi(x) + \delta^2$ is a subsolution of (0.1) in $B(y_0, 2\delta)$.

Define

$$w(x) = \begin{cases} \max\{\phi(x) + \delta^2, v(x)\} & \text{for } x \in B(y_0, \delta) \\ v(x) & \text{otherwise.} \end{cases}$$

Note that if $|x - y_0| = \delta$, then $v_*(x) \geq \phi(x) + \delta^2$. By the similar argument in the preceding proposition, we can see that w is a subsolution (note that our equation has a non-local term), and $w \in \mathcal{L}$. But $v(y_0) < w(y_0)$. ■

By Perron's method, we can find a solution u_0 , i.e., u_0^* is a subsolution (u.s.c.) and u_{0*} is a supersolution (l.s.c.). If the comparison principle holds in (0.1), then $u_0^* \leq u_{0*}$. On the other hand, $u_{0*} \leq u_0^*$ in general. Hence $u_{0*} = u_0 = u_0^*$ and u_0 is continuous. The following theorem gives a sufficient condition to assure that u_0 satisfies the boundary condition (0.3).

Theorem 1.1. *Assume (A.1)–(A.5). Suppose that there exists a supersolution W of (0.1) satisfying (1.1), and a continuous solution u_1 of*

$$(1.3) \quad F\left(x, u_1, \nabla u_1, \int_{\Omega} u_0^*(y) Q(dy, x)\right) = 0 \quad \text{in } \Omega$$

with

$$u_1(x) = \int_{\Omega} u_0^*(y) Q(dy, x) \quad \text{on } \partial\Omega.$$

Assume that the comparison principle holds for (0.1), i.e.,

$$u_0 \leq u_1 \text{ on } \partial\Omega \Rightarrow u_0 \leq u_1 \text{ in } \Omega.$$

If $u_1 \leq W$, then u_0 is a continuous solution of (0.1) satisfying the boundary condition (0.3).

Proof. We claim $u_1 \in \mathcal{L}$. Let $\phi \in C^1$ such that $u_1^* - \phi$ attains its maximum at y_0 , then

$$F\left(y_0, u_1(y_0), \nabla\phi(y_0), \int_{\Omega} u_0^*(y) Q(dy, y_0)\right) \leq 0.$$

Using $u_0 \leq u_1$ and (A.3) we have

$$F\left(y_0, u_1(y_0), \nabla\phi(y_0), \int_{\Omega} u_1(y)Q(dy, y_0)\right) \leq 0.$$

Also,

$$u_1(x) = \int_{\Omega} u_0^*(y)Q(dy, x) \leq \int_{\Omega} u_1(y)Q(dy, x) \quad \text{on } \partial\Omega.$$

Hence, we have the claim. By definition of u_0 and $u_0 \leq u_1$, we have $u_0 \equiv u_1$ in $\bar{\Omega}$. Therefore u_0 is a desired continuous solution. ■

2. Applications

In this section we apply Theorem 1.1 to prove existence results for various equations and systems associated with piecewise-deterministic processes.

Let

$$(2.1) \quad Lu(x) = -g(x) \cdot \nabla u(x) + \alpha(x)u(x) - \lambda(x) \int_{\Omega} (u(y) - u(x))Q(dy, x),$$

where \cdot is the usual inner product in \mathbf{R}^n , ∇u is the gradient vector of u and $Q(\cdot, x)$ is a probability measure.

We assume the following conditions:

- (H.1) Ω is a bounded domain with smooth boundary $\partial\Omega$.
- (H.2) $g(x): \Omega \rightarrow \mathbf{R}^n$ is Lipschitz continuous, and, $\alpha(x), \lambda(x): \bar{\Omega} \rightarrow \mathbf{R}$ are continuous.
- (H.3) There exists $\alpha_0 > 0$ such that $\alpha(x) \geq \alpha_0$ for $x \in \bar{\Omega}$.
- (H.4) $\lambda(x) > 0$ for $x \in \Omega$.
- (H.5) $Q(\cdot, x)$ satisfies (A.4) and (A.5).
- (H.6) $g(x) \cdot \eta(x) > 0$ for $x \in \partial\Omega$, where $\eta(x)$ is the outward unit normal at $x \in \partial\Omega$.

Note that hypothesis (H.2) can be weakened by using a more complicated hypothesis (see [3]), but that hypothesis is not the main point here.

Consider

$$(2.2) \quad Lu(x) = f(x) \quad \text{in } \Omega$$

$$(2.3) \quad u(x) = \int_{\Omega} u(y)Q(dy, x) \quad \text{on } \partial\Omega.$$

Theorem 2.1. *Assume (H.1)–(H.6) and that $f \in C(\bar{\Omega})$. Then there exists a unique continuous solution of equation (2.2) satisfying boundary condition (2.3).*

Proof. We apply Theorem 1.1 to

$$F(x, u, p, r) = -g(x) \cdot p + (\alpha(x) + \lambda(x))u - \lambda(x)r.$$

Note that (A.1)–(A.3) are satisfied,

$$w(x) = -\frac{\|f\|_\infty}{\alpha_0} \quad \text{is a subsolution, and}$$

$$W(x) = \frac{\|f\|_\infty}{\alpha_0} \quad \text{is a supersolution.}$$

By Propositions 1.1 and 1.2, there exists a solution u_0 of (2.2) satisfying

$$u_0(x) \leq \int_{\Omega} u_0^*(y)Q(dy, x).$$

Since $\int_{\Omega} u_0^*(y)Q(dy, x)$ is continuous with respect to x ,

$$F\left(x, u, p, \int_{\Omega} u_0^*(y)Q(dy, x)\right)$$

satisfies the conditions (A.1)–(A.5).

Consider the equation

$$-g \cdot \nabla u_1 + (\alpha + \lambda)u_1 - \lambda \int_{\Omega} u_0^*(y)Q(dy, x) = f \quad \text{in } \Omega$$

with Dirichlet boundary condition

$$u_1(x) = \int_{\Omega} u_0^*(y)Q(dy, x) \quad \text{on } \partial\Omega.$$

Comparison principle for this equation is known [2, 3, 4]. By (H.6) and the method of [13], we have the existence of super- and subsolutions. Then there exists a continuous solution u_1 of this equation.

Now applying Theorem 1.1, u_0 is a continuous solution satisfying

$$u_0 = \int_{\Omega} u_0(y)Q(dy, x) \quad \text{on } \partial\Omega.$$

The uniqueness is proved in Lenhart [14]. ■

Now we proceed to apply Theorem 1.1 to the obstacle problem:

$$(2.4) \quad \min \{Lu - f, u - \psi\} = 0 \quad \text{in } \Omega,$$

$$(2.5) \quad u(x) = \int_{\Omega} u(y)Q(dy, x) \quad \text{on } \partial\Omega.$$

Equation (2.4) with boundary condition (2.5) is the dynamic programming equation associated with the optimal stopping problem for PD processes. See [14] for more background.

Theorem 2.2. *Assume (H.1)–(H.6). Suppose that $f, \psi \in C(\bar{\Omega})$. Then there exists a unique continuous solution of equation (2.4) with boundary condition (2.5).*

Proof. Let $W = \max \{ \|f\|_\infty / \alpha_0, \|\psi\|_\infty \}$. Then W is a supersolution and $-W$ is a subsolution of the equation (2.4). Hence by Propositions 1.1 and 1.2, there exists a solution u_0 of (2.4) satisfying

$$(2.6) \quad u_0(x) \leq \int_{\Omega} u_0^*(y) Q(dy, x) \quad \text{on } \partial\Omega .$$

To check the hypothesis of Theorem 1.1, consider the following obstacle problem,

$$(2.7) \quad \min \left\{ -g \cdot \nabla u_1 + (\alpha + \lambda) u_1 - \lambda \int_{\Omega} u_0(y) Q(dy, x) - f, u_1 - \psi \right\} = 0 \quad \text{in } \Omega$$

$$(2.8) \quad u_1(x) = \int_{\Omega} u_0^*(y) Q(dy, x) \quad \text{on } \partial\Omega .$$

Note that the comparison principle holds.

Using (2.6) and $u_0 \geq \psi$ in Ω , the compatibility condition,

$$\psi(x) \leq \int_{\Omega} u_0^*(y) Q(dy, x) \quad \text{on } \partial\Omega$$

is satisfied.

First assume

$$(2.9) \quad h(x) = \int_{\Omega} u_0^*(y) Q(dy, x) \in C^1(\Omega) \cap C(\bar{\Omega})$$

and

$$(2.10) \quad h(x) = \int_{\Omega} u_0^*(y) Q(dy, x) > \psi(x) \quad \text{on } \partial\Omega .$$

In this case, problem (2.7)–(2.8) is equivalent to

$$(2.11) \quad \min \{ -g \cdot \nabla w_1 + (\alpha + \lambda) w_1 - \tilde{f}, w_1 - \tilde{\psi} \} = 0 \quad \text{in } \Omega$$

$$(2.12) \quad w_1(x) = 0 \quad \text{on } \partial\Omega ,$$

where $\tilde{f}, \tilde{\psi}$ satisfy the same properties as f, ψ and $\tilde{\psi} < 0$ on $\partial\Omega$. For simplicity, we write $f = \tilde{f}, \psi = \tilde{\psi}$. We show the existence of a solution to (2.11)–(2.12)

by Perron's method. Indeed, the solution of the linear equation

$$\begin{aligned} -g \cdot \nabla w + (\alpha + \lambda)w &= f & \text{in } \Omega, \\ w &= 0 & \text{on } \partial\Omega \end{aligned}$$

is a subsolution of (2.11)–(2.12).

To construct a supersolution, we follow a barrier construction argument from Oleinik-Radkevich [17] as in Ishii and Koike [13]. Since $\psi < 0$ on $\partial\Omega$, there exists a local barrier, ψ_z in $C(\Omega \cap V_z) \cap C^2(\Omega \cap V_z)$ where $z \in \partial\Omega$, V_z is a sufficiently small neighborhood of z ,

$$\begin{aligned} \psi_z(z) &= 0, \quad \psi_z \geq 0 & \text{on } \overline{\Omega \cap V_z}, \\ \psi_z &\geq \frac{\|f\|_\infty}{\alpha_0} & \text{on } \overline{\Omega} \cap \partial V_z, \\ -g \cdot \nabla \psi_z + (\alpha + \lambda)\psi_z &\geq f & \text{in } \Omega \cap V_z, \quad \text{and} \\ \psi_z &\geq \psi & \text{in } \Omega \cap V_z. \end{aligned}$$

Define

$$\hat{\psi}_z(x) = \begin{cases} \max \left\{ \psi_z(x), \max \left\{ \frac{\|f\|_\infty}{\alpha_0}, \|\psi\|_\infty \right\} \right\} & \text{in } \Omega \cap V_z \\ \max \left\{ \frac{\|f\|_\infty}{\alpha_0}, \|\psi\|_\infty \right\} & \text{otherwise,} \end{cases}$$

and

$$\hat{\psi}(x) = \inf \{ \hat{\psi}_z(x) \mid z \in \partial\Omega \}.$$

Then $\hat{\psi}$ is a supersolution. This implies that there exists a continuous solution of (2.7)–(2.8).

For general continuous boundary values h , not necessarily satisfying (2.9)–(2.10), we choose an approximating sequence $\{h_n\}$ such that $h_n \in C(\overline{\Omega}) \cap C^1(\Omega)$, $h_n > \psi$ on $\partial\Omega$ and $h_n \rightarrow h$ uniformly on $\overline{\Omega}$. Let u_n be a solution of (2.7)–(2.8) associated with boundary value h_n . By standard comparison argument, we have

$$\sup |u_n(x) - u_m(x)| \leq \sup |h_n(x) - h_m(x)|.$$

Hence $\{u_n\}$ converges to some $u \in C(\overline{\Omega})$ and by stability of viscosity solutions, we have that u is a solution of (2.7)–(2.8).

By the comparison result for obstacle problems, we have $u_1 \leq W$. Hence by Theorem 1.1, u_0 satisfies the boundary condition (0.3). The uniqueness follows from Lenhart [14]. ■

As a third example of using Theorem 1.1 to get existence results, we consider a system associated with a switching control of PD processes. We will approximate the system by decoupled equations which we solve by using Theorem 1.1.

A finite collection of PD processes, indexed by $a \in \{1, \dots, m\}$, are controlled in order to minimize the expected value of a cost functional with continuous and switching costs. The associated dynamic programming equation is the following system:

$$(2.13) \quad u = (u^1, \dots, u^m)$$

$$\max \{L^a u^a - f^a, u^a - M^a u\} = 0 \quad \text{in } \Omega,$$

$$(2.14) \quad u^a(x) = \int_{\Omega} u^a(y) Q^a(dy, x), \quad x \in \partial\Omega,$$

where

$$L^a u^a = -g^a \cdot \nabla u^a + \alpha^a u^a - \lambda^a \int_{\Omega} (u^a(y) - u^a(x)) Q^a(dy, x)$$

and

$$M^a u(x) = \min_{b \neq a} \{u^b(x) + k(a, b)(x)\}.$$

See Lenhart [14] and Lenhart and Liao [15] for more background on this control problem and for $W^{1,\infty}$ solutions to this system under stronger assumptions (in particular, α_0 is sufficiently large and the switching costs are constants). There u^a is the minimum cost functions with initial control setting a , meaning that the state process is initially a PD process with dynamics g^a , jump rate λ^a , and transition measure Q^a .

We make the following assumptions for all $a \in \{1, 2, \dots, m\}$.

(H.2)' $g^a: \Omega \rightarrow \mathbf{R}^n$ is Lipschitz continuous and $f^a, \lambda^a, \alpha^a: \Omega \rightarrow \mathbf{R}$ are continuous.

(H.3)' There exists $\alpha_0 > 0$ such that $\alpha^a(x) \geq \alpha_0 > 0$ on Ω .

(H.4)' $\lambda^a > 0$ on $\Omega, f^a \geq 0$ on Ω .

(H.5)' $Q^a(\cdot, x)$ is a probability measure on Ω , satisfying (A.4), (A.5), and

$$Q^a(dy, x) = Q^1(dy, x) \quad \text{for all } x \in \partial\Omega.$$

(H.6)' $g^a \cdot \eta > 0$ on $\partial\Omega$ where $\eta = \eta(x)$ is the outward normal at $x \in \partial\Omega$.

(H.7)' The switching costs are continuous, strictly positive on $\bar{\Omega}$ and strictly subadditive, i.e.,

$$k(a, c) < k(a, b) + k(b, c) \quad \text{for all } a, b, c \in \{1, \dots, m\} \text{ and for all } x \in \bar{\Omega}.$$

The agreement of the measures on $\partial\Omega$ in assumption (H.5)' is necessary

for the compatibility of the boundary condition and the implicit obstacles, $M^a u$. Hereafter, we denote $Q(\cdot, x) = Q^1(\cdot, x)$, $x \in \partial\Omega$ for simplicity.

Theorem 2.3. *Under assumptions (H.1), (H.2)'–(H.7)', there exists a unique solution to system (2.13) satisfying boundary condition (2.14).*

Proof. The comparison and uniqueness results in Lenhart [14] can be applied to the case here.

Define two maps,

$$\sigma, M: (C(\Omega))^m \rightarrow (C(\Omega))^m$$

by

$$M(v) = \left(\min_{b \neq 1} \{v^b + k(1, b)\}, \dots, \min_{b \neq m} \{v^b + k(m, b)\} \right)$$

and $\sigma(v) = w$ where w is the viscosity solution of

$$\begin{aligned} \max \{L^a w^a - f^a, w^a - v^a\} &= 0 \quad \text{in } \Omega, \\ w^a(x) &= \int_{\Omega} w^a(y) Q(dy, x), \quad x \in \partial\Omega \end{aligned}$$

(using Theorem 2.1 to get existence of w). The map $S = \sigma \circ M$ is increasing and concave. Define a supersolution, u_0 , to (2.13) as the solution of

$$\begin{aligned} L^a u_0^a &= f^a \quad \text{in } \Omega \\ \text{with } u_0^a &= \int_{\Omega} u_0^a(y) Q(dy, x) \quad \text{on } \partial\Omega. \end{aligned}$$

There exists $\gamma > 0$ such that

$$\gamma u_0^b \leq \inf_{a,x} k(a, b)(x) \quad \text{for all } b \in \{1, \dots, m\}.$$

Each component of $S(0)$ and u_0 satisfies

$$(2.15) \quad S(0) \geq \sigma(\gamma u_0) \geq \gamma \sigma(u_0) = \gamma u_0.$$

Thus we have the following property:

$$(2.16) \quad \begin{aligned} &\text{if } v - w \leq \beta v \text{ (componentwise) for some } \beta \in [0, 1], \\ &\text{then } Sv - Sw \leq (1 - \gamma)\beta Sv \text{ (componentwise)}. \end{aligned}$$

Property (2.16) is proved by starting with

$$(1 - \beta)v + \beta \cdot 0 \leq w,$$

applying S to both sides, and using concavity and (2.15). See [1, 10] for

similar arguments. Define $u_n = S^n u_0$. Using (2.16), first on

$$u_0 - u_1 \leq u_0 \quad \text{to obtain} \quad u_1 - u_2 \leq (1 - \gamma)u_1$$

and then inductively to obtain

$$(2.17) \quad u_{n-1} - u_n \leq (1 - \gamma)^{n-1} u_0.$$

By (2.17), we conclude the sequence converges uniformly,

$$u_n^k \searrow u^k \quad \text{as } n \rightarrow \infty.$$

The uniform convergence insures the convergence of the integral terms and the implicit obstacles, and that u is the desired solution to (2.13)–(2.14). ■

References

- [1] Bensoussan, A., Optimal impulsive control theory, *Lecture Notes in Control and Information Sci.*, **16** (1980), 17–41.
- [2] Crandall, M. G., Evans, L. C. and Lions, P. L., Some properties of viscosity solutions of Hamilton-Jacobi equations, *Trans. Amer. Math. Soc.*, **282** (1984), 487–502.
- [3] Crandall, M. G., Ishii, H. and Lions, P. L., Uniqueness of viscosity solutions revisited, *J. Math. Soc. Japan*, **39** (1987), 581–596.
- [4] Crandall, M. G. and Lions, P. L., Viscosity solutions of Hamilton-Jacobi equations, *Trans. Amer. Math. Soc.*, **277** (1983), 1–42.
- [5] Davis, M. H. A., Piecewise-deterministic Markov processes: A general class of non-diffusion models, *J. Royal Stat.*, **B46** (1984), 353–388.
- [6] Davis, M. H. A., Control of piecewise-deterministic processes via discrete-time dynamic programming, *Proc. 3rd Bad Honnef Symposium on Stochastic Differential Systems*, 1985.
- [7] Davis, M. H. A., Dempster, M. A. H., Sethi, S. P. and Vermes, D., Optimal capacity expansion under uncertainty, *Advances in Applied Probability*, **19** (1987), 156–176.
- [8] Gatarek, D., Impulsive control of a decreasing piecewise-deterministic process with long run average cost, *System Research Institute, Preprint 153*, Warsaw 1987.
- [9] Gugerli, U. S., Optimal stopping of a piecewise-deterministic Markov process, *Stochastics*, **19** (1986), 221–236.
- [10] Hanouzet, B. and Joly, J. L., Convergence uniforme des itérés définissant la solution d'une inéquation quasi-variationnelle abstraite, *C. R. Acad. Sci. Ser. A., Paris*, **286** (1978), 735–738.
- [11] Hordijk, A. and van Duyn Schouten, F. A., Average optimal policies in Markov decision drift processes with applications to a queueing and replacement model, *Advances in Applied Probability*, **15** (1983), 274–303.
- [12] Ishii, H., Perron's method for Hamilton-Jacobi equations, *Duke Math. J.*, **55** (1987), 369–384.
- [13] Ishii, H. and Koike, S., Viscosity solutions of a system of nonlinear second-order elliptic PDEs arising in switching game, *Funkcial. Ekvac.*, **34** (1991), 143–155.
- [14] Lenhart, S. M., Viscosity solutions associated with switching control problems for piecewise-deterministic processes, *Houston J. Math.*, **13** (1987), 405–426.
- [15] Lenhart, S. M. and Liao, Y. C., Integro-differential equations associated with optimal stopping time of a piecewise-deterministic process, *Stochastics*, **15** (1985), 183–207.

- [16] Lenhart, S. M. and Liao, Y. C., Switching control of piecewise-deterministic processes, *J. Optimization*, **59** (1988), 99–115.
- [17] Oleinik, A. O. and Radkevic, E. V., *Second Order Equations with Nonnegative Characteristic Form*, Amer. Math. Soc., Providence, Rhode Island and Plenum Press, New York, 1973.
- [18] Pliska, S. R., On a functional differential equations that arises in Markov control problem, *J. Differential Equations*, **28**, (1978), 390–405.
- [19] Pragarauskas, H., On first order singular Bellman equations, *Lecture Notes in Control and Information Sci.*, **36**, (1981), 175–188.
- [20] Pragarauskas, H., On the control of jump processes, *Lecture Notes in Control and Information Sci.*, **43**, (1982), 338–344.
- [21] Rosberg, Z., Variaga, P. and Walrand, J., Optimal control of service in tandem queues, *IEEE Trans. Auto. Control.*, **AC-27** (1982), 600–610.
- [22] Soner, H. M., Optimal control with state-space constraint II, *SIAM J. Control Optimization*, **24** (1986), 1110–1122.
- [23] Vermes, D., Optimal control of piecewise-deterministic Markov processes, *Stochastics*, **14** (1985), 165–208.

nuna adreso:

Suzanne M. Lenhart
University of Tennessee
Mathematics Department
Knoxville, TN 37996-1300
U.S.A.

Naoki Yamada
Department of Applied Mathematics
Faculty of Engineering
Miyazaki University
Kibana, Miyazaki 889-21
Japan

(Ricevita la 7-an de marto, 1990)