



Nontraumatic True Aneurysm of the Superficial Palmar Arch : A Case Report

Nagura, Issei ; Kokubu, Takeshi ; Fujioka, Hiroyuki ; Mifune, Yutaka ; Nishida, Kotaro ; Akisue, Toshihiro ; Kuroda, Ryosuke ; Kurosawa, ...

(Citation)

The Kobe journal of the medical sciences, 58(1):29-32

(Issue Date)

2012

(Resource Type)

departmental bulletin paper

(Version)

Version of Record

(JaLCD0I)

<https://doi.org/10.24546/81003961>

(URL)

<https://hdl.handle.net/20.500.14094/81003961>



Nontraumatic True Aneurysm of the Superficial Palmar Arch: A Case Report

ISSEI NAGURA^{*1}, TAKESHI KOKUBU², HIROYUKI FUJIOKA³,
YUTAKA MIFUNE², KOTARO NISHIDA², TOSHIHIRO AKISUE²,
RYOSUKE KURODA², and MASAHIRO KUROSAKA²

¹Department of Orthopaedic Surgery, Kobe Rosai Hospital,
4-1-23 Kagoike-dori, Chuo-ku, Kobe, 651-0053, Japan

²Department of Orthopaedic Surgery, Kobe University Graduate School of Medicine,
7-5-1 Kusunoki-cho, Chuo-ku, Kobe, 650-0017, Japan

³School of Rehabilitation, Hyogo University of Health Sciences,
1-3-6 Minatojima, Chuo-ku, Kobe, 650-8530, Japan

Received 9 March 2012/ Accepted 26 March 2012

Key Words: True aneurysm, superficial palmar arch, resection, Magnetic Resonance Angiography

ABSTRACT

A case of a 62 year-old woman with true aneurysm of the superficial palmar arch with no previous history of trauma is described. The aneurysm was resected without reconstruction based on intraoperative evidence of adequate finger perfusion after clumping the distal and proximal ends. Three years after the operation, there has been no recurrence of the aneurysm and the patient has neither complaints nor symptoms.

INTRODUCTION

True aneurysms in the hand are rare, and usually arise from the ulnar artery at the hypothenar eminence.³ In the hand, true aneurysms occurring at the superficial palmar arch have been rarely reported.¹⁻⁴ We present a case of true aneurysm of the superficial palmar arch with no previous history of trauma.

CLINICAL CASE

A 62 year-old woman presented with a mass on the mid-palmar area of the right hand, accompanied by discomfort for three years. No previous history of trauma was presented. The patient had no cold intolerance and no sensory disturbance. Range of motion of the fingers were full, however grip was limited due to the presence of the mass. Clinical examination showed a one cm palpable tumor with pulsation, however Allen's test was negative. A Magnetic Resonance Angiography (MRA) showed an aneurysm of the superficial palmar arch (Fig.1, 2).




Fig. 1. Magnetic Resonance Image of the hand

a. T1 weighted image b. T2 weighted image c. Gadolinium enhanced image



Fig. 2. Magnetic Resonance Angiography of the hand

MRA shows an aneurysm of the superficial palmar arch of the right hand.

During surgical exploration, a longitudinal skin incision was made directly over the aneurysm. Intra-operatively, a large fusiform aneurysm of the superficial palmar arch involving the common digital artery of the middle and ring finger was revealed (Fig.3). Satisfactory perfusion of the fingers was observed after clamping the both ends. After resection, the symptoms disappeared.

NONTRAUMATIC TRUE ANEURYSM



Fig. 3. Intra-operative view of the superficial palmar arch aneurysm

The histology demonstrated a true aneurysm where all three layers of the arterial wall were seen (Fig.4). The patient's postoperative course was uneventful on follow up three years after surgery.




Fig. 4. Microscopic findings

Light microscopy shows three layers in the vessel wall with hematoma consistent with a true aneurysm. No evidence of vasculitis was observed.

DISCUSSION

True aneurysms in the hand are rare and usually result from blunt trauma. A true aneurysm occurring at the superficial palmar arch which is not secondary to blunt trauma, as in our patient, is also rare. The usual treatment, when adequate circulation is observed, is to resect the aneurysm and ligate the arterial ends.^{1,3,4} If ischemia is suspected, the vascular reconstruction is performed.^{2,7} In our patient, preoperatively the patient had no cold

intolerance and there was intraoperative evidence of adequate finger perfusion, therefore we did not reconstruct the artery and, instead, opted to ligate the aneurysm.

Preoperative arteriography is the procedure of choice, because it gives essential information about the localization and the size of the aneurysm. Doppler ultrasound, digital plethymography and pulse oxymetry can also provide important information about the adequacy of collateral circulation. Krinsky et al reported that MRA with gadolinium provides excellent image quality not available with conventional angiography.⁵ Price et al reported that in-plane saturation is problematic in tortuous arteries and arteries with slow flow on MRA without gadolinium.⁶ However, it is possible to image arteries with fundamentally different contrast mechanism that does not depend upon blood flow in MRA with gadolinium. Gadolinium contrast agents improves the quality of MRA.⁶ Therefore, preoperative evaluation of MRA with gadolinium was also useful for preoperative planning.

In conclusion, true aneurysms of the superficial palmar arch in the hand, not secondary to blunt trauma, are rare. If cold intolerance is a presenting symptom, vascular reconstruction play an important role in relieving the symptoms. MRA with gadolinium is useful for preoperative planning for the treatment of the aneurysms.

ACKNOWLEDGEMENTS

The author thanks to Ms. Janina Tubby for English rewriting.

REFERENCES

1. **Aulicino PL, Hutton PMJ. and Du Puy TE.** True aneurysm – a case report and literature review. *J Hand Surg A* 1982;613-616.
2. **Estrella EP. and Lee EY.** Aneurysm of the superficial palmar arch: A Case Report. *J Hand Surg* 2008;**13**:179-182.
3. **Green DP.** True and false traumatic aneurysms in the hand; report of two cases and review of the literature. *J Bone Joint Surg (Am)*1973; **55**:120-128.
4. **Iossifidis AS. and Sutaria PD.** Aneurysm of the superficial palmar arch. A case report. *Int Orthop* 1995; **19**:403-404.
5. **Krinsky G. and Rofsky NM.** MR angiography of the aortic vessels and upper extremities. *Magn Reson Imaging Clin N Am* 1998;**6**:269-92.
6. **Prince MR.** Body MR angiography with gadolinium contrast agents. *Magn Reson Imaging Clin N Am* 1996;**4**:11-24.
7. **Rieck B, Mailander P, Kuske M, Machens HG. and Berger A.** True aneurysms of the palmar arch of the hand: a report of two cases. *Microsurgery* 1996;**17**:102-105.