

PDF issue: 2025-02-21

# On multidimensional inverse scattering for Stark Hamiltonians

Adachi, Tadayoshi Maehara, Katsuhiro

<mark>(Citation)</mark> Journal of Mathematical Physics,48(4):42101-42101

(Issue Date) 2007-04

(Resource Type) journal article

(Version) Version of Record

(URL) https://hdl.handle.net/20.500.14094/90000195



# On multidimensional inverse scattering for Stark Hamiltonians

# Tadayoshi Adachi<sup>a)</sup> and Katsuhiro Maehara

Department of Mathematics, Faculty of Science, Kobe University, 1-1 Rokkodai-cho, Nada-ku, Kobe-shi, Hyogo 657-8501, Japan and Graduate School of Science and Technology, Kobe University, 1-1 Rokkodai-cho, Nada-ku, Kobe-shi, Hyogo 657-8501, Japan

(Received 26 December 2006; accepted 5 February 2007; published online 6 April 2007)

Based on the Enss-Weder ["The geometrical approach to multidimensional inverse scattering," J. Math. Phys. **36**, 3902–3921 (1995)] time-dependent method, we study one of multidimensional inverse scattering problems for Stark Hamiltonians. We first show that when the space dimension is greater than or equal to 2, the high velocity limit of the scattering operator determines uniquely the potential such as  $|x|^{-\gamma}$  with  $\gamma > 1/2$  which is short range under the Stark effect. This is an improvement of previous results obtained by Nicoleau ["Inverse scattering for Stark Hamiltonians with short-range potentials," Asymptotic Anal. **35**, 349–359 (2003)] and Weder ["Multidimensional inverse scattering in an electric field," J. Funct. Anal. **139**, 441–465 (1996)]. Moreover, we prove that for a given long-range part of the potential under the Stark effect, the high velocity limit of the Dollard-type modified scattering operator determines uniquely the short-range part of the potential. © 2007 American Institute of Physics. [DOI: 10.1063/1.2713077]

#### I. INTRODUCTION

In this paper, we consider one of the inverse scattering problems for quantum systems in a constant electric field  $\mathcal{E} \in \mathbb{R}^n$ . Throughout this paper, we assume that  $n \ge 2$ . For brevity's sake, we suppose that  $\mathcal{E}=e_1=(1,0,\ldots,0)$ . The free Stark Hamiltonian under consideration is given by

$$H_0 = -\frac{1}{2}\Delta - x_1,$$
 (1.1)

acting on  $L^2(\mathbb{R}^n)$ , where  $x_1$  is the first component of  $x = (x_1, ..., x_n) \in \mathbb{R}^n$ . It is well known that  $H_0$  is essentially self-adjoint on  $S(\mathbb{R}^n)$ . The self-adjoint realization of  $H_0$  is also denoted by  $H_0$ . We here assume that the potential V is the multiplication operator by V(x), and that V(x) is represented as a sum of parts of very short range, short range, and long range under the Stark effect:

$$V(x) = V^{\rm vs}(x) + V^{\rm s}(x) + V^{\rm l}(x), \qquad (1.2)$$

where  $V^{vs} \in \mathcal{V}^{vs}$ ,  $V^s \in \mathcal{V}^s$ , and  $V^l \in \mathcal{V}^l$ . We here assume that  $\mathcal{V}^{vs}$  is the class of real-valued potentials  $V^{vs}(x)$  satisfying that  $V^{vs}(x) = V_1^{vs}(x) + V_2^{vs}(x)$  with  $V_2^{vs}(x)$  bounded,  $V_1^{vs}(x) - \Delta/2$ -bounded with relative bound less than 1, and  $V_1^{vs}(x)x_1 - \Delta/2$ -bounded, and that

$$\int_{0}^{\infty} \|V^{\rm vs}(x)(-\Delta+1)^{-1}F(|x|\ge R)\|_{\mathcal{B}(L^2)}\mathrm{d}R < \infty,$$
(1.3)

where  $F(|x| \ge R)$  is the characteristic function of  $\{x \in \mathbb{R}^n | |x| \ge R\}$ . Since  $V^{vs}(x)$  is a multiplication operator, the condition (1.3) is equivalent to

0022-2488/2007/48(4)/042101/12/\$23.00

**48**, 042101-1

© 2007 American Institute of Physics

a)Electronic mail: adachi@math.kobe-u.ac.jp

042101-2 T. Adachi and K. Maehara

J. Math. Phys. 48, 042101 (2007)

$$\int_{0}^{\infty} \|F(|x| \ge R) V^{\rm vs}(x) (-\Delta + 1)^{-1}\|_{\mathcal{B}(L^2)} \mathrm{d}R < \infty, \tag{1.4}$$

as is well known (see, e.g., Ref. 13). Moreover,  $\mathcal{V}^s$  is the class of real-valued potentials  $V^s(x)$  satisfying that  $V^s(x) \in C^1(\mathbb{R}^n)$  and that

$$\left|V^{s}(x)\right| \leq C \langle x \rangle^{-\gamma},\tag{1.5}$$

$$\left|\partial^{\beta}V^{s}(x)\right| \leq C_{\beta}\langle x\rangle^{-1-\alpha}, \quad \left|\beta\right| = 1,$$
(1.6)

with some  $1/2 < \alpha \le \gamma \le 1$ , where  $\langle x \rangle = \sqrt{1 + |x|^2}$ . Finally,  $\mathcal{V}^1$  is the class of real-valued potentials  $V^1(x)$  satisfying that  $V^1(x) \in C^2(\mathbb{R}^n)$  and that

$$|\partial^{\beta} V^{l}(x)| \leq C_{\beta} \langle x \rangle^{-\gamma_{D} - \mu |\beta|}, \quad |\beta| \leq 2,$$

with  $0 < \gamma_D \leq 1/2$  and  $1 - \gamma_D < \mu \leq 1$ .

We first consider the case where  $V^1=0$ . It is known that for  $V \in \mathcal{V}^{vs} + \mathcal{V}^s$ ,  $H=H_0+V$  is selfadjoint, and the wave operators

$$W^{\pm} = \underset{t \to \pm \infty}{\text{s-lim}} e^{itH} e^{-itH_0}$$
(1.7)

exist (as for two-body direct scattering under the Stark effect, see, e.g., Refs. 3, 8, 6, 10, 12, 14, 15, 17, and 18; as for *N*-body direct scattering under the Stark effect, see, e.g., Refs. 2 and 9). Then the scattering operator S=S(V) is defined by

$$S = (W^{+})^{*}W^{-}.$$
 (1.8)

Then one of the main results of this paper is represented as follows.

**Theorem 1.1:** Let  $V_1, V_2 \in \mathcal{V}^{vs} + \mathcal{V}^s$ . If  $S(V_1) = S(V_2)$ , then  $V_1 = V_2$ .

It is well known that  $V^{vs}$  is short range and  $V^s$  is long range in the absence of the external electric field, and that both  $V^{vs}$  and  $V^s$  are short range in the presence of a constant electric field. Thus this theorem implies that a certain potential that may be long range in the absence of the electric field can be determined by the scattering operator *S* under the Stark effect. This theorem was first proven by Weder<sup>16</sup> under the conditions  $\gamma > 3/4$  and  $n \ge 2$ . Later Nicoleau<sup>11</sup> proved this theorem for real-valued  $V \in C^{\infty}(\mathbb{R}^n)$  satisfying

$$\left|\partial^{\beta} V(x)\right| \leq C_{\beta} \langle x \rangle^{-\gamma - |\beta|},$$

with some  $\gamma > 1/2$ , under the condition  $n \ge 3$ . Our result improves their result. The key for improving the result of Weder<sup>16</sup> in Ref. 11 is to introduce the Dollard-type modifier  $e^{-i\int_0^t V^8(p_{\perp}s+e_1s^{2/2})ds}$  due to White,<sup>17</sup> where  $p=-i\nabla = (p_1, p_{\perp})$ . The assumption  $n-1 \ge 2$  is needed for his method. We will here use the Graf-type (or Zorbas-type) modifier  $e^{-i\int_0^t V^8(vs+e_1s^{2/2})ds}$  (see Refs. 6 and 19) with a certain constant vector  $v \in \mathbb{R}^n$  instead of  $e^{-i\int_0^t V^8(p_{\perp}s+e_1s^{2/2})ds}$  in order to deal with the case where  $n \ge 2$  and to relax the smoothness condition on potentials supposed by Nicoleau.<sup>11</sup> Our proof as well as theirs are based on the Enss-Weder time-dependent method.<sup>5</sup>

We next consider the case where  $V^{l} \neq 0$ . It is known that for  $V \in \mathcal{V}^{vs} + \mathcal{V}^{s} + \mathcal{V}^{l}$ ,  $H = H_0 + V$  is self-adjoint, and the Dollard-type modified wave operators due to Jensen-Yajima<sup>10</sup>

$$W_D^{\pm} = \operatorname{s-lim}_{t \to \pm \infty} e^{itH_0} M_D(t), \quad M_D(t) = e^{-i\int_0^t v^l (ps + e_1 s^2/2) ds}$$
(1.9)

as well as the Graf-type modified wave operators

042101-3 Inverse scattering for Stark Hamiltonians

J. Math. Phys. 48, 042101 (2007)

$$W_G^{\pm} = \underset{t \to \pm \infty}{\text{s-lim}} e^{itH} e^{-itH_0} M_G(t), \quad M_G(t) = e^{-i\int_0^t V^J(e_1 s^{2/2}) \mathrm{d}s},$$
 (1.10)

and the Dollard-type modified wave operators due to White<sup>17</sup>

$$W_W^{\pm} = \underset{t \to \pm \infty}{\text{s-lim}} e^{itH} e^{-itH_0} M_W(t), \quad M_W(t) = e^{-i\int_0^t V^{\rm d}(p_{\perp}s + e_1s^2/2) \mathrm{d}s}$$
(1.11)

exist by virtue of the condition  $\gamma_D + \mu > 1$  (see Refs. 1 and 17). Then the Dollard-type modified scattering operator  $S_D = S_D(V^1; V^{vs} + V^s)$  is defined by

$$S_D = (W_D^+)^* W_D^-. \tag{1.12}$$

The reason why  $W_D^{\pm}$  are used for defining the modified scattering operator in place of  $W_G^{\pm}$  and  $W_W^{\pm}$  is that  $W_D^{\pm}$  match the Enss-Weder time-dependent method as seen below. Then we obtain the following new result.

**Theorem 1.2:** Let  $V^{l} \in \mathcal{V}^{l}$  be given. Let  $V_{1}, V_{2} \in \mathcal{V}^{vs} + \mathcal{V}^{s}$ . If  $S_{D}(V^{l}; V_{1}) = S_{D}(V^{l}; V_{2})$ , then  $V_{1} = V_{2}$ . Moreover, any one of the Dollard-type modified scattering operators  $S_{D}$  determines uniquely the total potential V(x).

The plan of this paper is as follows: In Sec. II, we consider the case where  $V^1=0$ . In Sec. III, we consider the general case. The main purpose of these two sections is deriving the reconstruction formula (see Theorems 2.1 and 3.1) for potentials in the class  $\mathcal{V}^{vs} + \mathcal{V}^s + \mathcal{V}^s$ , which is fairly suitable for studying the scattering problems under the Stark effect. We note that Theorems 2.1 and 3.1 imply that the high velocity limit of the (modified) scattering operator determines uniquely the short-range part of the potential.

#### **II. SHORT-RANGE CASE**

In this section, we consider the case where  $V^{l}=0$ . The main purpose of this section is showing the following reconstruction formula, which yields the proof of Theorem 1.1.

**Theorem 2.1:** Let  $\hat{v} \in \mathbb{R}^n$  be given such that  $|\hat{v}|=1$  and  $|\hat{v} \cdot e_1| < 1$ . Set  $v = |v|\hat{v}$ . Let  $\eta > 0$  be given, and  $\Phi_0$ ,  $\Psi_0 \in L^2(\mathbb{R}^n)$  be such that  $\hat{\Phi}_0$ ,  $\hat{\Psi}_0 \in C_0^{\infty}(\mathbb{R}^n)$  with supp  $\hat{\Phi}_0$ , supp  $\hat{\Psi}_0 \subset \{\xi \in \mathbb{R}^n | |\xi| < \eta\}$ . Here  $\hat{\Phi}_0$  and  $\hat{\Psi}_0$  stand for the Fourier transforms of  $\Phi_0$  and  $\Psi_0$ , respectively. Set  $\Phi_v = e^{ivx}\Phi_0$  and  $\Psi_v = e^{ivx}\Psi_0$ . Then

$$\begin{split} \lim_{|v| \to \infty} & |v| (i[S, p_j] \Phi_v, \Psi_v) = \int_{-\infty}^{\infty} \left[ (V^{vs}(x + \hat{v}\tau) p_j \Phi_0, \Psi_0) - (V^{vs}(x + \hat{v}\tau) \Phi_0, p_j \Psi_0) \right. \\ & + i ((\partial_j V^s) (x + \hat{v}\tau) \Phi_0, \Psi_0) \right] \mathrm{d}\tau \end{split}$$
(2.1)

holds. Here  $p_i = -i\partial_i$ .

We will make preparations for the proof of Theorem 2.1. Throughout this paper, we need the following proposition due to  $\text{Enss}^4$  (see Proposition 2.10 in Ref. 4).

**Proposition 2.1:** For any  $f \in C_0^{\infty}(\mathbb{R}^n)$  with supp  $f \subset \{x \in \mathbb{R}^n | |x| < \eta\}$  for some  $\eta > 0$ , and any  $l \in \mathbb{N}$ , there exists a constant  $C_l$  dependent on f only such that

$$\left\|F(x \in \mathcal{M}')e^{it\Delta/2}f(p-v)F(x \in \mathcal{M})\right\| \le C_l(1+r+|t|)^{-l}$$

$$(2.2)$$

for  $v \in \mathbb{R}^n$ ,  $t \in \mathbb{R}$ , and measurable sets  $\mathcal{M}$ ,  $\mathcal{M}'$  with the property that  $r = \operatorname{dist}(\mathcal{M}', \mathcal{M} + vt) - \eta |t| \ge 0$ . Here  $F(x \in \mathcal{M})$  stands for the characteristic function of  $\mathcal{M}$ .

The following lemma has been proven by Weder.<sup>16</sup>

**Lemma 2.1:** Let v and  $\Phi_v$  be as in Theorem 2.1. Then

$$\int_{-\infty}^{\infty} \|V^{vs}(x)e^{-itH_0}\Phi_v\|dt = O(|v|^{-1})$$
(2.3)

holds as  $|v| \rightarrow \infty$  for  $V^{vs} \in \mathcal{V}^{vs}$ .

042101-4 T. Adachi and K. Maehara

J. Math. Phys. 48, 042101 (2007)

Proof: We will sketch the proof for consistency. By virtue of the Avron-Herbst formula<sup>3</sup>

$$e^{-itH_0} = e^{-it^3/6} e^{itx_1} e^{-ip_1 t^2/2} e^{it\Delta/2}$$
(2.4)

and

$$e^{-ivx}e^{it\Delta/2}e^{ivx} = e^{-iv^2t/2}e^{-ipvt}e^{it\Delta/2},$$
 (2.5)

one has

$$\|V^{\rm vs}(x){\rm e}^{-itH_0}\Phi_v\| = \|V^{\rm vs}(x+vt+e_1t^2/2){\rm e}^{it\Delta/2}\Phi_0\|.$$

Take  $f \in C_0^{\infty}(\mathbb{R}^n)$  such that  $f\hat{\Phi}_0 = \hat{\Phi}_0$  and  $\operatorname{supp} f \subset \{\xi \in \mathbb{R}^n | |\xi| < \eta\}$ . Then one has

$$\|V^{vs}(x+vt+e_1t^2/2)e^{it\Delta/2}\Phi_0\| = \|V^{vs}(x+vt+e_1t^2/2)(-\Delta+1)^{-1}e^{it\Delta/2}f(p)(-\Delta+1)\Phi_0\| \le I_1+I_2+I_3,$$
 with

$$\begin{split} I_1 &= \|V^{\rm vs}(x+vt+e_1t^2/2)(-\Delta+1)^{-1}\|_{\mathcal{B}(L^2)}\|F(|x| \ge 3\lambda|v||t|){\rm e}^{it\Delta/2}f(p)F(|x| \le \lambda|v||t|)\|_{\mathcal{B}(L^2)} \\ &\times \|(-\Delta+1)\Phi_0\|, \end{split}$$

$$\begin{split} I_2 &= \|V^{vs}(x+vt+e_1t^2/2)(-\Delta+1)^{-1}\|_{\mathcal{B}(L^2)}\|F(|x| \ge 3\lambda|v||t|)\mathrm{e}^{it\Delta/2}f(p)F(|x| > \lambda|v||t|)\langle x\rangle^{-2}\|_{\mathcal{B}(L^2)} \\ &\times \|\langle x\rangle^2(-\Delta+1)\Phi_0\|, \end{split}$$

$$I_{3} = \|V^{vs}(x + vt + e_{1}t^{2}/2)(-\Delta + 1)^{-1}F(|x| < 3\lambda|v||t|)\|_{\mathcal{B}(L^{2})}\|e^{it\Delta/2}f(p)(-\Delta + 1)\Phi_{0}\|_{\mathcal{B}(L^{2})}\|e^{it\Delta/2}f(p)(-\Delta + 1)\Phi_{0}\|_$$

where  $\lambda > 0$ , which is independent of |v|, will be determined below. Since

$$\|V^{\rm vs}(x+vt+e_1t^2/2)(-\Delta+1)^{-1}\|_{\mathcal{B}(L^2)} = \|V^{\rm vs}(x)(-\Delta+1)^{-1}\|_{\mathcal{B}(L^2)},$$

we have

$$I_1 + I_2 \le C(1 + |v||t|)^{-2},$$

for  $\lambda |v| \ge \eta$ . Here we used Proposition 2.1 for estimating  $I_1$  under the condition  $\lambda |v| \ge \eta$ . As for  $I_3$ , we note that

$$\begin{split} \|V^{vs}(x+vt+e_1t^2/2)(-\Delta+1)^{-1}F(|x|<3\lambda|v||t|)\|_{\mathcal{B}(L^2)} \\ &=\|V^{vs}(x)(-\Delta+1)^{-1}F(|x-vt-e_1t^2/2|<3\lambda|v||t|)\|_{\mathcal{B}(L^2)}. \end{split}$$

We put  $\delta = |\hat{v}e_1| < 1$ . Then we have

$$\begin{aligned} |vt + e_1 t^2 / 2|^2 &= |v|^2 t^2 + t^4 / 4 + t^3 v e_1 \ge |v|^2 |t|^2 + |t|^4 / 4 - \delta |v| |t|^3 \\ &= |t|^2 (|t|^2 - 4\delta |v||t| + 4\delta^2 |v|^2) / 4 + (1 - \delta^2) |v|^2 |t|^2 \ge (1 - \delta^2) |v|^2 |t|^2. \end{aligned}$$
(2.6)

If one takes  $\lambda$  as  $4\lambda = (1 - \delta^2)^{1/2} > 0$ , one has

$$F(|x - vt - e_1t^2/2| < 3\lambda |v||t|) = F(|x - vt - e_1t^2/2| < 3\lambda |v||t|)F(|x| \ge \lambda |v||t|).$$

Then we have

$$I_{3} \leq \|V^{\rm vs}(x)(-\Delta+1)^{-1}F(|x| \geq \lambda \|v\||t|)\|_{\mathcal{B}(L^{2})}\|(-\Delta+1)\Phi_{0}\|$$

Therefore we obtain

042101-5 Inverse scattering for Stark Hamiltonians

$$\int_{-\infty}^{\infty} (I_1 + I_2 + I_3) dt = O(|v|^{-1})$$

by assumption, which implies the lemma.

The following lemma is the key in this paper.

**Lemma 2.2:** Let v and  $\Phi_v$  be as in Theorem 2.1. Then

$$\int_{-\infty}^{\infty} \| (V^{s}(x) - V^{s}(vt + e_{1}t^{2}/2))e^{-itH_{0}}\Phi_{v} \| dt = \begin{cases} O(|v|^{-\alpha}) & \text{if } \alpha < 1\\ O(|v|^{-(1-\epsilon)}) & \text{if } \alpha = 1 \end{cases}$$
(2.7)

holds with any  $0 \le \epsilon \le 1$  as  $|v| \to \infty$  for  $V^{s} \in \mathcal{V}^{s}$ .

**Proof:** As in the proof of Lemma 2.1, take  $f \in C_0^{\infty}(\mathbb{R}^n)$  such that  $f\hat{\Phi}_0 = \hat{\Phi}_0$  and  $\operatorname{supp} f \subset \{\xi \in \mathbb{R}^n | |\xi| < \eta\}$ . Then one has

$$\left\| (V^{s}(x) - V^{s}(vt + e_{1}t^{2}/2))e^{-itH_{0}}\Phi_{v} \right\| = \left\| (V^{s}(x + vt + e_{1}t^{2}/2) - V^{s}(vt + e_{1}t^{2}/2))e^{it\Delta/2}f(p)\Phi_{0} \right\|$$

by virtue of Eqs. (2.4) and (2.5). This can be estimated as

$$||(V^{s}(x) - V^{s}(vt + e_{1}t^{2}/2))e^{-itH_{0}}\Phi_{v}|| \leq I_{1} + I_{2} + I_{3},$$

with

$$I_{1} = 2 \sup_{y \in \mathbb{R}^{n}} |V^{s}(y)| \|F(|x| \ge 3|v|^{\rho}|t|) e^{it\Delta/2} f(p)F(|x| \le |v|^{\rho}|t|) \|_{\mathcal{B}(L^{2})} \|\Phi_{0}\|,$$
  
$$I_{2} = 2 \sup_{y \in \mathbb{R}^{n}} |V^{s}(y)| \|F(|x| \ge 3|v|^{\rho}|t|) e^{it\Delta/2} f(p)F(|x| > |v|^{\rho}|t|) \langle x \rangle^{-2} \|_{\mathcal{B}(L^{2})} \|\langle x \rangle^{2} \Phi_{0}\|$$

$$I_3 = \|(V^{s}(x+vt+e_1t^2/2) - V^{s}(vt+e_1t^2/2))F(|x| < 3|v|^{\rho}|t|)\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f(p)\Phi_0\|_{\mathcal{B}(L^2)}\|e^{it\Delta/2}f$$

where  $0 < \rho < 1$ , which is independent of |v|, will be determined below. By using Proposition 2.1 for estimating  $I_1$  under the condition  $|v|^{\rho} \ge \eta$ , we have

$$I_1 + I_2 \le C(1 + |v|^{\rho}|t|)^{-2},$$

for  $|v|^{\rho} \ge \eta$ , which implies

$$\int_{-\infty}^{\infty} (I_1 + I_2) dt = O(|v|^{-\rho}).$$
(2.8)

We set  $\delta = |\hat{v} \cdot e_1| < 1$ . If  $|x| < 3|v|^{\rho}|t|$  and  $3|v|^{\rho-1} \le (1-\delta)/4$ ,

$$\begin{split} |x + vt + e_1 t^2 / 2|^2 &= |x + vt|^2 + t^4 / 4 + t^2 (x + vt) e_1 > (1 - 3|v|^{\rho - 1})^2 |v|^2 |t|^2 + |t|^4 / 4 - (\delta + 3|v|^{\rho - 1}) |v|| t|^3 \\ & \ge ((3 + \delta) / 4)^2 |v|^2 |t|^2 + |t|^4 / 4 - (1 + 3\delta) |v|| t|^3 / 4. \end{split}$$

Since  $|t|^4/4 - (1+3\delta)|v||t|^3/4 + ((1+3\delta)/4)^2|v|^2|t|^2 \ge 0$ , we have

$$|x + vt + e_1 t^2/2|^2 \ge \{((3 + \delta)/4)^2 - ((1 + 3\delta)/4)^2\}|v|^2|t|^2 = (1 - \delta^2)|v|^2|t|^2/2.$$

We here set  $c_1 = ((1 - \delta^2)/2)^{1/2}$ . Moreover, since  $((3 + \delta)/4)^2 |v|^2 |t|^2 - (1 + 3\delta) |v||t|^3/4 + ((1 + 3\delta)/(3 + \delta))^2 |t|^4/4 \ge 0$ , we have

$$|x + vt + e_1 t^2/2|^2 \ge \{1 - ((1 + 3\delta)/(3 + \delta))^2\}|t|^4/4 = 2(1 - \delta^2)|t|^4/(3 + \delta)^2.$$

We here set  $c_2 = (2(1-\delta^2))^{1/2}/(3+\delta)$ . Thus we obtain

042101-6 T. Adachi and K. Maehara

J. Math. Phys. 48, 042101 (2007)

$$|x + vt + e_1 t^2/2| \ge \max\{c_1 |v| |t|, c_2 |t|^2\},$$
(2.9)

if  $|x| < 3|v|^{\rho}|t|$  and  $3|v|^{\rho-1} \le (1-\delta)/4$ . Now we introduce  $V_{v,t}(x)$  as

$$V_{v,t}(x) = V^{s}(x)g(x/(|v||t|))$$

where  $g \in C^{\infty}(\mathbf{R}^n)$  such that

$$g(x) = \begin{cases} 1 & \text{if } |x| \ge c_1 \\ 0 & \text{if } |x| \le c_1/2. \end{cases}$$

Then  $I_3$  is estimated as

$$I_{3} \leq \|(V_{v,t}(x+vt+e_{1}t^{2}/2)-V_{v,t}(vt+e_{1}t^{2}/2))F(|x| < 3|v|^{\rho}|t|)\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L^{2})}\|\Phi_{0}\|_{\mathcal{B}(L$$

if  $3|v|^{\rho-1} \le (1-\delta)/4$ . By using

$$V_{v,t}(x+vt+e_1t^2/2) - V_{v,t}(vt+e_1t^2/2) = \int_0^1 (\nabla V_{v,t})(\theta x+vt+e_1t^2/2)xd\theta,$$

we estimate  $I_3$  as

$$I_{3} \leq 3 \|\Phi_{0}\||v|^{\rho}|t| \int_{0}^{1} \|(\nabla V^{s})(\theta x + vt + e_{1}t^{2}/2)F(|x| < 3|v|^{\rho}|t|)\|_{\mathcal{B}(L^{2})} d\theta + C \|\Phi_{0}\||v|^{\rho-1} \\ \times \int_{0}^{1} \|V^{s}(\theta x + vt + e_{1}t^{2}/2)F(|x| < 3|v|^{\rho}|t|)\|_{\mathcal{B}(L^{2})} d\theta.$$

By virtue of Eq. (2.9), we have

$$\int_{|t| \ge c_1 |v|/c_2} I_3 dt \le C' \left( |v|^{\rho} \int_{|t| \ge c_1 |v|/c_2} |t|^{-1-2\alpha} dt + |v|^{\rho-1} \int_{|t| \ge c_1 |v|/c_2} |t|^{-2\gamma} dt \right) \le O(|v|^{\rho-2\alpha}) + O(|v|^{\rho-2\gamma})$$
$$= O(|v|^{\rho-2\alpha})$$

because  $|\theta x| \leq |x|$  for  $0 \leq \theta \leq 1$ . We here used that  $1/2 < \alpha \leq \gamma$ . On the other hand,

$$\begin{split} \int_{|t| < c_1 |v|/c_2} I_3 \mathrm{d}t &\leq C |v|^{\rho} \int_{|t| < c_1 |v|/c_2} |t| (1+|v||t|)^{-1-\alpha} \mathrm{d}t + C |v|^{\rho-1} \int_{|t| < c_1 |v|/c_2} (1+|v||t|)^{-\gamma} \mathrm{d}t \\ &\leq 2C |v|^{\rho-1} \int_{|t| < c_1 |v|/c_2} (1+|v||t|)^{-\alpha} \mathrm{d}t \leq 2C |v|^{\rho-2} \int_{|\tau| < c_1 |v|^2/c_2} (1+|\tau|)^{-\alpha} \mathrm{d}\tau \\ &\leq \begin{cases} O(|v|^{\rho-2\alpha}) & \text{if } \alpha < 1 \\ O(|v|^{\rho-2} \log |v|) & \text{if } \alpha = 1 \end{cases} \end{split}$$

is obtained. Therefore we have

$$\int_{-\infty}^{\infty} I_3 \mathrm{d}t \leq \begin{cases} O(|v|^{\rho-2\alpha}) & \text{if } \alpha < 1\\ O(|v|^{\rho-2}\log|v|) & \text{if } \alpha = 1. \end{cases}$$
(2.10)

Noticing Eqs. (2.8) and (2.10), and taking  $\rho$  as

$$\rho = \begin{cases} \alpha & \text{if } \alpha < 1 \\ 1 - \epsilon & \text{if } \alpha = 1, \end{cases}$$

with  $0 \le \epsilon \le 1$ , we obtain the lemma.

### 042101-7 Inverse scattering for Stark Hamiltonians

#### J. Math. Phys. 48, 042101 (2007)

We now introduce auxiliary wave operators,

$$\Omega_{G,v}^{\pm} = \underset{t \to \pm \infty}{\text{s-lim}} e^{itH} U_{G,v}(t), \quad U_{G,v}(t) = e^{-itH_0} e^{-iJ_0^{L} V^{\text{s}}(vs + e_1 s^2/2) \text{d}s}.$$

We here emphasize that the Graf-type modifier  $e^{-i f_0^\ell V^8(vs+e_1s^2/2)ds}$  commutes with any operators. This fact will be used frequently. Then we have the following.

**Lemma 2.3:** Let v and  $\Phi_v$  be as in Theorem 2.1. Then

$$\sup_{t \in \mathbf{R}} \| (e^{-itH} \Omega_{G,v}^{-} - U_{G,v}(t)) \Phi_{v} \| = \begin{cases} O(|v|^{-\alpha}) & \text{if } \alpha < 1\\ O(|v|^{-(1-\epsilon)}) & \text{if } \alpha = 1 \end{cases}$$
(2.11)

holds with any  $0 < \epsilon < 1$  as  $|v| \rightarrow \infty$ .

*Proof:* Noting that  $\|(e^{-itH}\Omega_{G,v}^- - U_{G,v}(t))\Phi_v\| = \|(\Omega_{G,v}^- - e^{itH}U_{G,v}(t))\Phi_v\|,$ 

$$\begin{split} \|(\Omega_{G,v}^{-} - e^{itH}U_{G,v}(t))\Phi_{v}\| &\leq \int_{-\infty}^{t} \|V^{vs}(x)U_{G,v}(\tau)\Phi_{v}\|d\tau + \int_{-\infty}^{t} \|(V^{s}(x) - V^{s}(v\tau + e_{1}\tau^{2}/2))U_{G,v}(\tau)\Phi_{v}\|d\tau \\ &\leq \int_{-\infty}^{\infty} \|V^{vs}(x)e^{-i\tau H_{0}}\Phi_{v}\|d\tau + \int_{-\infty}^{\infty} \|(V^{s}(x) - V^{s}(v\tau + e_{1}\tau^{2}/2))e^{-i\tau H_{0}}\Phi_{v}\|d\tau \\ \end{split}$$

yields the lemma by virtue of Lemmas 2.1 and 2.2. We here used the commutativity of  $e^{-i\int_0^t V^{\delta}(vs+e_1s^2/2)ds}$  mentioned above.

*Proof of Theorem 2.1:* Since the proof is quite similar to the one of Theorem 2.4 in Ref. 16, we sketch it.

We first note that S is represented as

$$S = (W^{+})^{*}W^{-} = I_{G,v}(\Omega_{G,v}^{+})^{*}\Omega_{G,v}^{-}, \quad I_{G,v} = e^{-i\int_{-\infty}^{\infty} V^{s}(v\,\theta + e_{1}\,\theta^{2}/2)\,\mathrm{d}\theta}.$$

Since  $[S, p_j] = [S - I_{G,v}, p_j - v_j], (p_j - v_j)\Phi_v = (p_j\Phi_0)_v$ , and

$$i(S - I_{G,v})\Phi_v = I_{G,v}i(\Omega_{G,v}^+ - \Omega_{G,v}^-)^*\Omega_{G,v}^-\Phi_v = I_{G,v}\int_{-\infty}^{\infty} U_{G,v}(t)^*V_t(x)e^{-itH}\Omega_{G,v}^-\Phi_v dt,$$

with  $V_t(x) = V^{vs}(x) + V^{s}(x) - V^{s}(vt + e_1t^2/2)$ , we have

$$|v|(i[S,p_j]\Phi_v,\Psi_v) = I_{G,v}\{I(v) + R(v)\}$$

with

$$I(v) = |v| \int_{-\infty}^{\infty} \left[ (V_t(x)U_{G,v}(t)(p_j\Phi_0)_v, U_{G,v}(t)\Psi_v) - (V_t(x)U_{G,v}(t)\Phi_v, U_{G,v}(t)(p_j\Psi_0)_v) \right] dt,$$

$$\begin{split} R(v) &= \left| v \right| \int_{-\infty}^{\infty} \left[ ((\mathrm{e}^{-itH} \Omega_{G,v}^{-} - U_{G,v}(t))(p_{j} \Phi_{0})_{v}, V_{t}(x) U_{G,v}(t) \Psi_{v}) \right. \\ & - \left( (\mathrm{e}^{-itH} \Omega_{G,v}^{-} - U_{G,v}(t)) \Phi_{v}, V_{t}(x) U_{G,v}(t)(p_{j} \Psi_{0})_{v} \right) \right] \mathrm{d}t. \end{split}$$

By Lemmas 2.1, 2.2, and 2.3, one has

$$|R(v)| = \begin{cases} O(|v|^{1-2\alpha}) & \text{if } \alpha < 1\\ O(|v|^{-1+2\epsilon}) & \text{if } \alpha = 1. \end{cases}$$

We here used the commutativity of  $e^{-if_0^t V^s(vs+e_1s^2/2) ds}$ . Since  $\alpha > 1/2$  by assumption and one can take  $\varepsilon$  as  $0 < \varepsilon < 1/2$ ,

042101-8 T. Adachi and K. Maehara

$$\lim_{|v|\to\infty} R(v) = 0$$

holds. Using the commutativity of  $e^{-i\int_0^t V^s(vs+e_1s^2/2) ds}$ , the Avron-Herbst formula (2.4) and (2.5), I(v) is rewritten as

$$I(v) = |v| \int_{-\infty}^{\infty} \left[ (V_t(x+vt+e_1t^2/2)e^{it\Delta/2}p_j\Phi_0, e^{it\Delta/2}\Psi_0) - (V_t(x+vt+e_1t^2/2)e^{it\Delta/2}\Phi_0, e^{it\Delta/2}p_j\Psi_0) \right] dt.$$

Since

$$(\{V^{s}(x+vt+e_{1}t^{2}/2) - V^{s}(vt+e_{1}t^{2}/2)\}e^{it\Delta/2}p_{j}\Phi_{0}, e^{it\Delta/2}\Psi_{0}) - (\{V^{s}(x+vt+e_{1}t^{2}/2) - V^{s}(vt+e_{1}t^{2}/2)\}e^{it\Delta/2}\Phi_{0}, e^{it\Delta/2}p_{j}\Psi_{0}) = i((\partial_{j}V^{s})(x+vt+e_{1}t^{2}/2)e^{it\Delta/2}\Phi_{0}, e^{it\Delta/2}\Psi_{0})$$

and  $\partial_i V^s \in \mathcal{V}^{vs}$  by assumption, I(v) is rewritten as

$$\begin{split} I(v) &= |v| \int_{-\infty}^{\infty} \left[ (V^{vs}(x+vt+e_1t^2/2)e^{it\Delta/2}p_j\Phi_0, e^{it\Delta/2}\Psi_0) - (V^{vs}(x+vt+e_1t^2/2)e^{it\Delta/2}\Phi_0, e^{it\Delta/2}p_j\Psi_0) \right. \\ &+ i((\partial_j V^s)(x+vt+e_1t^2/2)e^{it\Delta/2}\Phi_0, e^{it\Delta/2}\Psi_0) \right] dt = \int_{-\infty}^{\infty} l_v(\tau)d\tau \end{split}$$

with

$$\begin{split} l_v(\tau) &= (V^{vs}(x+\hat{v}\,\tau+e_1(\tau/|v|)^{2/2})\mathrm{e}^{i(\pi/|v|)\Delta/2}p_j\Phi_0, \mathrm{e}^{i(\pi/|v|)\Delta/2}\Psi_0) - (V^{vs}(x+\hat{v}\,\tau) \\ &+ e_1(\tau/|v|)^{2/2})\mathrm{e}^{i(\pi/|v|)\Delta/2}\Phi_0, \mathrm{e}^{i(\pi/|v|)\Delta/2}p_j\Psi_0) + i((\partial_j V^s)(x+\hat{v}\,\tau) \\ &+ e_1(\tau/|v|)^{2/2})\mathrm{e}^{i(\pi/|v|)\Delta/2}\Phi_0, \mathrm{e}^{i(\pi/|v|)\Delta/2}\Psi_0). \end{split}$$

Since

$$|l_{v}(\tau)| \leq C\{\|V^{vs}(x)(-\Delta+1)^{-1}F(|x| \geq \lambda|\tau|)\|_{\mathcal{B}(L^{2})} + (1+|\tau|)^{-2} + (1+|\tau|)^{-1-\alpha}\}$$

is obtained as in the proof of Lemma 2.1, we see that

$$\lim_{|v| \to \infty} I(v) = \int_{-\infty}^{\infty} \left[ (V^{vs}(x + \hat{v}\tau)p_j \Phi_0, \Psi_0) - (V^{vs}(x + \hat{v}\tau)\Phi_0, p_j \Psi_0) + i((\partial_j V^s)(x + \hat{v}\tau)\Phi_0, \Psi_0) \right] d\tau$$

by the Lebesgue dominated convergence theorem. Since

$$\begin{split} |vt + e_1 t^2 / 2|^2 &= |v|^2 t^2 + t^4 / 4 + t^3 v e_1 \geqslant |v|^2 |t|^2 + |t|^4 / 4 - \delta |v| |t|^3 = (|v|^2 |t|^2 - \delta |v| |t|^3 + \delta^2 |t|^4 / 4) \\ &+ (1 - \delta^2) |t|^4 / 4 \geqslant (1 - \delta^2) |t|^4 / 4, \end{split}$$

as well as Eq. (2.6) holds, we see that  $\lim_{|v|\to\infty} I_{G,v}=1$  by the Lebesgue dominated convergence theorem because of  $\gamma > 1/2$ . These imply the theorem.

By virtue of Theorem 2.1 and the Plancherel formula associated with the Radon transform (see Ref. 7), Theorem 1.1 can be shown in the same way as in the proof of Theorem 1.2 of Ref. 16 (see also Ref. 5). Thus we omit the proof of Theorem 1.1.

## **III. LONG-RANGE CASE**

The main purpose of this section is showing the following reconstruction formula, which yields the proof of Theorem 1.2.

**Theorem 3.1:** Let  $\hat{v} \in \mathbb{R}^n$  be given such that  $|\hat{v}|=1$  and  $|\hat{v} \cdot e_1| < 1$ . Set  $v = |v|\hat{v}$ . Let  $\eta > 0$  be given, and  $\Phi_0$ ,  $\Psi_0 \in L^2(\mathbb{R}^n)$  be such that  $\hat{\Phi}_0$ ,  $\hat{\Psi}_0 \in C_0^{\infty}(\mathbb{R}^n)$  with supp  $\hat{\Phi}_0$ , supp  $\hat{\Psi}_0 \subset \{\xi \in \mathbb{R}^n | |\xi| < \eta\}$ . Set  $\Phi_v = e^{ivx}\Phi_0$  and  $\Psi_v = e^{ivx}\Psi_0$ . Then

$$\begin{split} \lim_{|v| \to \infty} & |v| (i[S_D, p_j] \Phi_v, \Psi_v) = \int_{-\infty}^{\infty} \left[ (V^{vs}(x + \hat{v}\tau) p_j \Phi_0, \Psi_0) - (V^{vs}(x + \hat{v}\tau) \Phi_0, p_j \Psi_0) \right. \\ & + i ((\partial_j V^s)(x + \hat{v}\tau) \Phi_0, \Psi_0) + i ((\partial_j V^l)(x + \hat{v}\tau) \Phi_0, \Psi_0) \right] \mathrm{d}\tau \tag{3.1}$$

holds.

We first need the following lemma.

**Lemma 3.1:** Let v and  $\Phi_v$  be as in Theorem 3.1. Then

$$\sup_{t \in \mathbf{R}} \|\langle x \rangle^2 M_{D,v}(t) \Phi_0\| = O(1)$$
(3.2)

holds as  $|v| \rightarrow \infty$ , where  $M_{D,v}(t) = e^{-i\int_0^t v^l (ps+vs+e_1s^2/2) ds}$ .

*Proof:* As in the proof of Lemma 2.1, take  $f \in C_0^{\infty}(\mathbb{R}^n)$  such that  $f\hat{\Phi}_0 = \hat{\Phi}_0$  and  $\operatorname{supp} f \subset \{\xi \in \mathbb{R}^n | |\xi| < \eta\}$ . Since  $x = i\partial_p$ , one has

$$\begin{split} \|\|x\|^{2}M_{D,v}(t)f(p)\langle x\rangle^{-2}\|_{\mathcal{B}(L^{2})} &\leq \|M_{D,v}(t)f(p)\|x\|^{2}\langle x\rangle^{-2}\|_{\mathcal{B}(L^{2})} + 2\left\|M_{D,v}(t)\left(\int_{0}^{t}s(\nabla V^{1})(ps+vs+e_{1}s^{2}/2)ds\right)f(p)x\langle x\rangle^{-2}\right\|_{\mathcal{B}(L^{2})} \\ &+ e_{1}s^{2}/2)ds\right)f(p)x\langle x\rangle^{-2}\right\|_{\mathcal{B}(L^{2})} + 2\|M_{D,v}(t)(\nabla f)(p)x\langle x\rangle^{-2}\|_{\mathcal{B}(L^{2})} \\ &+ 2\left\|M_{D,v}(t)\left(\int_{0}^{t}s(\nabla V^{1})(ps+vs+e_{1}s^{2}/2)ds\right)(\nabla f)(p)\langle x\rangle^{-2}\right\|_{\mathcal{B}(L^{2})} \\ &+ \|M_{D,v}(t)(\Delta f)(p)\langle x\rangle^{-2}\|_{\mathcal{B}(L^{2})} + \left\|M_{D,v}(t)\left(\int_{0}^{t}s(\nabla V^{1})(ps+vs+e_{1}s^{2}/2)ds\right)^{2}f(p)\langle x\rangle^{-2}\right\|_{\mathcal{B}(L^{2})} \\ &+ e_{1}s^{2}/2)ds\right)^{2}f(p)\langle x\rangle^{-2}\Big\|_{\mathcal{B}(L^{2})} + \left\|M_{D,v}(t)\left(\int_{0}^{t}s^{2}(\Delta V^{1})(ps+vs+e_{1}s^{2}/2)ds\right)f(p)\langle x\rangle^{-2}\right\|_{\mathcal{B}(L^{2})}. \end{split}$$

Set  $\delta = |\hat{v} \cdot e_1| < 1$ . If  $|\xi| \le \eta$  and  $\eta/|v| \le (1-\delta)/4$ , then

$$\begin{aligned} |\xi t + vt + e_1 t^2 / 2|^2 &= |\xi + v|^2 t^2 + t^4 / 4 + t^3 (\xi + v) e_1 > (1 - \eta / |v|)^2 |v|^2 |t|^2 + |t|^4 / 4 - (\delta + \eta / |v|) |v| |t|^3 \\ &\ge ((3 + \delta) / 4)^2 |v|^2 |t|^2 + |t|^4 / 4 - (1 + 3\delta) |v| |t|^3 / 4. \end{aligned}$$
(3.3)

Since  $((3+\delta)/4)^2 |v|^2 |t|^2 - (1+3\delta) |v||t|^3/4 + ((1+3\delta)/(3+\delta))^2 |t|^4/4 \ge 0$ , we have

$$|\xi t + vt + e_1 t^2/2|^2 \ge 2(1 - \delta^2)|t|^4/(3 + \delta)^2 = (c_2|t|^2)^2.$$
(3.4)

This implies the lemma because  $1-2(\gamma_D+\mu) < -1$  and  $2-2(\gamma_D+2\mu)=2-4(\gamma_D+\mu)+2\gamma_D < -1$  by assumption.

**Lemma 3.2:** Let v and  $\Phi_v$  be as in Theorem 3.1. Then

$$\int_{-\infty}^{\infty} \|V^{\rm vs}(x)U_D(t)\Phi_v\| dt = O(|v|^{-1})$$
(3.5)

holds as  $|v| \rightarrow \infty$  for  $V^{vs} \in \mathcal{V}^{vs}$ .

Proof: By virtue of the Avron-Herbst formula (2.4) and (2.5), one has

042101-10 T. Adachi and K. Maehara

J. Math. Phys. 48, 042101 (2007)

$$\|V^{\rm vs}(x)U_D(t)\Phi_v\| = \|V^{\rm vs}(x+vt+e_1t^2/2){\rm e}^{it\Delta/2}M_{D,v}(t)\Phi_0\|.$$

Then the lemma can be proven in the same way as in the proof of Lemma 2.1, by virtue of Lemma 3.1.  $\hfill \Box$ 

The following lemma can be also proven in the same way as in the proof of Lemma 2.2. Thus we omit the proof.

**Lemma 3.3:** Let v and  $\Phi_v$  be as in Theorem 3.1. Then

$$\int_{-\infty}^{\infty} \left\| (V^{s}(x) - V^{s}(vt + e_{1}t^{2}/2))U_{D}(t)\Phi_{v} \right\| dt = \begin{cases} O(|v|^{-\alpha}) & \text{if } \alpha < 1\\ O(|v|^{-(1-\epsilon)}) & \text{if } \alpha = 1 \end{cases}$$
(3.6)

holds with any  $0 \le \epsilon \le 1$  as  $|v| \to \infty$  for  $V^{s} \in \mathcal{V}^{s}$ .

The following lemma is the key in this section.

**Lemma 3.4:** Let v and  $\Phi_v$  be as in Theorem 3.1. Then

$$\int_{-\infty}^{\infty} \| (V^{l}(x) - V^{l}(pt - e_{1}t^{2}/2))U_{D}(t)\Phi_{v} \| dt = O(|v|^{-\rho})$$
(3.7)

holds with some  $1/2 < \rho < 1$  as  $|v| \to \infty$  for  $V^{l} \in \mathcal{V}^{l}$ .

Proof: Before showing the lemma, we note that

$$e^{-itH_0}V^{l}(pt + e_1t^2/2) = V^{l}(pt - e_1t^2/2)e^{-itH_0}$$
(3.8)

holds by virtue of the Avron-Herbst formula (2.4).

As in the proof of Lemma 2.1, take  $f \in C_0^{\infty}(\mathbb{R}^n)$  such that  $f\hat{\Phi}_0 = \hat{\Phi}_0$  and  $\operatorname{supp} f \subset \{\xi \in \mathbb{R}^n | |\xi| < \eta\}$ . By virtue of the Avron-Herbst formula (2.4) and (2.5), one has

$$\begin{split} &\|(V^{l}(x) - V^{l}(pt - e_{1}t^{2}/2))U_{D}(t)\Phi_{v}\| = \|(V^{l}(x + vt + e_{1}t^{2}/2) - V^{l}(pt + vt + e_{1}t^{2}/2))e^{it\Delta/2}f(p)M_{D,v}(t)\Phi_{0}\|. \\ &\text{Set } \delta = |\hat{v} \cdot e_{1}| < 1. \text{ If } |\xi| \leq \eta \text{ and } \eta/|v| \leq (1 - \delta)/4, \text{ then} \end{split}$$

$$|\xi t + vt + e_1 t^2/2|^2 \ge (1 - \delta^2)|v|^2|t|^2/2 = (c_1|v||t|)^2$$

as well as Eq. (3.4) hold by virtue of Eq. (3.3). Thus we obtain

$$|\xi t + vt + e_1 t^2/2| \ge (c_1 |v||t|)^{\nu} \times (c_2 |t|^2)^{1-\nu} = c_1^{\nu} c_2^{1-\nu} |v|^{\nu} |t|^{2-\nu},$$
(3.9)

where  $0 < \nu < 1$ , which is independent of |v|, will be determined below. Now we introduce  $V_{v,t}^{l}(x)$  as

$$V_{v,t}^{l}(x) = V^{l}(x)g_{\nu}(x/(|v|^{\nu}|t|^{2-\nu})), \qquad (3.10)$$

where  $g_{\nu} \in C^{\infty}(\mathbf{R}^n)$  such that

$$g_{\nu}(x) = \begin{cases} 1, & |x| \ge c_1^{\nu} c_2^{1-\nu} \\ 0, & |x| \le c_1^{\nu} c_2^{1-\nu}/2. \end{cases}$$

If  $|x| < 3|v|^{\rho}|t|$  and  $3|v|^{\rho-1} \le (1-\delta)/4$  for  $0 < \rho < 1$ , then we also obtain

$$|x + vt + e_1 t^2/2| \ge c_1^{\nu} c_2^{1-\nu} |v|^{\nu} |t|^{2-\nu}$$

by virtue of Eq. (2.9). Then we have

$$\begin{aligned} \| (V^{1}(x) - V^{1}(pt - e_{1}t^{2}/2))U_{D}(t)\Phi_{v} \| \\ &= \| (V^{1}(x + vt + e_{1}t^{2}/2) - V^{1}(pt + vt + e_{1}t^{2}/2))e^{it\Delta/2}f(p)M_{D,v}(t)\Phi_{0} \| \\ &= \| (V^{1}(x + vt + e_{1}t^{2}/2) - V^{1}_{v,t}(pt + vt + e_{1}t^{2}/2))e^{it\Delta/2}f(p)M_{D,v}(t)\Phi_{0} \| \end{aligned}$$

$$\leq \| (V^{l}(x+vt+e_{1}t^{2}/2) - V^{l}_{v,t}(x+vt+e_{1}t^{2}/2)) e^{it\Delta/2}f(p)M_{D,v}(t)\Phi_{0} \| \\ + \| (V^{l}_{v,t}(x+vt+e_{1}t^{2}/2) - V^{l}_{v,t}(pt+vt+e_{1}t^{2}/2)) e^{it\Delta/2}f(p)M_{D,v}(t)\Phi_{0} \| \\ \leq I_{1}+I_{2}+I_{3},$$

with

$$I_1 = \sup_{y \in \mathbb{R}^n} |V^{l}(y)| ||F(|x| \ge 3|v|^{\rho}|t|) e^{it\Delta/2} f(p)F(|x| \le |v|^{\rho}|t|) ||_{\mathcal{B}(L^2)} ||M_{D,v}(t)\Phi_0||,$$

$$I_{2} = \sup_{y \in \mathbb{R}^{n}} |V^{1}(y)| ||F(|x| \ge 3|v|^{\rho}|t|) e^{it\Delta/2} f(p)F(|x| > |v|^{\rho}|t|) \langle x \rangle^{-2} ||_{\mathcal{B}(L^{2})} ||\langle x \rangle^{2} M_{D,v}(t) \Phi_{0}||,$$

$$I_{3} = \left\| \left( V_{v,t}^{l}(x+pt+vt+e_{1}t^{2}/2) - V_{v,t}^{l}(pt+vt+e_{1}t^{2}/2) \right) f(p) M_{D,v}(t) \Phi_{0} \right\|_{2}$$

where  $0 \le \rho \le 1$ , which is independent of |v|, will be determined below. By virtue of Proposition 2.1 for estimating  $I_1$  under the condition  $|v|^{\rho} \ge \eta$  and Lemma 3.1, one has

$$I_1 + I_2 \le C(1 + |v|^{\rho}|t|)^{-2}$$

for  $|v|^{\rho} \ge \eta$ , which implies

$$\int_{-\infty}^{\infty} (I_1 + I_2) dt = O(|v|^{-\rho}).$$

By the Baker-Campbell-Hausdorff formula,

$$\begin{split} V^{l}_{v,t}(x+pt+vt+e_{1}t^{2}/2) &- V^{l}_{v,t}(pt+vt+e_{1}t^{2}/2) \\ &= \int_{0}^{1} \big[ (\nabla V_{v,t})(\theta x+pt+vt+e_{1}t^{2}/2)x+it(\Delta V_{v,t})(\theta x+pt+vt+e_{1}t^{2}/2)/2 \big] \mathrm{d}\theta, \end{split}$$

we estimate  $I_3$  as

$$\begin{split} I_{3} &\leq C\{(|v|^{\nu}|t|^{2-\nu})^{-(\gamma_{D}+\mu)} + (|v|^{\nu}|t|^{2-\nu})^{-(\gamma_{D}+1)} + |v|^{-\nu(\gamma_{D}+2\mu)}|t|^{-(2-\nu)(\gamma_{D}+2\mu)+1} + |v|^{-\nu(\gamma_{D}+\mu+1)} \\ &\times |t|^{-(2-\nu)(\gamma_{D}+\mu+1)+1} + |v|^{-\nu(\gamma_{D}+2)}|t|^{-(2-\nu)(\gamma_{D}+2)+1}\}. \end{split}$$

On the other hand, one has

$$I_3 \leq 2 \sup_{y \in \mathbf{R}^n} |V^1(y)| \|\Phi_0\|.$$

Thus, by a straightforward computation, we obtain

$$\begin{split} \int_{-\infty}^{\infty} I_{3} \mathrm{d}t &= O(|v|^{-\nu/(2-\nu)}) + O(|v|^{-\nu(\gamma_{D}+2\mu)/\{(2-\nu)(\gamma_{D}+2\mu)-1\}}) + O(|v|^{-\nu(\gamma_{D}+\mu+1)/\{(2-\nu)(\gamma_{D}+\mu+1)-1\}}) \\ &+ O(|v|^{-\nu(\gamma_{D}+2)/\{(2-\nu)(\gamma_{D}+2)-1\}}) \\ &= O(|v|^{-\nu/(2-\nu)}), \end{split}$$

if  $(2-\nu)(\gamma_D+2\mu)>2$ . Since  $0<\gamma_D\le 1/2$  and  $\gamma_D+\mu>1$ , one has  $\gamma_D+2\mu>2-\gamma_D\ge 3/2$ . Take  $1/2<\rho<1$  such that  $\gamma_D+2\mu>\rho+1$ , and make  $\nu$  as  $\nu/(2-\nu)=\rho$ . Then  $(2-\nu)(\gamma_D+2\mu)>2$  are satisfied by  $\rho+1=2/(2-\nu)$ , and

042101-12 T. Adachi and K. Maehara

----

$$\int_{-\infty}^{\infty} \| (V^{l}(x) - V^{l}(pt - e_{1}t^{2}/2))U_{D}(t)\Phi_{v} \| dt = O(|v|^{-\rho})$$
(3.11)

holds.

By virtue of (3.8), Lemmas 3.2, 3.3, and 3.4, the following lemma can be obtained as Lemma 2.3. Thus we omit the proof.

**Lemma 3.5:** Let v and  $\Phi_{v}$  be as in Theorem 3.1. Let  $\rho$  be as in Lemma 3.4. Then

$$\sup_{t \in \mathbf{R}} \| (e^{-itH} \Omega_D^- - U_D(t)) \Phi_v \| = O(|v|^{-\min\{\alpha, \rho\}})$$
(3.12)

holds as  $|v| \rightarrow \infty$ .

Noting that  $1-2 \min\{\alpha, \rho\} < 0$  by assumption, Theorem 3.1 can be proven in the same way as in the proof of Theorem 2.1. Moreover, Theorem 1.2 can be shown in the same way as in the proof of Theorem 1.2 of Weder<sup>16</sup> (see also Ref. 5). Thus we omit the proofs.

#### ACKNOWLEDGMENT

The first author is partially supported by the Grant-in-Aid for Young Scientists of MEXT 17740078.

- <sup>1</sup>Adachi, T., "Long-range scattering for three-body Stark Hamiltonians," J. Math. Phys. 35, 5547–5571 (1994).
- <sup>2</sup>Adachi, T., and Tamura, H., "Asymptotic completeness for long-range many-particle systems with Stark effect. II," Commun. Math. Phys. **174**, 537–559 (1996).
- <sup>3</sup>Avron, J. E., and Herbst, I. W., "Spectral and scattering theory of Schrödinger operators related to the Stark effect," Commun. Math. Phys. **52**, 239–254 (1977).
- <sup>4</sup>Enss, V., "Propagation properties of quantum scattering states," J. Funct. Anal. 52, 219–251 (1983).
- <sup>5</sup>Enss, V., and Weder, R., "The geometrical approach to multidimensional inverse scattering," J. Math. Phys. **36**, 3902–3921 (1995).
- <sup>6</sup>Graf, G. M., "A remark on long-range Stark scattering," Helv. Phys. Acta 64, 1167–1174 (1991).
- <sup>7</sup>Helgason, S., Groups and Geometric Analysis (Academic, New York, 1984).
- <sup>8</sup>Herbst, I. W., "Unitary equivalence of Stark Hamiltonians," Math. Z. 155, 55–70 (1977).
- <sup>9</sup>Herbst, I. W., Møller, J. S., and Skibsted, E., "Asymptotic completeness for *N*-body Stark Hamiltonians," Commun. Math. Phys. **174**, 509–535 (1996).
- <sup>10</sup> Jensen, A., and Yajima, K., "On the long-range scattering for Stark Hamiltonians," J. Reine Angew. Math. **420**, 179–193 (1991).
- <sup>11</sup>Nicoleau, F., "Inverse scattering for Stark Hamiltonians with short-range potentials," Asymptotic Anal. **35**, 349–359 (2003).
- <sup>12</sup>Perry, P., *Scattering Theory by the Enss Method*, Mathematical Reports Vol. 1 (Harwood Academic, Chur, 1983).
- <sup>13</sup>Reed, M., and Simon, B., *Methods of Modern Mathematical Physics III: Scattering Theory* (Academic, New York, 1979).
- <sup>14</sup> Simon, B., "Phase space analysis of simple scattering system: Extensions of some work of Enss," Duke Math. J. 46, 119–168 (1979).
- <sup>15</sup> Veselić, K., and Weidmann, J., "Potential scattering in a Homogeneous Electrostatic Field," Math. Z. **156**, 93–104 (1977).
- <sup>16</sup>Weder, R., "Multidimensional inverse scattering in an electric field," J. Funct. Anal. **139**, 441–465 (1996).
- <sup>17</sup>White, D., "Modified wave operators and Stark Hamiltonians," Duke Math. J. **68**, 83–100 (1992).
- <sup>18</sup> Yajima, K., "Spectral and scattering theory for Schrödinger operators with Stark-effect," J. Fac. Sci., Univ. Tokyo, Sect. 1A **26**, 377–390 (1979).
- <sup>19</sup>Zorbas, J., "Scattering theory for Stark Hamiltonians involving long-range potentials," J. Math. Phys. **19**, 577–580 (1978).