

PDF issue: 2025-07-11

Proximal translation of the radius following arthroplasty of the distal radioulnar joint in Hajdu-Cheney syndrome

Fujioka, Hiroyuki ; Tanaka, Juichi ; Yoshiya, Shinichi ; Kokubu, Takeshi ; Fujita, Kenji ; Makino, Takeshi ; Mizuno, Kosaku

(Citation) Journal of Shoulder and Elbow Surgery, 12(1):97-100

(Issue Date) 2003-01

(Resource Type) journal article

(Version) Accepted Manuscript

(URL) https://hdl.handle.net/20.500.14094/90000297



1	Proximal Translation of the Radius Following Arthroplasty of
2	the Distal Radioulnar Joint in Hajdu-Cheney Syndrome
3	
4	Hiroyuki Fujioka M.D., Juichi Tanaka M.D.*, Shinichi Yoshiya M.D.,
5	Takeshi Kokubu M.D., Kenji Fujita M.D., Takeshi Makino M.D.,
6	Kosaku Mizuno M.D.
7	
8	
9	Department of Orthopaedic Surgery, Kobe University Graduate School of
10	Medicine, 7-5-1, Kusunoki-cho, Chuo-ku, Kobe 650-0017 Japan
11	
12	* Department of Orthopaedic Surgery, Hyogo Medical College, 1-1, Mukogawa-cho,
13	Nishinomiya, 663-8501 Japan
14	
15	
16	
17	Corresponding Author: Hiroyuki Fujioka
18	Department of Orthopaedic Surgery, Kobe University Graduate School of
19	Medicine
20	Address: 7-5-1, Kusunoki-cho, Chuo-ku, Kobe 650-0017 Japan
21	
	Tel: 81-78-382-5985 Fax: 81-78-351-6930

1 INTRODUCTION

Several surgical procedures for the derangement of the distal radioulnar joint $\mathbf{2}$ (DRUJ) have been developed such as the Darrach's, the Sauvé-Kapandji's, and the 3 Bowers' procedures.^{1,3} After arthroplasty of the DRUJ, limitation in the range of 4 motion and wrist pain have been well documented, however, few reports related to $\mathbf{5}$ 6 problems of the elbow after these surgical procedures have been reported. In the 7 present paper, we report asymptomatic marked proximal translation of the radius at the elbow joint after treatment of dorsal dislocation of the DRUJ using the 8 Sauvé-Kapandji's procedure (S-K procedure) in a patient with Hajdu-Cheney 9 syndrome, an idiopathic skeletal dysplasia with osteoporosis characterized by 10short stature, joint laxity, short clubbed fingers and toes, premature loss of teeth, 11 and defective vision and hearing.^{5,9,13} 12

13

14 CASE REPORT

The patient was a 46-year-old woman. The patient was the first-born and the 15product of a full-term normal delivery. There was no family history of bone disease 1617and development of her intelligence was normal. In the patient's later childhood the patient's was diagnosed as having the Hajdu-Cheney syndrome from short 1819 stature, osseous dysplasia with osteoporosis, short clubbed fingers and toes, premature loss of teeth, defective hearing, acro-osteolysis, wide cranial sutures, 20and multiple wormian bones. The patient was right handed and had experienced 21mild pain in the right wrist and inability to extend the little finger 6 months 22before the patient visited us. Although the wrist pain did not increased, the 23patient experienced inability to extend the middle and ring fingers and visited us. 24

1 Physical examination revealed dorsal displacement and tenderness of the distal end of the ulna. Full range of motion in extension and flexion of the wrist and $\mathbf{2}$ elbow, or in pronation of the forearm was noted. However supination of the 3 forearm was limited to 60° due to wrist pain. Active motion of the thumb and the 4 index finger was not limited. The patient could fully flex all joints of the middle, $\mathbf{5}$ 6 ring and little fingers, and extend their distal and proximal interphalangeal joints. 7 However, extension of their metacarpophalangeal joints was restricted to 50° (Figure 1). Passive range of motion of these fingers was fully maintained. Relief 8 and tenodesis effects of the extensor tendons of the fingers were not detected 9 clearly. Grip strength in the right hand was 12 kg and in the unaffected left hand, 1019kg. General joint laxity was detected. Anteroposterior view of the radiograph of 11 the hand showed osteolysis of the distal phalanges, positive variance and 12deformity of the ulna and dissociation of the DRUJ (Figure 2). The radiograph of 13 14the forearm showed slight bowing deformity of the ulna and incongruity of the humeroradial joint, however, discrepancy of the length of these paired bones was 15not evident (Figure 3). 16

17The patient was treated surgically. A curved dorsal incision was made on the dorsal aspect of the wrist and the extensor retinaculum was incised over the 18extensor digitorum communis (EDC). Very mild tenosynovitis and rupture of the 19 extensor tendons, EDC (III, IV, V) and extensor digiti minimi, was observed. The 20ruptured extensor tendons were reconstructed with a tendon graft.³ The graft of 21the palmaris longus was connected to the distal tendon stumps by an intercalary 22weave and then attached to the proximal tendon with an interlace weave 23technique in slightly greater extension than the normal cascade. Dorsal 24

dislocation of the DRUJ was managed by the S-K procedure.¹ A bone segment of the ulna was excised in order to create a space for the pseudoarthrosis site and articular surfaces of the radius and the ulna were excised until cancellous bone. The head of the ulna was opposed to the radius and pinned to it with two Kirschner wires.

6 The patient started active finger exercises at 3 weeks postoperative. By 6 months 7 after surgery, there was no pain or limitation in the range of motion of the wrist or any of the fingers. Two years after surgery, the patient had no complaint 8 concerning the patient's wrist, fingers, and elbow with full functional recovery. 9 Radiographs of the wrist showed complete arthrodesis of the DRUJ and the 10pseudoarthrosis of the ulna (Figure 4). However, radiographs of the elbow 11 showed marked anterior dislocation of the radial head and proximal translation of 12the radius (Figure 5). 13

14

15 DISCUSSION

Although Hajdu-Cheney syndrome is a congenital disorder, the correct diagnosis 1617is rarely made until later childhood when characteristic symptoms occur. Clinical features included distinctive facial appearance, short stature, generalized 18osteoporosis, joint laxity, premature loss of teeth, retarded puberty, and defective 19 vision and hearing.^{5,9,13} On the radiographs, osteolysis of the distal phalanges of 20the hands and feet, wide cranial sutures, and multiple wormian bones were 21observed. Mostly all of the clinical features were detected in the patient described 22in the present report. 23

24 Rupture of the extensor tendons and disorder of the wrist joint has not been

reported in this syndrome. Rupture of the extensor tendons in rheumatoid arthritis is caused by the mechanical force associated with deformity of the wrist and tenosynovitis.³ In the case presented, tenosynovitis of the extensor tendons was not severe and rupture occurred in the dominant hand. Mechanism of extensor tendon rupture was mainly considered to the mechanical force by the dorsal dislocated ulna and it was successfully treated with an intercalary tendon graft using the palmaris longus.

In Hajdu-Cheney syndrome, bowing deformities and cortical defects of the paired long bones, for example the tibia, fibula, radius, and ulna causes incongruity and arthrosis of the knee and ankle joint at both ends of these paired bones.^{5,9} In the case presented, cortical defects of the radius and the ulna were not observed, however, bowing deformity was observed in the ulna.

Proximal translation of the radius was caused by resection of the radial head due 13 to lack of bony stabilization at the humeroradial joint.^{4,7,10} Fracture of the radial 14head and injury of the interosseous membrane caused translation of the radius 15with limitation in the range of motion of the wrist, forearm, and elbow.^{2,6} A case 16report of osteoarthritis of the humeroradial joint due to proximal migration of the 1718radius 25 years after an extensive Darrach's procedure at the age of the 15 19 suggested that the interosseous membrane plays an important role to the proximal translation of the radius.¹¹ The patients who had the posttraumatic 2021disorder of the wrist with simultaneous involvement of the elbow was treated by the S-K procedure associated with insertion of a silicone implant of the radial 22head and they obtained a good functional recover of the wrist and elbow joints.¹² 23Biomechanical analysis of pressure distribution in the humeroradial joint using 24

fresh frozen human cadavers revealed that the force transmitted to the capitellum increased after the S-K procedure, however, that there was no significant differences between the forces before and after the S-K procedure.⁸ By contrast the force transmitted to the capitellum after the S-K procedure with incision of the interosseous membrane significantly increased compared with the forces before the opertation.

7 In the patient reported here, anterior subluxation and proximal translation of the radial head had been promoted severely after surgical treatment of the DRUJ by 8 the S-K procedure. It was speculated that dorsal dislocation of the ulna at the 9 DRUJ and subluxation of the radial head was caused by incongruity of the 10radioulnar joint due to deformity of the forearm bones, joint laxity, and 11 insufficiency of the interosseous membrane. Instability of the wrist due to the 12pseudoarthrosis site at the distal ulna created by the S-K procedure promoted 13 14proximal translation of the radius and instability of the forearm. The patient has no complaint of the elbow at present. However, we should monitor closely 15translation of the radius after arthroplasty of the wrist in the patient with general 1617joint laxity and insufficiency of the interosseous membrane such as observed in Hajdu-Cheney syndrome because there is the possibility of pain or instability of 18the elbow occurring. We think that the wrist arthroplasty with resection of the 19 ulna such as the S-K procedure should be performed very carefully in the patient 2021with general joint laxity and insufficiency of the interosseous membrane.

1 REFERENCES

2	1.	Bowers WH. The distal radioulnar joint. In Green DP, editor. Operative Hand
3		Surgery 3 rd edition. New York; Churchill Livingstone; 1993. p. 973-1019.
4	2.	Essex-Lopresti P. Fractures of the radial head with distal radio-ulnar
5		dislocation. Report of two cases. J Bone Joint Surg Br 1951; 33: 244-7.
6	3.	Felden P, Millender LH, Nalebuff EA. Rheumatoid arthritis in the hand and
7		wrist. In Green DP, editor. Operative Hand Surgery 3rd edition. New York;
8		Churchill Livingstone; 1993. p. 1587-690.
9	4.	Goldberg I, Peylan J, Yosipovitch Z. Late results of excision of the radial head
10		for an isolated closed fracture. J Bone Joint Surg Am 1986; 68: 675-9.
11	5.	Herscovici Jr. D, Bowen JR, Scott Jr. CI. Cervical instability as an unusual
12		manifestation of Hadju-Cheney syndrome of acroosteolysis. Clin Orthop 1990;
13		255: 111-6.
14	6.	Hotchkiss RN. Injuries to the interosseous ligament of the forearm. Hand Clin
15		1994; 10: 391-8.
16	7.	McDougall A, White J. Subluxation of the inferior radio-ulnar joint
17		complicating fracture of the radial head . J Bone Joint Surg Br 1957; 39:
18		278-87.
19	8.	Ofuchi S, Takahashi K, Yamagata M, Rokkaku T, Moriya H, Hara T. Pressure
20		distribution in the humeroradial joint and force transmission to the capitellum
21		during rotation of the forearm: effects of the Sauvé-Kapandji procedure and
22		incision of the interosseous membrane. J Orthop Sci 2001; 6: 33-8.
23	9.	O'Reilly MAR, Shaw DG. Hajdu-Cheney syndrome. Ann Rheum Dis 1994; 53:
24		276-9.

 $\overline{7}$

1	10. Sowa DT, Hotchikiss RN, Weiland AJ. Symptomatic proximal translation of
2	the radius following radial head resection. Clin Orthop 1995; 317: 106-13.
3	11. Stuer P, De Smet L, Fabry G. Osteoarthritis of the elbow following an
4	extensive Darrach procedure. Acta Orthop Belgica 1996; 62: 62-5.
5	12. Taleisnik J. The Sauvé-Kapandji's procedure. Clin Orthop 1992; 275: 110-23.
6	13. Weleber RG, Beals RK. The Hajdu-Cheney syndrome. J Pediatr 1976; 88:
7	243-9.

1	LEGENDS
2	Figure 1: The clinical appearance of the right hand.
3	Extension of the MP joints of the middle, ring, and little fingers was limited.
4	
5	Figure 2: Radiographs of the hand (R indicates the right hand and L the left).
6	The anteroposterior view of the hand showed osteolysis of the distal phalanges,
7	positive variance and deformity of the ulna, and dissociation of the DRUJ.
8	
9	Figure 3. The radiographs of the forearm. Anterolateral (A) and lateral (B) view.
10	Bowing deformity of the ulna and incongruity of the humeroradial joint were
11	observed, however, there was no discrepancy of length of these paired bones.
12	
13	Figure 4: Radiograph of the wrist after surgical treatment. Anterorposterior (A)
14	and lateral (B) view.
15	Complete arthrodesis of the DRUJ and the pseudoarthrosis of the ulna was
16	shown.
17	
18	Figure 5: Radiograph of the elbow after surgical treatment of the wrist.
19	Anterorposterior (A) and lateral (B) view.
20	Marked proximal translation and anterior dislocation of the radial head were
21	shown.