

PDF issue: 2025-02-22

Risk comparison of the Stein-rule estimator in a linear regression model with omitted relevant regressors and multivariatet errors under the Pitman nearness criterion

Namba, Akio Ohtani, Kazuhiro

(Citation) Statistical Papers, 48(1):151-162

(Issue Date) 2007-01

(Resource Type) journal article

(Version) Accepted Manuscript

(URL) https://hdl.handle.net/20.500.14094/90000338



# Risk Comparison of the Stein-Rule Estimator in a Linear Regression Model with Omitted Relevant Regressors and Multivariate *t* Errors under the Pitman Nearness Criterion\*

Akio Namba and Kazuhiro Ohtani

Graduate School of Economics Kobe University Rokko, Nada-ku, Kobe 657-8501 Japan

#### Abstract

In this paper we consider a linear regression model with omitted relevant regressors and multivariate t error terms. The explicit formula for the Pitman nearness criterion of the Stein-rule (SR) estimator relative to the ordinary least squares (OLS) estimator is derived. It is shown numerically that the dominance of the SR estimator over the OLS estimator under the Pitman nearness criterion can be extended to the case of the multivariate t error distribution when the specification error is not severe. It is also shown that the dominance of the SR estimator over the OLS estimator cannot be extended to the case of the multivariate t error distribution when the specification error is severe.

<sup>\*</sup>This research is partially supported by the Grants-in-Aid for 21st Century COE program.

# **1** Introduction

In the context of linear regression, the ordinary least squares (OLS) estimator is known as the best linear unbiased estimator (BLUE). However, the Stein-rule (SR) estimator for regression coefficients proposed by Stein (1956) and James and Stein (1961) dominates the OLS estimator in terms of mean squared error (MSE) of prediction. Since the findings of Stein (1956) and James and Stein (1961), lots of shrinkage estimators have been proposed, and their sampling properties have been examined.

When shrinkage estimators are compared with the OLS estimator, the MSE has usually been used as a criterion of comparison. However, there are several studies in which another criterion is used to compare the sampling performances of these estimators. For example, using the Pitman nearness criterion, Keating and Czitrom (1989) showed by numerical evaluations that the SR estimator is uniformly preferred to the OLS estimator. This indicates that the dominance of the SR estimator over the OLS estimator holds even when the Pitman nearness criterion is used instead of MSE. The Pitman nearness criterion is also used in many studies. Some examples are Rao, Keating and Mason (1988), Sen, Kubokawa and Saleh (1989), Srivastava and Srivastava (1993), and Chaturvedi and Bhatti (1998).

Although it is assumed in these studies that the model is specified correctly, the model may be specified incorrectly. One example of such a specification error is to exclude some relevant independent variables in the specified model. It is also assumed in these studies that the error terms obey a normal distribution. However, as is stated in Fama (1965), and Blattberg and Gonedes (1974), many economic data may be generated by a distribution with fatter tails than a normal distribution. One example of such a distribution is a multivariate t distribution.

Mittelhammer (1984), Ohtani (1993b), and Namba (2002) examined the MSE performances of the Stein-rule (SR) estimator and its variants when relevant independent variables are omitted. They showed that although the SR estimator has smaller MSE than the OLS estimator in a wide region of the parameter space, the dominance of the SR estimator over the OLS estimator does not hold necessarily when there are omitted variables. On the other hand, there are many studies on the sampling performances of estimators when error terms in a linear regression model obey a multivariate t distribution. Some examples

are Zellner (1976), Prucha and Kelejian (1984), Ullah and Zinde-Walsh (1984), Judge et al. (1985), Sutradhar and Ali (1986), Sutradhar (1988), Singh (1988, 1991), Giles (1991, 1992), Ohtani (1991, 1993a), Ohtani and Giles (1993), Ohtani and Hasegawa (1993), Namba (2001), and Namba and Ohtani (2002).

In this paper we consider a linear regression model when relevant independent variables are omitted in the specified model and error terms obey a multivariate *t* distribution. Using the above model, we compare the sampling performances of the SR estimator and the OLS estimator under the Pitman nearness criterion. Thus, our analysis is an extension of Keating and Czitrom (1989) to two directions: omitted relevant independent variables and multivariate *t* error terms. In the next section we introduce a model and estimators. In section 3 we derive the exact formula of the Pitman nearness criterion of the SR estimator relative to the OLS estimator. In section 4, using this exact formula, we compare the SR estimator and the OLS estimator numerically. Our numerical results show that the dominance of the SR estimator over the OLS estimator under the Pitman nearness criterion can be extended to the case of the multivariate *t* error distribution when the specification error is not severe. The numerical results also show that the dominance of the SR estimator over the OLS estimator over the OLS estimator over the OLS estimator over the OLS estimator error is not severe.

#### 2 Model and Estimators

Consider a linear regression model,

$$y = X_1\beta_1 + X_2\beta_2 + u$$
  
=  $X\beta + u$ , (1)

where y is an  $n \times 1$  vector of observations on a dependent variable,  $X_1$  and  $X_2$  are  $n \times k_1$  and  $n \times k_2$ matrices of observations on nonstochastic independent variables and  $X = [X_1, X_2]$ ,  $\beta_1$  and  $\beta_2$  are  $k_1 \times 1$ and  $k_2 \times 1$  vectors of regression coefficients and  $\beta' = [\beta'_1, \beta'_2]$ . We assume that  $X_1$  and  $X = [X_1, X_2]$  are of full column rank. Also, in this paper, we assume that u has a multivariate t distribution with the probability density function (p.d.f) given by

$$p(u|\nu,\sigma) = \frac{\nu^{\nu/2}\Gamma((\nu+n)/2)}{\pi^{n/2}\Gamma(\nu/2)\sigma^n} \frac{1}{\{\nu + u'u/\sigma^2\}^{(n+\nu)/2}},$$
(2)

where  $\sigma$  is a scale parameter and  $\nu$  is a degrees of freedom parameter. It is easy to show that E[u] = 0 and  $E[uu'] = [\nu\sigma^2/(\nu-2)]I_n$  for  $\nu > 2$ . When  $\nu \to \infty$ , the p.d.f of *u* approaches that of a normal distribution with mean 0 and covariance matrix  $\sigma^2 I_n$ . As is shown in Zellner (1976), the multivariate *t* distribution can be viewed as a mixture of multivariate normal and inverted gamma distributions:

$$p(u|\nu,\sigma) = \int_0^\infty P_N(u|\tau) P_{IG}(\tau|\nu,\sigma) d\tau,$$
(3)

where

$$P_N(u|\tau) = (2\pi\tau^2)^{-n/2} \exp[-u'u/2\tau^2],$$
(4)

$$P_{IG}(\tau|\nu,\sigma) = \frac{2(\nu\sigma^2/2)^{\nu/2}}{\Gamma(\nu/2)} \tau^{-(\nu+1)} \exp[-\nu\sigma^2/2\tau^2].$$
(5)

Suppose that the matrix of regressors  $X_2$  is omitted mistakenly and the model is specified as

$$y = X_1\beta_1 + u^*, \quad u^* = X_2\beta_2 + u.$$
 (6)

Then, based on the misspecified model, the ordinary least squares (OLS) estimator of  $\beta_1$  is

$$b_1 = S_{11}^{-1} X_1' y, (7)$$

where  $S_{11} = X'_1 X_1$ . Also, the Stein-rule (SR) estimator proposed by Stein (1956) and James and Stein (1961) is defined as

$$b_{SR1} = \left(1 - \frac{ae_1'e_1}{b_1'S_{11}b_1}\right)b_1,\tag{8}$$

where  $e_1 = y - X_1 b_1$ , and *a* is a constant such that  $0 \le a \le 2(k_1 - 2)/(n - k_1 + 2)$ . If we use the loss function

$$L(\bar{\beta}_1) = (X_1\bar{\beta}_1 - X\beta)'(X_1\bar{\beta}_1 - X\beta), \tag{9}$$

where  $\bar{\beta}_1$  is any estimator of  $\beta_1$ , and no relevant regressors are omitted, then Stein (1956) showed that the SR estimator dominates the OLS estimator. Moreover, as is shown in James and Stein (1961), if there are no omitted regressors, the MSE of the SR estimator is minimized when  $a = (k_1 - 2)/(n - k_1 + 2)$ . Thus, we use this value of *a* hereafter. Also, Ohtani (1993a) and Namba (2002) showed by numerical evaluations that although the SR estimator has smaller MSE than the OLS estimator in a wide region of the parameter space, the dominance of the SR estimator over the OLS estimator does not hold necessarily when there are omitted variables. In the next section, we derive the explicit formula for the Pitman nearness criterion of the SR estimator relative to the OLS estimator.

### **3** Pitman Nearness Criterion

Using (9), the Pitman nearness criterion of the SR estimator relative to the OLS estimator is given by

$$PN(b_{SR1}, b_1) = Pr[L(b_{SR1}) \le L(b_1)],$$
(10)

where  $Pr[\cdot]$  denotes the probability and  $L(\cdot)$  is defined in (9). If  $PN(b_{SR1}, b_1) \ge 1/2$ , the SR estimator is preferable to the OLS estimator in the Pitman nearness sense.

If  $k_1$  is an even integer, we can derive the explicit formula of  $PN(b_{SR1}, b_1)$  in a similar way to Keating and Czitrom (1989). As is shown in appendix, the explicit formula for  $PN(b_{SR1}, b_1)$  is

$$PN(b_{SR1}, b_1) = \nu^{\nu/2} \sum_{i=0}^{\infty} \sum_{j=0}^{m+i-1} \frac{\theta_1^{m+2i-r-1} \theta_2^j}{(2\theta_1 + \theta_2 + \nu)^{\nu/2 + m + 2i + j - r - 1}} \\ \times \frac{\Gamma(\nu/2 + m + 2i + j - r - 1)}{\Gamma(\nu/2)\Gamma(m + i - r)i!j!} [1 - I_{\frac{a}{2+a}}(r + 1, n/2 - m + j)],$$
(11)

where  $\theta_1 = \beta' S \beta / 4\sigma^2$ ,  $\theta_2 = \beta' X' M_1 X \beta / \sigma^2$ ,  $M_1 = I - X_1 S_{11}^{-1} X'_1$ ,  $m = k_1/2$ ,  $k_1$  is an even integer and  $I_x(a,b)$  is the incomplete beta function ratio defined as

$$I_x(a,b) = \frac{1}{B(a,b)} \int_0^x t^{a-1} (1-t)^{b-1} dt.$$
(12)

Using this formula, we examine the Pitman nearness of the SR estimator relative to the OLS estimator in the next section.

# 4 Numerical Analysis

In this section, we compare the SR estimator with the OLS estimator based on the Pitman nearness criterion using (11). The noncentrality parameters are expressed as  $\theta_1 = \theta/4$  and  $\theta_2 = \theta(1 - R_1^2)$ , where  $\theta = \beta' S \beta / \sigma^2$ ,  $R_1^2 = \beta' X' X_1 S_{11}^{-1} X'_1 X \beta / \beta' S \beta$  and S = X' X. As is discussed in Ohtani (1993a),  $\theta$  is the noncentrality parameter which appeared in the test for the null hypothesis that all the regression coefficients are zeros in the correctly specified model. Also,  $R_1^2$  is interpreted as the coefficient of determination in regression of  $X\beta$  on  $X_1$ . Thus, if  $R_1^2$  is close to unity, the magnitude of model misspecification is regarded as small, and vice versa. Thus, in the numerical evaluations, we use the following parameter values;  $\theta$  = various values,  $R_1^2 = 0.1, 0.3, 0.5, 0.7, 0.9, 1.0, m = 2, 3, 4, n = 20, 30, 40, and v = 3, 5, 7, 10,$  $20, 30, <math>\infty$ . The numerical evaluations were executed on a personal computer using the FORTRAN code. The double infinite series in (11) were judged to converge when the increment of the series got smaller than  $10^{-12}$ .

The results for m = 2, 4, n = 20, 30, and  $\nu = 3, 20$  are shown in Tables 1 to 4. We see from Table 1 that the SR estimator is preferred to the OLS estimator for all values of  $\theta$  and  $R_1^2$  considered here when m = 2, n = 20, and v = 3. In particular, when  $\theta$  is close to 0 and  $R_1^2$  is close to 1.0, the SR estimator is much preferred to the OLS estimator. However, the efficiency gets small as  $\theta$  gets large and  $R_1^2$  gets small. When v increases from 3 to 20, the OLS estimator is preferred to the SR estimator for  $\theta = 50.0$ and  $R_1^2 = 0.1$ . This indicates that as the tails of the error distribution get flat, the preference for the SR estimator gets strong under the Pitman nearness criterion. We see from Table 2 that as n increases from 20 to 40, the SR estimator is preferred to the OLS estimator for all values of  $\theta$  and  $R_1^2$  considered here even when v = 20, though the performance of efficiency is similar to that for n = 20. We see from Table 3 that the OLS estimator is preferred to the SR estimator for large values of  $\theta$  and small values of  $R_1^2$ when m = 4, n = 20, and v = 3. In particular, when  $\theta$  is close to 50.0 and  $R_1^2$  is close to 0.1, the OLS estimator is much preferred to the SR estimator. This indicates that the region of  $\theta$  and  $R_1^2$  such that the SR estimator is preferred to the OLS estimator gets much narrow as *m* increases from 2 to 4. However, we see from Table 4 that as n increases from 20 to 40, the SR estimator is preferred to the OLS estimator for all values of  $\theta$  and  $R_1^2$  considered here, though the performance of efficiency is similar to that for m =2, n = 20 and v = 3 given in Table 1.

From the above results, we see that when the specification error is not severe ( $R_1^2$  is close to 1.0), the SR estimator is preferred to the OLS estimator under the Pitman nearness criterion even when the error terms obey the multivariate *t* distribution. Thus, the dominance of the SR estimator over the OLS estimator under the Pitman nearness criterion can be extended to the case of the multivariate *t* error distribution when the specification error is not severe. The results also show that when  $R_1^2$  is close to 0.1, the OLS estimator is preferred to the SR estimator for some values of  $\theta$ , though the region of  $\theta$  and  $R_1^2$  such that the OLS estimator is preferred to the SR estimator gets narrow as *n* increases. Thus, the dominance of the SR estimator over the OLS estimator under the Pitman nearness criterion cannot be extended to the case of the multivariate *t* error distribution when the specification error is severe.

# Appendix

Using (7), (8) and (9), we have

$$\Pr[L(b_{SR1}) \leq L(b)] = \Pr\left[\left(X_1b_1 - \frac{X\beta}{2}\right)'\left(X_1b_1 - \frac{X\beta}{2}\right) \geq \frac{\beta'S\beta}{4} + \frac{ae_1'e_1}{2}\right].$$
(13)

First, we derive the formula for  $\Pr[L(b_{SR1}) \leq L(b_1)|\tau]$ , assuming that  $\tau$  is given. If we define  $v_1 = (X_1b_1 - X\beta/2)'(X_1b_1 - X\beta/2)/\tau^2$  and  $v_2 = e'_1e_1/\tau^2$ , then  $v_1 \sim \chi_{k_1}'^2(\lambda_1)$  and  $v_2 \sim \chi_{n-k_1}'^2(\lambda_2)$  for given  $\tau$ , where  $\chi_f'^2(\lambda)$  is the noncentral chi-square distribution with f degrees of freedom and noncentrality parameter  $\lambda$ ,  $\lambda_1 = \beta'S\beta/4\tau^2$ , and  $\lambda_2 = \beta'X'M_1X\beta/\tau^2$ . Thus, using  $v_1$  and  $v_2$ , we have

$$\Pr[L(b_{SR1}) \leq L(b_1)|\tau] = \Pr\left[\frac{v_1 - \lambda_1}{v_2} \geq \frac{a}{2}|\tau\right].$$
(14)

Since  $v_1$  and  $v_2$  are mutually independent for given  $\tau$ , we have

$$\Pr\left[\frac{v_1 - \lambda_1}{v_2} \ge \frac{a}{2} \middle| \tau\right] = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} K_{ij} \iint_{\mathcal{R}} v_1^{k_1/2 + i - 1} v_2^{(n-k_1)/2 + j - 1} \exp\left[-\frac{v_1 + v_2}{2}\right] dv_1 dv_2,$$
(15)

where *R* is the region such that  $(v_1 - \lambda_2)/v_2 \ge a/2$ ,

$$K_{ij} = \frac{w_i(\lambda_1)w_j(\lambda_2)}{2^{n/2+i+j}\Gamma(k_1/2+i)\Gamma((n-k_1)/2+j)},$$
(16)

and  $w_i(\lambda) = \exp(-\lambda/2)(\lambda/2)^i/i!$ .

Making use of the change of variables,  $w = (v_1 - \lambda_1)/v_2$  and  $z = v_2$ , the integral in (15) reduces to

$$\int_{0}^{\infty} \int_{a/2}^{\infty} (wz + \lambda_1)^{k_1/2 + i - 1} z^{(n-k_1)/2 + j} \exp\left[-\frac{wz + z + \lambda_1}{2}\right] dw \, dz.$$
(17)

When  $k_1 = 2m$ , where *m* is a positive integer, (17) reduces to

$$\sum_{r=0}^{m+i-1} {}_{m+i-1}C_r \lambda_1^{m+i-1-r} \exp(-\lambda_1/2) \int_0^\infty \int_{a/2}^\infty w^r z^{n/2+r-m+j} \exp\left[-\frac{(1+w)z}{2}\right] dw \, dz,\tag{18}$$

where  $_{p}C_{m} = \frac{p!}{m!(p-m)!}$ .

Again, making use of the change of a variable,  $z_1 = (1 + w)z/2$ , the integral in (18) reduces to

$$2^{n/2+r-m+j+1}\Gamma(n/2+r-m+j+1)\int_{a/2}^{\infty}\frac{w^r}{(1+w)^{n/2+r-m+j+1}}dw.$$
(19)

Further, making use of the change of a variable, t = w/(1 + w), we obtain

$$\Pr\left[\frac{v_1 - \lambda_1}{v_2} \ge \frac{a}{2} \middle| \tau\right] = \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \sum_{r=0}^{m+i-1} \frac{(\lambda_1/2)^{m+2i-r-1} (\lambda_2/2)^j \exp[-(\lambda_1 + \lambda_2/2)]}{\Gamma(m+i-r)i! j!} \times \left[1 - I_{\frac{a}{2+a}}(r+1, n/2 - m+j)\right].$$
(20)

Utilizing the relation,

$$\Pr[L(b_{SR1}) \leq L(b)] = \int_0^\infty \Pr[L(b_{SR1}) \leq L(b)|\tau] P_{IG}(\tau|\nu,\sigma) d\tau$$
$$= \int_0^\infty \Pr\left[\frac{\nu_1 - \lambda_1}{\nu_2} \geq \frac{a}{2} |\tau\right] P_{IG}(\tau|\nu,\sigma) d\tau,$$
(21)

and making use of the change of variable,  $t_1 = (2\eta_1 + \eta_2 + \nu\sigma^2)/2\tau^2$ , where  $\eta_1 = \beta' S\beta/4$  and  $\eta_2 = \beta' X' M_1 X\beta$ , we obtain (11) in the text.

## References

- Blattberg, R., and N. Gonedes, 1974, "A Comparison of the Stable and Student Distributions as Statistical Models for Stock Prices," *Journal of Business*, pp. 244–280.
- Chaturvedi, A., and M. I. Bhatti, 1998, "On the Double *k*-Class Estimators in Linear Regression," *Journal* of *Quantitative Economics*, 14, 53–58.
- Fama, E., 1965, "The Behavior of Stock Market Prices," Journal of Business, 38, 34–105.
- Giles, J. A., 1991, "Pre-testing for Linear Restrictions in a Regression Model With Spherically Symmetric Disturbances," *Journal of Econometrics*, 50, 377–398.
- Giles, J. A., 1992, "Estimation of the Error Variance After a Preliminary-test of Homogeneity in a Regression Model With Spherically Symmetric Disturbances," *Journal of Econometrics*, 53, 345–361.
- James, W., and C. Stein, 1961, "Estimation With Quadratic Loss," in *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability*, vol. 1, pp. 361–379. University of California Press, Berkeley.
- Judge, G., S. Miyazaki, and T. Yancey, 1985, "Minimax Estimators for the Location Vectors of Spherically Symmetric Densities," *Econometric Theory*, 1, 409–417.

- Keating, J. P., and V. Czitrom, 1989, "A Comparison of James-Stein Regression With Least Squares in the Pitman Nearness Sense," *Journal of Statistical Computation and Simulation*, 34, 1–9.
- Keating, J. P., and R. L. Mason, 1988, "James-Stein Estimation from an Alternative Perspective," American Statistician, 42, 160–164.
- Mittelhammer, R. C., 1984, "Restricted Least Squares, Pre-test, OLS and Stein Rule Estimators: Risk Comparisons Under Model Misspecification," *Journal of Econometrics*, 25, 151–164.
- Namba, A., 2001, "MSE Performance of the 2SHI Estimator in a Regression Model with Multivariate *t* Error Terms," *Statistical Papers*, 42, 81–96.
- Namba, A., 2002, "PMSE Performance of the Biased Estimators in a Linear Regression Model When Relevant Regressors are Omitted," *Econometric Theory*, 18, 1086–1098.
- Namba, A., and K. Ohtani, 2002, "MSE Performance of the Double k-class Estimator of Each Individual Regression Coefficient under Multivariate t-Errors," in Aman Ullah, Alan T.K. Wan, and Anoop Chaturvedi (ed.), Handbook of Applied Econometrics and Statistical Inference . chap. 15, pp. 305– 326, Marcel Dekker.
- Ohtani, K., 1991, "Small Sample Properties of the Interval Constrained Least Squares Estimator When Error Terms Have a Multivariate *t* Distribution," *Journal of the Japan Statistical Society*, 21, 197–204.
- Ohtani, K., 1993a, "A Comparison of the Stein-rule and Positive-part Stein-rule Estimators in a Misspecified Linear Regression Model," *Econometric Theory*, 9, 668–679.
- Ohtani, K., 1993b, "Testing for Equality of Error Variances between Two Linear Regressions with Independent Multivariate *t* Errors," *Journal of Quantitative Economics*, 9, 85–97.
- Ohtani, K., 1996, "Further Improving the Stein-rule Estimator Using the Stein Variance Estimator in a Misspecified Linear Regression Model," *Statistics and Probability Letters*, 29, 191–199.
- Ohtani, K., 1998, "MSE Performance of the Minimum Mean Squared Error Estimators in a Linear Regression Model when Relevant Regressors are Omitted," *Journal of Statistical Computation and Simulation*, 61, 61–75.
- Ohtani, K., and J. Giles, 1993, "Testing Linear Restrictions on Coefficients in a Linear Regression Model With Proxy Variables and Spherically Symmetric Disturbances," *Journal of Econometrics*, 57, 393– 406.
- Ohtani, K., and H. Hasegawa, 1993, "On Small Sample Properties of  $R^2$  in a Linear Regression Model With Multivariate *t* Errors and Proxy Variables," *Econometric Theory*, 9, 504–515.
- Prucha, I. R., and H. H. Kelejian, 1984, "The Structure of Simultaneous Equation Estimators: A Generalization Towards Nonnormal Disturbances," *Econometrica*, 52, 721–736.
- Rao, C., J. Keating, and R. L. Mason, 1986, "The Pitman Nearness Criterion and Its Determination," *Communications in Statistics-Theory and Methods*, 15, 3173–3196.

- Sen, P. K., T. Kubokawa, and A. E. Saleh, 1989, "The Stein Paradox in the Sense of the Pitman Measure of Closeness," *Annals of Statistics*, 17, 1375–1386.
- Singh, R. S., 1988, "Estimation of Error Variance in Linear Regression Models with Error Having Multivariate Student-*t* Distribution with Unknown Degrees of Freedom," *Economics Letters*, 27, 47–53.
- Singh, R. S., 1991, "James-Stein Rule Estimators in Linear Regression Models With Multivariate-*t* Distributed Error," *The Australian Journal of Statistics*, 33, 145–158.
- Srivastava, A., and V. Srivastava, 1993, "Pitman Closeness for Stein-rule Estimators of Regression Coefficient," *Statistics and Probability Letters*, 18, 85–89.
- Stein, C., 1956, "Inadmissibility of the Usual Estimator for the Mean of a Multivariate Normal Distribution," in *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability*, vol. 1, pp. 197–206. University of California Press, Berkeley.
- Sutradhar, B. C., 1988, "Testing Linear Hypothesis With t Error Variable," Sankhyā, Series B, Indian Journal of Statistics, 50, 175–180.
- Sutradhar, B. C., and M. M. Ali, 1986, "Estimation of the Parameters of a Regression Model With a Multivariate *t* Error Variable," *Communications in Statistics-Theory and Methods*, 15, 429–450.
- Ullah, A., and V. Zinde-Walsh, 1984, "On the Robustness of LM, LR, and W Tests in Regression Models," *Econometrica*, 52, 1055–1066.
- Zellner, A., 1976, "Bayesian and Non-Bayesian Analysis of the Regression Model with Multivariate Student-*t* Error Terms," *Journal of the American Statistical Association*, 71, 400–405.

		$R_1^2$					
ν	$\theta$	0.1	0.3	0.5	0.7	0.9	1.0
3	0.0	0.9221	0.9221	0.9221	0.9221	0.9221	0.9221
	1.0	0.8893	0.8910	0.8926	0.8943	0.8959	0.8967
	2.0	0.8600	0.8632	0.8664	0.8695	0.8727	0.8743
	4.0	0.8125	0.8182	0.8239	0.8297	0.8354	0.8382
	6.0	0.7759	0.7837	0.7914	0.7992	0.8069	0.8108
	8.0	0.7466	0.7561	0.7655	0.7750	0.7845	0.7892
	10.0	0.7223	0.7333	0.7442	0.7552	0.7662	0.7717
	15.0	0.6757	0.6896	0.7037	0.7179	0.7320	0.7391
	20.0	0.6412	0.6576	0.6743	0.6910	0.7078	0.7162
	25.0	0.6139	0.6325	0.6513	0.6703	0.6894	0.6990
	30.0	0.5914	0.6119	0.6326	0.6536	0.6747	0.6853
	40.0	0.5556	0.5792	0.6033	0.6278	0.6525	0.6649
	50.0	0.5275	0.5538	0.5808	0.6083	0.6362	0.6501
20	0.0	0.9221	0.9221	0.9221	0.9221	0.9221	0.9221
	1.0	0.8888	0.8905	0.8921	0.8938	0.8955	0.8963
	2.0	0.8572	0.8605	0.8638	0.8671	0.8704	0.8720
	4.0	0.8025	0.8087	0.8149	0.8211	0.8273	0.8303
	6.0	0.7589	0.7674	0.7760	0.7845	0.7930	0.7972
	8.0	0.7239	0.7344	0.7449	0.7554	0.7658	0.7710
	10.0	0.6954	0.7075	0.7196	0.7318	0.7439	0.7500
	15.0	0.6422	0.6577	0.6732	0.6888	0.7044	0.7122
	20.0	0.6047	0.6228	0.6411	0.6595	0.6779	0.6871
	25.0	0.5759	0.5964	0.6170	0.6377	0.6586	0.6690
	30.0	0.5527	0.5751	0.5979	0.6207	0.6438	0.6553
	40.0	0.5164	0.5423	0.5686	0.5953	0.6221	0.6355
	50.0	0.4883	0.5172	0.5466	0.5765	0.6066	0.6218

Table 1: Results for m = 2 and n = 20.

		$R_1^2$					
ν	$\theta$	0.1	0.3	0.5	0.7	0.9	1.0
3	0.0	0.9157	0.9157	0.9157	0.9157	0.9157	0.9157
	1.0	0.8860	0.8868	0.8876	0.8884	0.8892	0.8896
	2.0	0.8601	0.8616	0.8631	0.8646	0.8661	0.8669
	4.0	0.8186	0.8213	0.8241	0.8268	0.8295	0.8308
	6.0	0.7871	0.7908	0.7945	0.7982	0.8019	0.8037
	8.0	0.7621	0.7666	0.7711	0.7757	0.7802	0.7824
	10.0	0.7416	0.7469	0.7521	0.7573	0.7625	0.7652
	15.0	0.7029	0.7096	0.7163	0.7231	0.7298	0.7332
	20.0	0.6748	0.6828	0.6908	0.6987	0.7067	0.7107
	25.0	0.6531	0.6621	0.6712	0.6802	0.6893	0.6938
	30.0	0.6354	0.6454	0.6554	0.6654	0.6755	0.6805
	40.0	0.6079	0.6195	0.6312	0.6430	0.6547	0.6606
	50.0	0.5868	0.5999	0.6131	0.6263	0.6395	0.6461
20	0.0	0.9157	0.9157	0.9157	0.9157	0.9157	0.9157
	1.0	0.8854	0.8862	0.8870	0.8878	0.8886	0.8890
	2.0	0.8573	0.8589	0.8605	0.8620	0.8636	0.8644
	4.0	0.8094	0.8124	0.8153	0.8183	0.8212	0.8227
	6.0	0.7717	0.7758	0.7798	0.7839	0.7879	0.7899
	8.0	0.7417	0.7467	0.7517	0.7567	0.7617	0.7641
	10.0	0.7175	0.7233	0.7291	0.7348	0.7406	0.7435
	15.0	0.6732	0.6806	0.6880	0.6955	0.7029	0.7066
	20.0	0.6427	0.6515	0.6602	0.6689	0.6777	0.6821
	25.0	0.6200	0.6298	0.6397	0.6496	0.6595	0.6645
	30.0	0.6020	0.6128	0.6237	0.6347	0.6456	0.6511
	40.0	0.5746	0.5873	0.6000	0.6127	0.6254	0.6318
	50.0	0.5542	0.5684	0.5826	0.5969	0.6112	0.6184

Table 2: Results for m = 2 and n = 40.

		$R_1^2$						
ν	$\theta$	0.1	0.3	0.5	0.7	0.9	1.0	
3	0.0	0.9422	0.9422	0.9422	0.9422	0.9422	0.9422	
	1.0	0.9222	0.9251	0.9280	0.9308	0.9335	0.9348	
	2.0	0.9006	0.9068	0.9129	0.9188	0.9245	0.9273	
	4.0	0.8571	0.8701	0.8828	0.8951	0.9069	0.9127	
	6.0	0.8159	0.8352	0.8542	0.8726	0.8904	0.8990	
	8.0	0.7780	0.8029	0.8276	0.8518	0.8752	0.8865	
	10.0	0.7433	0.7732	0.8031	0.8326	0.8613	0.8751	
	15.0	0.6687	0.7085	0.7494	0.7906	0.8310	0.8506	
	20.0	0.6077	0.6548	0.7044	0.7553	0.8060	0.8306	
	25.0	0.5568	0.6094	0.6659	0.7250	0.7847	0.8139	
	30.0	0.5136	0.5702	0.6323	0.6985	0.7664	0.7997	
	40.0	0.4438	0.5056	0.5760	0.6539	0.7359	0.7766	
	50.0	0.3898	0.4542	0.5303	0.6173	0.7112	0.7584	
20	0.0	0.9422	0.9422	0.9422	0.9422	0.9422	0.9422	
	1.0	0.9225	0.9254	0.9281	0.9309	0.9336	0.9349	
	2.0	0.9013	0.9073	0.9132	0.9190	0.9246	0.9273	
	4.0	0.8566	0.8697	0.8824	0.8946	0.9065	0.9122	
	6.0	0.8118	0.8320	0.8516	0.8705	0.8886	0.8974	
	8.0	0.7688	0.7956	0.8218	0.8472	0.8715	0.8832	
	10.0	0.7283	0.7612	0.7937	0.8252	0.8554	0.8700	
	15.0	0.6390	0.6848	0.7307	0.7759	0.8197	0.8407	
	20.0	0.5651	0.6206	0.6773	0.7342	0.7898	0.8166	
	25.0	0.5035	0.5663	0.6318	0.6986	0.7645	0.7965	
	30.0	0.4515	0.5197	0.5924	0.6677	0.7429	0.7795	
	40.0	0.3687	0.4438	0.5273	0.6165	0.7078	0.7525	
	50.0	0.3057	0.3843	0.4750	0.5753	0.6801	0.7319	

Table 3: Results for m = 4 and n = 20.

		$R_1^2$						
ν	$\theta$	0.1	0.3	0.5	0.7	0.9	1.0	
3	0.0	0.9374	0.9374	0.9374	0.9374	0.9374	0.9374	
	1.0	0.9234	0.9246	0.9258	0.9269	0.9281	0.9287	
	2.0	0.9091	0.9116	0.9140	0.9165	0.9189	0.9201	
	4.0	0.8811	0.8863	0.8915	0.8965	0.9014	0.9038	
	6.0	0.8552	0.8630	0.8706	0.8781	0.8855	0.8891	
	8.0	0.8314	0.8416	0.8517	0.8615	0.8711	0.8758	
	10.0	0.8096	0.8221	0.8344	0.8464	0.8582	0.8639	
	15.0	0.7625	0.7798	0.7971	0.8140	0.8306	0.8388	
	20.0	0.7232	0.7448	0.7662	0.7875	0.8083	0.8185	
	25.0	0.6897	0.7149	0.7401	0.7651	0.7897	0.8018	
	30.0	0.6606	0.6889	0.7174	0.7458	0.7739	0.7877	
	40.0	0.6117	0.6452	0.6794	0.7138	0.7480	0.7648	
	50.0	0.5717	0.6095	0.6485	0.6880	0.7275	0.7469	
20	0.0	0.9374	0.9374	0.9374	0.9374	0.9374	0.9374	
	1.0	0.9235	0.9247	0.9258	0.9270	0.9281	0.9287	
	2.0	0.9092	0.9116	0.9140	0.9164	0.9188	0.9199	
	4.0	0.8802	0.8854	0.8905	0.8955	0.9004	0.9028	
	6.0	0.8519	0.8599	0.8678	0.8754	0.8829	0.8866	
	8.0	0.8252	0.8358	0.8463	0.8566	0.8666	0.8715	
	10.0	0.8001	0.8133	0.8263	0.8391	0.8515	0.8576	
	15.0	0.7448	0.7638	0.7825	0.8009	0.8188	0.8275	
	20.0	0.6985	0.7225	0.7461	0.7693	0.7921	0.8032	
	25.0	0.6593	0.6875	0.7154	0.7430	0.7700	0.7832	
	30.0	0.6255	0.6574	0.6891	0.7206	0.7514	0.7665	
	40.0	0.5697	0.6077	0.6460	0.6841	0.7217	0.7402	
	50.0	0.5248	0.5678	0.6115	0.6554	0.6989	0.7203	

Table 4: Results for m = 4 and n = 40.