

PDF issue: 2025-06-01

Saturated humidity accelerates lateral root development in rice (Oryza sativa L.) seedlings by increasing phloem-based auxin transport

Tory, Chhun ; Uno, Yuichi ; Taketa, Shin ; Azuma, Tetsushi ; Ichii, Masahiko ; Okamoto, Takashi ; Tsurumi, Seiji

(Citation) Journal of Experimental Botany,58(7):1695-1704

(Issue Date) 2007-03-23

(Resource Type) journal article

(Version) Version of Record

(URL) https://hdl.handle.net/20.500.14094/90001060



RESEARCH PAPER

Saturated humidity accelerates lateral root development in rice (*Oryza sativa* L.) seedlings by increasing phloem-based auxin transport

Tory Chhun¹, Yuichi Uno², Shin Taketa³, Tetsushi Azuma², Masahiko Ichii³, Takashi Okamoto¹ and Seiji Tsurumi^{1,*}

¹ Center for Supports to Research and Education Activities Isotope Division, Kobe University, Kobe, 657-8501 Japan

² Faculty of Agriculture, Kobe University, Kobe, 657-8501 Japan

³ Faculty of Agriculture, Kagawa University, Miki, Kagawa, 761-0795 Japan

Received 4 October 2006; Revised 16 January 2007; Accepted 22 January 2007

Abstract

Auxin transport plays a significant role modifying plant growth and development in response to environmental signals such as light and gravity. However, the effect of humidity on auxin transport is rarely documented. It is shown here that the transport of labelled indole-3-acetic acid (IAA) from the shoot to the root is accelerated in rice (Oryza sativa L. ssp. indica cv. IR8) seedlings grown under saturated humidity (SHseedlings) compared with plants grown under normal humidity (NH-seedlings). The development of lateral roots in SH-seedlings was greatly enhanced compared with NH-seedlings. Removal of the shoot from SHseedlings reduced the density of lateral roots, and the application of IAA to the cut stem restored the lateral root density, while the decapitation of NH-seedlings did not alter lateral root development. Phloem-based auxin transport appeared responsible for enhanced lateral root formation in SH-seedlings since (i) the rate of IAA transport from the shoot to the root tip was greater than 3.5 cm h^{-1} and (ii) naphthylphthalamic acid (NPA)-induced reduction of polar auxin transport in the shoot did not influence the number of lateral roots in SH-seedlings. It is proposed that high humidity conditions accelerate the phloem-based transport of IAA from the leaf to the root, resulting in an increase in the number of lateral roots.

Key words: *AUX1*-like gene, auxin transport, humidity, lateral root development, *Oryza sativa* L. ssp. *indica* cv. IR8, phloem transport, rice seedlings.

Introduction

Auxins play a pivotal part in plant growth and development, including cell enlargement and division, lateral branching of shoots and roots, vascular differentiation, gravitropism, and early embryonic development (Davies, 1995; Hobbie, 1998). The auxin indole-3-acetic acid (IAA) is unique amongst plant hormones in being directionally transported (Lomax *et al.*, 1995). In addition to directional transport, auxin can also move through phloem (Morris and Kadir, 1972). These auxin pathways are not constant processes but change thoroughout the plant life cycle or in response to environmental stimuli such as light and gravity (Lomax *et al.*, 1995).

In shoot tissues, polar auxin transport occurs in a basipetal direction from the shoot apex to the base. In contrast, auxin transport is more complex in roots, where IAA is transported acropetally through the central cylinder and basipetally via the outer layers of root cells (Tsurumi and Ohwaki, 1978; Estelle, 1996; Marchant *et al.*, 1999; Muday, 2001; Blilou *et al.*, 2005; Swarup *et al.*, 2005). Recent knowledge about auxin transport has been primarily obtained through a series of studies on auxin influx and efflux carriers using *Arabidopsis* mutants. Auxin is taken up by plant cells via a carrier protein termed AUX1

* To whom correspondence should be addressed. E-mail: tsurumis@scitec.kobe-u.ac.jp or tsurumis@silver.kobe-u.ac.jp Abbreviations: NH, normal humidity; NPA, naphthylphthalamic acid; SH, saturated humidity.

© The Author [2007]. Published by Oxford University Press [on behalf of the Society for Experimental Biology]. All rights reserved. For Permissions, please e-mail: journals.permissions@oxfordjournals.org

(Bennett *et al.*, 1996; Swarup *et al.*, 2004; Yang *et al.*, 2006) and is mobilized out of cells via auxin efflux carriers encoded by a family of *PIN* genes (Gälweiler *et al.*, 1998; Benková *et al.*, 2003; Petrásek *et al.*, 2006) or *PGPs* (Geisler *et al.*, 2005). Mutations of *AUX1* and *AGR1/EIR1/PIN2/WAV6* (Chen *et al.*, 1998; Luschnig *et al.*, 1998; Müller *et al.*, 1998; Utsuno *et al.*, 1998; Marchant *et al.*, 1999) disrupt gravity response in roots and/or lateral root formation, demonstrating that auxin transport plays a critical role in root growth and development.

Auxin synthesized in young apical parts and leaves is transported to the roots through the phloem or a polar transport system (Reed *et al.*, 1998; Marchant *et al.*, 2002; Ljung *et al.*, 2005). However, the respective role of the two auxin pathways for root development is still unclear.

In the present study, it is shown that the saturated humidity surrounding the shoot of rice seedlings influences lateral root development by impacting phloem-based auxin transport from the shoot to the root. This represents a novel mechanism for how an environmental stimulus could modify root architecture by influencing auxin transport.

Materials and methods

Plant materials and growth condition

Seeds of rice cultivar IR8 (*Oryza sativa* L. ssp. *indica*) were surface-sterilized in a 0.2% (v/v) Benomyl (Dupont Company, Tokyo, Japan) solution for 24 h, rinsed, and soaked in water overnight in the dark at 30 °C. Germinated seeds were transferred to a floating net and grown hydroponically in a 400 ml glass cup under continuous white light at an irradiance of approximately 100 µmol $m^{-2} s^{-1}$ for 4 d at 25 °C in a growth room, where the relative humidity was kept at 60%. For saturated humidity, glass cups were placed in a closed glass chamber of 3.0 l with or without a continuous flow (1.8 l min⁻¹) of water-saturated fresh air (Fig. 1; see



Fig. 1. Growth conditions for NH-seedlings propagated at 60% relative humidity (A) versus SH-seedlings grown at 100% relative humidity in a closed glass chamber (B). Rice seedlings were grown hydroponically in a 400 ml glass cup under continuous white light at an irradiance of approximately 100 μ mol m⁻² s⁻¹ for 4 d at 25 °C in a room, where the relative humidity was maintained at 60%.

Supplementary Fig. S1 at *JXB* online). Relative humidity was measured with a digital humidity meter (HN-K, Chino Corp)

Transport of labelled IAA from the shoot to the root

Auxin transport from the shoot to the root was performed mainly as described by Chhun et al. (2004) with a slight modification. The shoots of 4-d-old seedlings were decapitated and the cut stem of 0.5 cm in length was capped with a small plastic tip containing 3 μ l of 10 mM MES buffer (pH 5.7) supplemented with 1 µM 5-[³H]IAA (specific activity 740 MBq μ mol⁻¹, American Radiolabelled Chemicals, Inc., St Louis, MO, USA). The [³H]IAA-treated seedlings were incubated for various transport periods under 60% or 100% relative humidity. Radioactivity was measured either on the whole root or after dividing the root into three or four segments. The length of root segments was approximately 1 cm while the length of the root tip segment was slightly longer than 1 cm depending on the length of the whole root. Ten whole roots or root segments were combined, and were placed into 5 ml of scintillation fluid overnight. Radioactivity was counted with a scintillation counter (model LS6500, Beckman Instruments, Fullerton, CA, USA).

Basipetal transport of labelled IAA in root segments

Basipetal auxin transport in roots was measured as described by Chhun *et al.* (2005) with a slight modification. An apical root segment of 1.6 cm in length was obtained from a 4-d-old seedling and placed vertically on a small filter paper containing 1 μ M [³H]IAA in a 1.5 ml Eppendorf plastic tube after decapitating the apical 1 mm in order to keep the contact of the apical end of the root segment with the donor filter paper. After 1 h incubation at room temperature, a 3 mm segment was cut from the basal end of the root segment. Five 3 mm root segments were combined and the radioactivity counted.

Application of auxin and NPA to shoot

Two-day-old SH-seedlings were decapitated by removing the apex to retain 0.3 cm of shoot tissue. The decapitated shoot was capped with a small plastic tube containing 5 μ l of 0 μ M and 10 μ M IAA. IAA-treated seedlings were grown under 100% relative humidity for an additional 2 d and the number of lateral roots was measured.

For *N*-(1-naphthyl)phthalamic acid (NPA, Tokyo Kasei Co., Tokyo, Japan) treatment, NPA was applied to the basal part of the shoot as a ring after mixing with lanolin.

Cloning of four AUX1-like genes of rice (cv. IR8)

By searching the DNA Data Bank of Japan database (DDBJ), four AUX1-like genes were found (GenBank accession numbers AK100090, AK103524, AK111849, and AK102295) in rice (ssp. japonica). Specific primers for the AUX1-like genes were designed on the basis of DNA sequences in the DDBJ database (see Supplementary Table S2 at JXB online). Messenger RNA (mRNA) was extracted from roots of 7-d-old seedlings of rice cv. IR8 according to the manufacturer's instruction (QuickPrep Micro mRNA Purification Kit, Amersham Biosciences Ltd, UK). Reverse transcription was then performed with reverse transcriptase (Power-Script RT, BD Biosciences, Palo Alto, CA, USA) in a total volume of 20 µl at 42 °C for 90 min. PCR amplification was performed with 1 µg of cDNA essentially based on the manufacturer manual (Ex Taq, TaKaRa, Shiga, Japan). Four cDNA fragments were obtained, termed OsRAU1 (related to AUX1), OsRAU2, OsRAU3, and OsRAU4, which were cloned into the pCR2.1 TOPO vector using the TOPO TA cloning Kit (Invitrogen, Carlsbad, CA, USA) as described in the instruction manual. Positive identification was performed by the dye-primer cycle sequencing method using a BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA) and a DNA sequence analyser (ABI PRISM-Avant Genetic Analyzer, Applied Biosystems). Sequence comparisons with databases were performed at the US NCBIs GenInfo Network BLAST server.

Gene expression analysis

Total RNA was extracted from various rice organs including the shoot, embryo, upper and lower seminal roots, lateral roots, and crown roots using the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany). Reverse transcription was then performed in a total volume of 20 μ l using 5 μ g of DNase-treated total RNA as template at 42 °C for 90 min. PCR amplification was performed with 1 μ g of cDNA essentially based on the manufacturer's manual (Ex Taq, TaKaRa, Shiga, Japan). For gene expression analysis, specific primers for *OsRAU1*, *OsRAU2*, *OsRAU3*, and *OsRAU4* genes were designed (see Supplementary Table S2 at *JXB* online) to generate 300 bp sized products while the rice actin gene was used as a positive control. RT-PCR conditions were carefully manipulated to measure transcripts at the exponential phase of amplification. The PCR products in the exponential range of amplification were separated on an 1.2% agarose gel and stained with ethidium bromide.

Results

Rice seedling root morphology is modified in response to saturated humidity

Rice seedlings were grown hydroponically under 60% relative humidity in the light at 25 °C for 4 d (termed NH-seedlings). Seedlings were also grown under saturated humidity in a closed glass chamber (termed SH-seedlings) (Fig. 1). As shown in Table 1 and Fig. 2, it was observed that the root morphologies of SH-seedlings were distinct from those of NH-seedlings. The number of lateral roots per plant and the density of lateral roots were greatly increased in SH-seedlings. Furthermore, the length of the lateral roots was longer in SH-seedlings than in NH-seedlings and

Table 1. Characteristics of NH- and SH-seedlings grown under normal and saturated humidity, respectively

Rice seedlings were grown hydroponically in a 400 ml glass cup under 60% relative humidity (NH-seedlings) or saturated humidity (SH-seedlings) in continuous white light at 25 °C for 4 d (Fig. 1). Lateral roots emerged at least 0.5 mm from the root surface were counted. Data represent averages (\pm SE) of 30 seedlings from three independent experiments. *, ** indicate significant differences from NH-seedlings at the 5% and 1% levels, respectively, as judged by the Student's *t* test.

Characters	NH-seedlings	SH-seedlings
Lateral roots (plant ⁻¹)	10.00±1.20	18.60±1.50**
Lateral roots $(cm^{-1})^a$	2.86 ± 0.30	5.09±0.20**
Lateral root length $(cm)^{b}$	0.26 ± 0.02	$0.60 \pm 0.04 **$
Root length (cm)	4.05 ± 0.10	3.67±0.20*
Root tip angle $(degree)^c$	5.70 ± 0.90	31.15±5.20**
Plant height (cm)	1.42 ± 0.08	2.91±0.10**

^{*a*} Density of the lateral roots was calculated by dividing the number of lateral roots by the length of the seminal root.

^b The average of the five longest lateral roots was determined for each plant and then the average for 10 plants was obtained.

^c The direction of gravity is shown as 0 degrees.

dense crown roots appeared on the basal part of shoots in SH-seedlings (Fig. 2), whereas the length of seminal roots of SH-seedlings was shorter compared with NH-seedlings. The direction of SH-root tip growth deviated from the gravity vector, suggesting that the response to gravity was altered in the SH-roots, whereas NH-roots grew downward in response to gravity.

Since the SH-seedlings were grown in a closed chamber for 4 d, it was possible that the seedlings could be influenced by some gaseous factors accumulated in the chamber. To rule out this possibility, the air inside the chamber was refreshed with a continuous flow of watersaturated air (see Supplementary Fig. S1 and Table S1 at *JXB* online). It was observed that this did not alter SHroot architecture, suggesting that saturated humidity represents the cause of the root morphology of SHseedlings including enhanced lateral root development and root tip curving.

Excision of SH-seedlings shoot tissue blocks humidity-induced changes in root architectures

Lateral root development in *Arabidopsis* is controlled by auxin derived from the shoot (Reed *et al.*, 1998; Bhalerao *et al.*, 2002). To test this possibility in this experimental system, the effect of removing the shoot apex on the lateral root development in rice seedlings was examined. The shoot of 2-d-old seedlings was removed and root morphology was observed 2 d after decapitation (Fig. 3A). The removal of the shoot apex of SH-seedlings reduced the density of lateral roots compared with intact seedlings and, unexpectedly, the roots grew downward as if the gravity response was restored (Fig. 3B). In contrast, the removal of the shoot apex of NH-seedlings did not alter lateral root density and root gravity response as compared with intact seedlings (Fig. 3).



Fig. 2. Four-day-old seedlings grown under 60% relative humidity (A, NH-seedlings) and 100% relative humidity (B, SH-seedlings). SH-seedlings display dense and longer lateral roots, enhanced shoot height and root curling compared with NH-seedlings.

Application of auxin to the decapitated SH-seedlings restores lateral root density

The effect of removing the shoot apex on lateral root development could be due to a reduction in auxin supply from the shoot. To test this idea IAA was applied to the decapitated shoot of 2-d-old SH-seedlings and lateral root density was measured 2 d after IAA application. As predicted by our model, application of 10 μ M IAA increased the density of lateral roots in the SH-seedlings to a level equivalent to intact seedlings (Fig. 4). These results raise the possibility that saturated humidity increases auxin transport from the shoot to the root, and that decapitation of the shoot nullifies the auxin supply from this source.

The transport of labelled IAA from shoot to root is enhanced in SH-seedlings

To test the idea that saturated humidity alters auxin transport, the transport of labelled IAA from shoot to root



tissues was examined in 4-d-old seedlings. As shown in Fig. 5, a greater amount of $[{}^{3}H]IAA$ was transported from the shoot to the root tissues in SH-seedlings compared with NH-seedlings. The difference was significant at 1 h and became greater after 2 h incubation, implying that saturated humidity accelerated the auxin transport from the shoot to the root.

To analyse the distribution of labelled IAA in roots, radioactivity was measured in each 1 cm root segment.



Fig. 4. Effect of IAA on lateral root density when IAA was applied to the decapitated stem of SH-seedlings. The shoot apex of 2-d-old SH-seedlings was removed before mature lateral root formation. The decapitated stem was then capped with a small tube containing 5 μ l of 0 μ M or 10 μ M IAA. Lateral root density was measured following 2 d incubation. Data are averaged (±SE) for 30 seedlings. b is significantly different from a at *P* <0.01 as judged by the Student *t* test.



Fig. 3. The effect of removing the shoot apex in NH- and SH-seedlings on lateral root density (A) and root gravity response (B). The shoots of 2-d-old NH- and SH-seedlings were removed and the root morphology was observed 2 d after decapitation. Shoot decapitation was done two or three times a day by excising the shoot apex with a razor blade to avoid new growth. The direction of gravity is shown as 0 degree. Data are averaged (\pm SE) for 30 seedlings. Two asterisks represent significant differences from intact seedlings at *P* <0.01 as judged by the Student *t* test.

Fig. 5. Time-course of [³H]IAA movement from shoot to root in NHand SH-seedlings. [³H]IAA was applied to decapitated shoots of 4-d-old NH- and SH-seedlings under NH- and SH-condition, respectively, and the radioactivity of whole root was measured after various transport periods. Data are averaged (\pm SE) from at least 30 roots. One asterisk and two asterisks represent significant differences from NH-seedlings at *P* <0.05 and *P* <0.01, respectively, as judged by the Student *t* test.

After various transport periods, roots were divided into four segments for NH-seedlings and into three segments for SH-seedlings. As shown in Fig. 6, labelled auxin reached the apical root segment in both seedling treatments after 1 h incubation. Since the distance from the application site of the shoot to the apical root segment in NH-seedlings was about 3.5 cm, the velocity of labelled IAA from the shoot to the root tip is estimated to be 3.5 cm h⁻¹ or greater. Such rapid transport of IAA is characteristic of phloem (rather than polar) transport (Goldsmith *et al.*, 1974).

In NH-roots (Fig. 6), the radioactivity per segment decreased from segment 1 to segment 3 and this pattern did not change for 4 h. In contrast, in SH-seedlings, the radioactivity of segment 2 increased greatly after 2 h incubation and reached the same level as segment 1 at 3 h. Segment 2 of SH-roots represents the position where lateral roots emerge under SH-conditions (Fig. 2). In NH-seedlings, lateral roots were formed only in segment 1, whereas in SH-seedlings they emerged in a wider region including segments 1 and 2.

NPA does not block the humidity-induced increase in lateral root number and density in SH-seedlings

It could be predicted that if auxin transport from the shoot to the root were mediated by a polar transport system, the auxin transport inhibitor NPA would block auxin movement. To address this issue, the transport of labelled auxin from the shoot to the root was examined after treating the basal part of shoots in 4-d-old SH-seedlings with NPA lanolin. As shown in Fig. 7A, NPA slightly but signifi-



Fig. 6. Distribution of radioactivity in roots when $[{}^{3}H]IAA$ was applied to the shoots of NH- (A) and SH- (B) seedlings. $[{}^{3}H]IAA$ was applied to decapitated shoots of 4-d-old NH- and SH-seedlings under NH- and SH-condition, respectively. After various transport periods, the root was divided into four (A) or three (B) segments and the radioactivity of each segment was measured. The length of each root segment was l cm except at the root tip. The length of the tip segment was slightly longer than 1 cm depending on the length of whole root. Data represent averages (\pm SE) of 30 roots from three independent experiments.

cantly reduced [³H]IAA movement from the shoot to the root by 15%, implying that only a small proportion of auxin transport from shoot to root is mediated by a polar transport system.

To evaluate the role of polar auxin transport in shoot tissues in root morphology, the effect of NPA was examined on lateral root formation. The basal part of the shoot of 2-d-old seedlings was treated with NPA lanolin in a ring



Fig. 7. Effect of an auxin transport inhibitor, NPA, on [³H]IAA transport from shoot to root (A) and lateral root number and density (B). The NPA solution was mixed with lanolin and applied to the basal part of the shoot as a ring. (A) [³H]IAA was applied to the decapitated shoot of 4-d-old SH-seedlings, which were treated with NPA, and the radioactivity of the whole root was measured after a 4 h transport period. (B) Two-day-old SH-seedlings, treated with NPA, were grown for a further 2 d and the number of lateral roots was measured. To confirm the effect of NPA, the inhibitor was re-applied every day to the basal part of the shoot. Data represent averages (\pm SE) of 40 seedlings from four independent experiments. The asterisk represents significant differences from 0 μ M-NPA treated seedlings at *P* <0.05 as judged by the Student *t* test.

and the number of lateral roots was counted 2 d later. To ensure the effectiveness of NPA treatment, the NPA lanolin ring was applied every day to the basal part of the shoot. Surprisingly, the number or density of lateral roots per plant was not reduced in NPA-treated seedlings (Fig. 7B). These results suggest that the phloem-based transport of auxin is responsible for humidity-induced lateral root development. However, it is still possible that some signal from the shoot, other than auxin, is acting in this system.

Basipetal transport of IAA in roots is reduced in SH-seedlings compared with NH-seedlings

SH-seedlings exhibit altered gravitropic growth (Fig. 2). Since the basipetal transport of auxin in the roots (from the root tip backward to the base) is the important factor for gravitropism (Rashotte *et al.*, 2000; Swarup *et al.*, 2005), the basipetal transport of labelled IAA was measured in the roots of 4-d-old SH- and NH-seedlings (Fig. 8). The basipetal [³H]IAA transport in the roots was suppressed in SH-seedlings compared with NH-seedlings, consistent with the reduced gravitropic response of SH-seedling root tips.

Sequence and expression analysis of four AUX1-like genes in rice (ssp. indica cv. IR8)

AUX1 gene expression in Arabidopsis leaves has been proposed to regulate the phloem-based IAA transport by facilitating hormone loading into the leaf vascular system (Marchant *et al.*, 2002). In order to examine the possible involvement of AUX1-like genes in the phloem-based IAA transport from the shoot to the root in rice, cloning and expression analysis of homologous gene were carried out. By searching the DDBJ database, four AUX1-like genes were found in rice (ssp. *japonica* cv. Nipponbare); AK100090 on chromosome 1, AK103524 on chromosome 3, AK111849 on chromosome 5, and AK102295 on chromosome 10. These genes shared a high degree of identity at the amino acid level with AtAUX1; 83.7, 77.5, 82.5, and 76.6%, respectively. Primer sets were designed to clone the whole open reading frame of each gene in rice (ssp. indica cv. IR8) by RT-PCR. The deduced amino acid sequences of the four obtained cDNAs were aligned with AtAUX1 (see Supplementary Fig. S2 at JXB online). The difference in sequence between IR8 and Nipponbare was only three or four bases per gene and one or two amino acids per protein as for the AUX1-like genes. Amino acids, which were substituted in the 13 aux1 missense alleles (Swarup et al., 2004), are conserved in four homologous genes in rice (see Supplementary Fig. S2 at JXB online). Based on their sequence identity, these cDNAs were termed OsRAU1 (GenBank accession number AB275160), OsRAU2 (AB275161), OsRAU3 (AB275162), and OsRAU4 (AB275163).

To examine the role of AUX1-like genes of rice, the expression of OsRAU1, OsRAU2, OsRAU3, and OsRAU4 genes was monitored in various organs of rice seedlings by performing RT-PCR to estimate their transcript levels. As shown in Fig. 9, the OsRAU1 transcript was observed in all organs tested and particularly high levels of the OsRAU1 transcript accumulated in the upper root and lateral root samples. The OsRAU2 transcript was observed in shoots, the upper root, lower root, and lateral root samples. Transcripts were present at low to moderate levels in upper and lower roots, lateral roots, and crown root samples for OsRAU3, and in the shoot and upper roots for OsRAU4.



Fig. 8. The basipetal transport of $[{}^{3}\text{H}]\text{IAA}$ in NH- and SH-roots. $[{}^{3}\text{H}]\text{IAA}$ was applied to the apical end of 1.5 cm root segments for 1 h and the radioactivity accumulating at the basal 3 mm end of the segment was measured. Data represent averages (\pm SE) of 40 seedlings from four independent experiments. Two asterisks represent significant differences from NH-seedlings at *P* <0.05 as judged by the Student *t* test.



Fig. 9. Expression of *AUX1*-like genes, *OsRAU1*, *OsRAU2*, *OsRAU3*, and *OsRAU4*, in various organs of rice (cv. IR8). Semi-quantitative RT-PCR analysis was performed and the *actin* gene was used as control. One μ g of cDNA was used for PCR amplification. Total RNA was extracted from various organs. The embryo was obtained from seeds before benomyl treatment. Shoot and basal and apical seminal root samples were prepared from 3-d-old seedlings. The apical root was taken from the apical 1 cm root tip and the basal root includes the remaining part of the root. Lateral and crown roots were prepared from 5-d-old seedlings.

The expression of the *AUX1*-like genes in 4-d-old SH-seedlings was then compared with that in NH-seedlings. Interestingly, SH-seedlings exhibited a greater abundance of the *OsRAU1* transcript in shoot tissues compared with NH-seedlings (Fig. 10). In contrast to *OsRAU1*, the expression levels of *OsRAU2*, *OsRAU3*, and *OsRAU4* in SH-seedlings were the same as those in NH-seedlings.

Discussion

It is reported here that the shoot and root morphologies of rice seedlings grown under saturated humidity are distinct from those under normal humidity (Fig. 2). The shoot height of SH-seedlings also increased compared with NHseedlings (Table 1), consistent with a previous observation that shoot elongation of floating rice was enhanced under SH-conditions (Azuma et al., 1991). Although ethylene and gibberellins promoted internodal elongation of rice seedlings, Azuma et al. (1991) found that a moist environment was required for ethylene-induced internodal elongation in floating rice while the stimulatory effect of gibberellins was independent of moisture conditions. Ethylene's requirement for high humidity conditions suggests that this signal is involved in promoting the shoot height of SH-seedlings. In the present study, it has been shown that the SH-seedlings exhibit altered root morphology, including dense root branching and root agravitropism (Fig. 2; Table 1), despite the roots having been kept in water, implying that the highly humid air surrounding the shoot influences root development. Ethylene is not likely to cause the humidity-induced changes in root morphology because the SH-induced increase in lateral root density was also observed, even when the air surrounding the shoot was continuously refreshed with water-saturated air (see Supplementary Table S1 at JXB online).



Fig. 10. Comparison of transcription levels of AUXI-like genes, OsRAUI, OsRAU2, OsRAU3, and OsRAU4, in shoot and root tissues of 4-d-old NH- and SH-seedlings. Semi-quantitative RT-PCR analysis was performed and the *actin* gene was used as control. One μ g of cDNA was used for PCR amplification.

It is proposed that the SH-induced increase in lateral root density is due to an enhanced translocation of auxin from the shoot to the root tissue as discussed below. First, removal of the shoot apex of SH-seedlings greatly reduced lateral root density in the seminal root (Fig. 3A), indicating that the humidity-induced changes in root morphology are dependent on the presence of shoot tissues. In contrast, removal of shoot tissues did not alter lateral root density in NH-seedlings. Secondly, application of IAA to the decapitated shoot of SH-seedlings restored the density of lateral roots to the intact SH-seedling level. These results indicate that a supply of auxin from shoot to root tissues is required for the humidity-induced increase in lateral root density. It is interesting to note that the requirement of the shoot for lateral root development was only observed in SH-roots (and not NH-roots). This difference in the role of the shoot raises the possibility that the auxin supply from shoot to root tissues is enhanced in SH-seedlings. This idea was supported by the finding that the transport of labelled IAA from the shoot to the root was greater in SHseedlings compared with NH-seedlings (Fig. 5). Furthermore, the elevated level of transport of labelled IAA from the shoot to the root tissues in SH-seedlings resulted in an enhanced accumulation of radioactivity in the basal part of the roots (Fig. 6), just where the lateral roots emerged under SH-conditions (Fig. 2). These results are consistent with the idea that SH-induced lateral root formation is ascribed to the accelerated transport of auxin from the shoot to the root.

Auxin transport from shoot to root tissues could occur either through phloem or through polar transport routes. Polar transport of auxin occurs at a velocity of about 10 mm h^{-1} in the roots of *Phaseolus vulgaris*, *Pisum* sativum, and Arabidopsis thaliana (Iversen et al., 1971; Rowntree and Morris, 1979; Rashotte et al., 2003), whereas the rate of phloem-mediated auxin transport is over 5 cm h^{-1} in the roots of *Populus tremula* and *Vicia* faba (Eliasson, 1972; Tsurumi and Wada, 1980). Measuring the rate of auxin transport is one of the useful ways to discriminate between these transport routes (Goldsmith et al., 1974). When labelled IAA was applied to the shoot, in these experiments, radioactivity was detectable in the root apex within the first hour of transport in either NH- or SH-seedlings (Fig. 6). The longer the transport period, the greater the level of radioactivity that accumulated in the root apex. Based on the distance of translocation from the decapitated shoot to the root tip and the detection of radioactivity in the root apex within 1 h (Fig. 6), the velocity of auxin translocation from the shoot to the root tip was estimated as 3.5 cm h^{-1} or greater, consistent with the idea that auxin transport from the shoot to the root was mediated by the phloem.

Treatment with NPA, an inhibitor of polar auxin transport, represents another method to distinguish the

transport pathway (Lomax et al., 1995). Application of NPA to the shoot base reduced the transport of labelled IAA from the shoot to the root by 15% compared with its control (Fig. 7A), indicating that polar transport represents only a minor route for the translocation of auxin from the shoot to the root. Consistent with this conclusion, similar treatment with NPA in 2-d-old seedlings did not impair the humidity-induced increase in lateral root number and density (Fig. 7B). In contrast to NPA treatment of shoot tissues, NPA application to the root has been shown to reduce lateral root numbers (Reed et al., 1998). It remains possible that a proportion of the phloem auxin in the shoot may move into the polar transport system in roots. Indeed, the migration of auxin from the phloem to polar systems has been observed in aerial tissues (Cambridge and Morris, 1996).

It is interesting to note that the transcript abundance of an *AUX1*-like gene, *OsRAU1*, was elevated in the shoot of SH-seedlings compared with NH-seedlings, whereas, in roots, the *OsRAU1* level was unchanged (Fig. 10). The amino acid sequence of OsRAU1 protein is 83.7% identical to AtAUX1 which has been proposed to play a role in phloem-based IAA transport pathway (Parry et al., 2001*b*; Swarup *et al.*, 2001; Marchant *et al.*, 2002). On the other hand, AtAUX1 does not seem to be involved in the polar auxin transport system in shoots (Parry *et al.*, 2001*a*). Further experiments are required to prove the involvement of OsRAU1 in the enhanced phloem translocation of labelled auxin.

Labelled IAA applied to shoot tissues accumulated in the basal part of roots in SH-seedlings (Fig. 6), consistent with the previous observation reporting the accumulation of labelled auxin in lateral roots and their primordia (Rowntree and Morris, 1979; Tsurumi and Wada, 1980; Kerk and Feldman, 1995). The accumulation of labelled IAA in primordia may be due to the expression of the AUX1 protein in primordia (Marchant *et al.*, 2002). Although basipetal IAA transport in root tips was reduced in SH-seedlings (Fig. 8), auxin accumulation in the basal part of roots in Fig. 6B is not likely to be due to transport inhibition, because this would result in a reduction in lateral root numbers (Reed *et al.*, 1998).

In SH-seedlings, the direction of the root tip deviated from the gravity vector (Fig. 2). This pattern of root growth was not observed in NH-seedlings. It was found that the basipetal transport of labelled IAA was greatly reduced in SH-roots compared with NH-roots (Fig. 8), suggesting that reduced basipetal IAA transport was the cause of the gravitropic defect. Although the removal of the shoot apex of SH-seedlings reduced root curvature and restored normal downward root growth as for NHroots (Fig. 3B), shoot auxin is not likely to be the cause of root curving because the application of auxin to the cut stem did not induce such curving in roots (data not shown). In contrast to SH-seedlings, the removal of the shoot apex of NH-seedlings did not alter lateral root density compared with intact seedlings (Fig. 3). The role of shoot auxin on lateral root formation is dependent on plant age (Bhalerao *et al.*, 2002). In young NH-seedlings, seed-derived auxin (Hall, 1980) may be sufficient for lateral root development. The removal of rice seedlings from their seed severely reduced lateral root number and led to their death 10 d after germination (data not shown). This result is consistent with a previous report that radioactivity from [¹⁴C]IAA applied to the seed of *Vicia faba* seedlings accumulated in lateral root primordia along with vascular bundles (Tsurumi and Wada, 1980).

In the present study, the shoot environment is the cause of SH-induced root morphology because the roots of both SH- and NH-seedlings had been kept in water. The seminal root of SH-seedlings is greatly branched and the root tip is curved (Fig. 2). Decapitation of the shoot in SH-seedlings restored the gravitropic response of the seminal root tip as well as normalizing root branching (Fig. 3), implying that the shoot surrounded by moist air is the source of signals responsible for the observed changes in root morphogenesis. The phloem-based transport of auxin from shoot to root tissues is likely to promote root branching. Auxin functions as a mediator in phototropism and gravitropism by moving from sensor tissues to responding tissues in the shoot and the roots (Swarup et al., 2005; Whippo and Hangarter, 2006). In the present experiments, the shoot represents the sensing organ for the humid environment and the root functions as the responding organ. The transmitted signal is also likely to be auxin, which is delivered via the phloem from shoot to root tissues. This represents a novel example of how an environmental signal can cause morphological changes by altering auxin transport from shoot to root tissues in rice seedlings.

Supplementary data

Supplementary data in the form of two figures and two tables can be found at *JXB* online.

Fig. S1. Schematic illustrations of growth condition for NH-seedlings grown under 60% relative humidity and SH-seedlings grown under 100% relative humidity.

Fig. S2. Predicted amino acid sequence of *AUX1*-like genes in rice (ssp. *indica* cv. IR8).

Table S1. Effect of refreshing the shoot environment on root morphology of SH-seedlings.

Table S2. List of primers for isolation and expression of *AUX1*-like genes.

Acknowledgements

We thank Dr Malcolm J Bennett of Nottingham University (Nottingham, UK) for critical reading of this manuscript, Dr Keiko Kosuge of Kobe University for DNA sequence analysis, and Dr Tetsushi Iwasaki of Kobe University for his valuable suggestions. This work was supported in part by Grants-in-Aid for Scientific Research 1604177 from the Ministry of Education, Culture, Sports, Science and Technology of Japan to ST.

References

- Azuma T, Mihara F, Uchida N, Yasuda T, Yamaguchi T. 1991. Influence of humidity on ethylene-induced internodal elongation in floating rice. *Plant and Cell Physiology* **32**, 307–309.
- Benková E, Michniewicz M, Sauer M, Teichmann T, Seifertová D, Jürgens G, Friml J. 2003. Local, efflux-dependent auxin gradients as a common module for plant organ formation. *Cell* **115**, 591–602.
- Bennett MJ, Marchant A, Green HG, May ST, Ward SP, Millner PA, Walker AR, Schulz B, Feldmann KA. 1996. Arabidopsis *AUX1* gene: a permease-like regulator of root gravitropism. *Science* 273, 948–950.
- Bhalerao RP, Eklöf J, Ljung K, Marchant A, Bennett M, Sandberg G. 2002. Shoot-derived auxin is essential for early lateral root emergence in *Arabidopsis* seedlings. *The Plant Journal* 29, 325–332.
- Blilou I, Xu J, Wildwater M, Willemsen V, Paponov I, Friml J, Heidstra R, Aida M, Palme K, Scheres B. 2005. The PIN auxin efflux facilitator network controls growth and patterning in *Arabidopsis* roots. *Nature* **433**, 39–44.
- Cambridge AP, Morris DA. 1996. Transfer of exogenous auxin from the phloem to the polar auxin transport pathway in pea (*Pisum sativum* L.). *Planta* **199**, 583–588.
- Chen R, Hilson P, Sedbrook J, Rosen E, Caspar T, Masson PH. 1998. The Arabidopsis thaliana AGRAVITROPIC 1 gene encodes a component of the polar-auxin-transport efflux carrier. Proceedings of the National Academy of Sciences, USA 95, 15112–15117.
- Chhun T, Taketa S, Ichii M, Tsurumi S. 2005. Involvement of *ARM2* in the uptake of indole-3-butyric acid in rice (*Oryza sativa* L.). *Plant and Cell Physiology* **46**, 1161–1164.
- Chhun T, Taketa S, Tsurumi S, Ichii M. 2004. Different behaviour of indole-3-acetic acid and indole-3-butylric acid in stimulating lateral root development in rice (*Oryza sativa* L.). *Plant Growth Regulation* **43**, 135–143.
- **Davies PJ.** 1995. The plant hormones: their nature, occurrence and functions. In: Davies PJ, ed. *Plant hormones: physiology, biochemistry and molecular biology*, 2nd edn. Dordrecht, The Netherlands: Kluwer Academic Publishers, 1–12.
- Eliasson L. 1972. Translocation of shoot-applied indolylacetic acid into the roots of *Populus tremula*. *Physiologia Plantarum* 27, 412–416.
- Estelle M. 1996. Plant tropism: the ins and outs of auxin. *Current Biology* 6, 1589–1591.
- Gälweiler L, Guan C, Müller A, Wisman E, Mendgen K, Yephremov A, Palme K. 1998. Regulation of polar auxin transport by AtPIN1 in *Arabidopsis* vascular tissue. *Science* 282, 2226–2229.
- Geisler M, Blakeslee JJ, Bouchard R, *et al.* 2005. Cellular efflux of auxin catalyzed by the Arabidopsis MDR/PGP transporter AtPGP1. *The Plant Journal* **44**, 179–194.
- **Goldsmith MHM, Cataldo DA, Karn J, Brenneman T, Trip P.** 1974. The rapid non-polar transport of auxin in the phloem of intact *Coleus* plants. *Planta* **116**, 301–317.
- Hall PJ. 1980. Indole-3-acetyl-*myo*-inositol in kernels of *Oryza* sativa. Phytochemistry 19, 2121–2123.
- Hobbie LJ. 1998. Auxin: molecular genetic approaches in *Arabidopsis. Plant Physiology and Biochemistry* **36**, 91–102.
- **Iversen T-H, Aasheim T, Pedersen K.** 1971. Transport and degradation of auxin in relation to geotropism in roots of *Phaseolus vulgaris. Physiologia Plantarum* **25**, 417–424.

- Kerk NM, Feldman LJ. 1995. A biochemical model for the initiation and maintenance of the quiescent center: implications for organization of root meristems. *Development* 121, 2825–2833.
- Ljung K, Hull AK, Celenza J, Yamada M, Estelle M, Normanly J, Sandberg G. 2005. Sites and regulation of auxin biosynthesis in *Arabidopsis* roots. *The Plant Cell* 17, 1090–1104.
- Lomax TL, Muday GK, Rubery PH. 1995. Auxin transport. In: Davies PJ, ed. *Plant hormones; physiology, biochemistry and molecular biology*, 2nd edn. Dordrecht, The Netherlands: Kluwer Academic Publishers, 509–530.
- Luschnig C, Gaxiola RA, Grisafi P, Fink GR. 1998. EIR1, a root-specific protein involved in auxin transport, is required for gravitropism in *Arabidopsis thaliana*. *Genes and Development* **12**, 2175–2187.
- Marchant A, Bhalerao R, Casimiro I, Eklöf J, Casero PJ, Bennett M, Sandberg G. 2002. AUX1 promotes lateral root formation by facilitating indole-3-acetic acid distribution between sink and source tissues in the *Arabidopsis* seedling. *The Plant Cell* 14, 589–597.
- Marchant A, Kargul J, May ST, Muller P, Delbarre A, Perrot-Rechenmann C, Bennett MJ. 1999. AUX1 regulates root gravitropism in *Arabidopsis* by facilitating auxin uptake within root apical tissues. *EMBO Journal* **18**, 2066–2073.
- Morris DA, Kadir GO. 1972. Pathways of auxin transport in the intact pea seedling (*Pisum sativum* L.). *Planta* **107**, 171–182.
- Muday GK. 2001. Auxins and tropism. Journal of Plant Growth Regulation 20, 226–243.
- Müller A, Guan C, Gälweiler L, Tänzler P, Huijser P, Marchant A, Parry G, Bennett M, Wisman E, Palme K. 1998. *AtPIN2* defines a locus of *Arabidopsis* for root gravitropism control. *EMBO Journal* **17**, 6903–6911.
- Parry G, Delbarre A, Marchant A, Swarup R, Napier R, Perrot-Rechenmann C, Bennett MJ. 2001a. Novel auxin transport inhibitors phenocopy the auxin influx carrier mutation aux1. The Plant Journal 25, 399–406.
- Parry G, Marchant A, May S, et al. 2001b. Quick on the uptake: characterization of a family of plant auxin influx carriers. *Journal* of Plant Growth Regulation 20, 217–225.
- Petrásek J, Mravec J, Bouchard R, *et al.* 2006. PIN proteins perform a rate-limiting function in cellular auxin efflux. *Science* **312**, 914–918.
- Rashotte AM, Brady SR, Reed RC, Ante SJ, Muday GK. 2000. Basipetal auxin transport is required for gravitropism in roots of *Arabidopsis. Plant Physiology* **122**, 481–490.
- Rashotte AM, Poutpart J, Waddell CS, Muday GK. 2003. Transport of the two natural auxins, indole-3-butyric acid and indole-3-acetic acid, in *Arabidopsis*. *Plant Physiology* **133**, 761–772.
- Reed RC, Brady SR, Muday GK. 1998. Inhibition of auxin movement from the shoot into the root inhibits lateral root development in *Arabidopsis*. *Plant Physiology* **118**, 1369–1378.
- **Rowntree RA, Morris DA.** 1979. Accumulation of ¹⁴C from exogenous labelled auxin in lateral root primordia of intact pea seedlings (*Pisum sativum* L.). *Planta* **144**, 463–466.
- Swarup R, Friml J, Marchant A, Ljung K, Sandberg G, Palme K, Bennett M. 2001. Localization of the auxin permease AUX1 suggests two functionally distinct hormone transport pathways operate in the *Arabidopsis* root apex. *Genes and Devolopment* 15, 2648–2653.
- Swarup R, Kargul J, Marchant A, *et al.* 2004. Structure–function analysis of the presumptive *Arabidopsis* auxin permease AUX1. *The Plant Cell* **16**, 3069–3083.
- Swarup R, Kramer EM, Perry P, Knox K, Leyser HMO, Haseloff J, Beemster GTS, Bhalerao R, Bennett MJ. 2005.

Root gravitropism requires lateral root cap and epidermal cells for transport and response to a mobile auxin signal. Nature Cell Biology 7, 1057–1065.

- **Tsurumi S, Ohwaki Y.** 1978. Transport of ¹⁴C-labeled indoleacetic acid in *Vicia* root segments. *Plant and Cell Physiology* **19**, 1195– 1206.
- Tsurumi S, Wada S. 1980. Transport of shoot- and cotyledonapplied indole-3-acetic acid to Vicia faba root. Plant and Cell Physiology 21, 803-816.
- Utsuno K, Shikanai T, Yamada Y, Hashimoto T. 1998. AGR, an Agravitropic locus of Arabidopsis thaliana, encodes a novel membrane-protein family member. Plant and Cell Physiology 39, 1111-1118.
- Whippo CW, Hangarter RP. 2006. Phototropism: bending towards enlightenment. The Plant Cell 18, 1110-1119.
- Yang Y, Hammes UZ, Taylor CG, Schachtman DP, Nielsen E. 2006. High-affinity auxin transport by the AUX1 influx carrier protein. Current Biology 16, 1123–1127.

JXB