



Global attractors for doubly nonlinear evolution equations with non-monotone perturbations

Akagi, Goro

(Citation)

Journal of Differential Equations, 250(4):1850-1875

(Issue Date)

2011-02-15

(Resource Type)

journal article

(Version)

Accepted Manuscript

(URL)

<https://hdl.handle.net/20.500.14094/90001966>



Global attractors for doubly nonlinear evolution equations with non-monotone perturbations

GORO AKAGI*

Abstract

This paper addresses the analysis of dynamical systems generated by doubly nonlinear evolution equations governed by subdifferential operators with non-monotone perturbations in a reflexive Banach space setting. In order to construct global attractors, an approach based on the notion of generalized semiflow is employed instead of the usual semi-group approach, since solutions of the Cauchy problem for the equation might not be unique. Moreover, the preceding abstract theory is applied to a generalized Allen-Cahn equation as well as a semilinear parabolic equation with a nonlinear term involving gradients.

Keyword: Doubly nonlinear evolution equation, generalized semiflow, global attractor, reflexive Banach space, generalized Allen-Cahn equation

MSC: 37L30, 35B41, 34G25, 35K65

1 Introduction

Let V and V^* be a real reflexive Banach space and its dual space, respectively, and let H be a Hilbert space whose dual space H^* is identified with itself

*Department of Machinery and Control Systems, School of Systems Engineering, Shibaura Institute of Technology, 307 Fukasaku, Minuma-ku, Saitama-shi, Saitama 337-8570 Japan (e-mail: g-akagi@sic.shibaura-it.ac.jp). Supported in part by the Shibaura Institute of Technology grant for Project Research (No. 211459 (2006), 211455 (2007)), and the Grant-in-Aid for Young Scientists (B) (No. 19740073), Ministry of Education, Culture, Sports, Science and Technology.

such that

$$V \hookrightarrow H \equiv H^* \hookrightarrow V^* \quad (1.1)$$

with continuous and densely defined canonical injections. Let φ and ψ be proper lower semi-continuous convex functions from V into $(-\infty, \infty]$, and let $\partial_V \varphi, \partial_V \psi : V \rightarrow 2^{V^*}$ be subdifferential operators of φ and ψ respectively. Moreover, let B be a (possibly) non-monotone and multi-valued operator from V into V^* .

We deal with the following Cauchy problem (denoted by (CP)) for a doubly nonlinear evolution inclusion:

$$\partial_V \psi(u'(t)) + \partial_V \varphi(u(t)) + \lambda B(u(t)) \ni f \text{ in } V^*, \quad 0 < t < \infty, \quad (1.2)$$

$$u(0) = u_0, \quad (1.3)$$

where $f \in V^*$ and $u_0 \in D(\varphi) := \{u \in V; \varphi(u) < \infty\}$ are given data, with a parameter $\lambda \in [0, 1]$, which controls the smallness of the perturbation (i.e., no perturbation if $\lambda = 0$ and no restriction if $\lambda = 1$). Equation (1.2) can be regarded as a perturbation problem for a monotone system $\partial_V \psi(u'(t)) + \partial_V \varphi(u(t)) \ni f$ in V^* , which has been well studied (see, e.g., [5], [11], [20], [21], [29], [6, 7], [39], [28], [30] and the references given therein), and we emphasize that the operator $B(\cdot)$ is non-monotone and has no gradient structure. We first review some existence result of global (in time) strong solutions of (CP) for any $\lambda \in [0, 1]$ (i.e., no restriction on λ) under appropriate conditions such as the coerciveness and the boundedness of $\partial_V \psi$, the precompactness of sub-level sets of φ , and the boundedness and the compactness of B (see Theorem 3.4).

The first purpose of this paper is to study the long-time behavior of global solutions for (CP), in particular, the existence of global attractors. There are a number of works for the study of dynamical systems generated by evolution equations (see, e.g., [24], [25], [12], [38]). Most of them are concerned with single-valued dynamical systems, that is, the case that the solution is uniquely determined by initial data, and they are based on a usual semi-group approach. However, the doubly nonlinear problem (CP) might admit multiple solutions; indeed, Colli [20] gave a simple example that (CP) possesses multiple solutions even if $B \equiv 0$ and $\partial_V \varphi$ is linear. Moreover, the author also exhibited the non-uniqueness of solutions for some doubly nonlinear problems in [1]. Hence (CP) possibly generates a dynamical system with a multi-valued evolution operator, which is called *multi-valued dynamical system*. There are several general theories for handling multi-valued dynamical systems in [35], [17], [8], [26] (see also [36], [27], [40], [16], [41], [9], [19], [32], [18] and references therein). We particularly employ the notion

of generalized semiflow proposed by J.M. Ball [8] to analyze the long-time behavior of solutions for (CP).

There are only a few contributions to the long-time behavior of solutions for doubly nonlinear problems such as (CP). In a Hilbert space setting (i.e., $V = V^* = H$), Segatti [34] proved the global (in time) solvability of (CP) with $B(u) = -u$ under a linear growth condition for $\partial_H \psi$, and moreover, he proved the existence of global attractors by using the notion of generalized semiflow and establishing a dissipative estimate for a Lyapunov functional $J(u) := \varphi(u) - \frac{\lambda}{2}|u|_H^2$. Moreover, Schimperna, Segatti and Stefanelli [33] studied the well-posedness and the long-time behavior of solutions for a concrete doubly nonlinear gradient system in $L^2(\Omega)$ of a non-convex energy functional.

In our setting, the form of the perturbation has not been explicitly specified, and moreover, it might have no gradient structure. Furthermore, we work in a Banach space setting (i.e., $V \neq V^*$). We use techniques recently developed by the author in [3] to handle severe nonlinearities arising from the double nonlinearity and the Banach space setting of (CP), and then, we verify that the set of all solutions of (CP) forms a generalized semiflow in Ball's sense without any restrictions on the parameter $\lambda \in [0, 1]$ (see Theorem 4.1). Moreover, we prove the existence of global attractors for the generalized semiflow in some critical case by imposing a smallness constraint on $\lambda > 0$ (see Theorem 5.1 and Remark 5.5). We also prove the existence of global attractors for an arbitrary $\lambda \in [0, 1]$ in subcritical cases (see Corollary 5.6).

The second purpose of the present paper is to apply the preceding theory to generalized Allen-Cahn equations proposed by Gurtin [23]. We first investigate the asymptotic behavior of solutions $u = u(x, t)$ for a quasilinear Allen-Cahn equation of the form

$$\alpha(u_t) - \Delta_m u + \partial_r W(x, u) \ni f \quad \text{in } \Omega \times (0, \infty), \quad (1.4)$$

where Ω is a bounded domain of \mathbb{R}^N , $f : \Omega \rightarrow \mathbb{R}$ is given, $\alpha(r) = |r|^{p-2}r$ with $p \geq 2$ and Δ_m stands for the so-called m -Laplace operator given by

$$\Delta_m \phi(x) = \nabla \cdot (|\nabla \phi(x)|^{m-2} \nabla \phi(x)), \quad 1 < m < \infty.$$

Moreover, $W(x, r)$ denotes a double-well potential given by

$$W(x, r) = j(r) + \lambda \int_0^r g(x, \rho) d\rho \quad \text{for } x \in \Omega, r \in \mathbb{R}$$

with a proper lower semicontinuous convex function $j : \mathbb{R} \rightarrow [0, +\infty]$, $\lambda \in [0, 1]$ and a Carathéodory function $g : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ possibly non-monotone in the second variable. Then $\partial_r W(x, r)$ stands for the derivative in r of the potential $W(x, r)$, more precisely,

$$\partial_r W(x, r) = \partial j(r) + \lambda g(x, r) \quad \text{for } x \in \Omega \text{ and } r \in \mathbb{R},$$

where ∂j is the subdifferential of j . We reduce the initial-boundary value problem for (1.4) into (CP) in $V = L^p(\Omega)$ by setting $\partial_V \psi(u) = \alpha(u(\cdot))$, $\partial_V \varphi(u) = -\Delta_m u + \partial j(u(\cdot))$ and $B(u) = g(\cdot, u(\cdot))$. It is crucial in our analysis to set $V = L^p(\Omega)$ so that the mapping $\partial_V \psi : u \mapsto \alpha(u(\cdot))$ is coercive and bounded from V into V^* .

We next handle the following semilinear problem with a nonlinear term involving the gradient of u :

$$\alpha(u_t) - \Delta u + N(x, u, \nabla u) = f \quad \text{in } \Omega \times (0, \infty), \quad (1.5)$$

where $N(x, r, \mathbf{p}) = \partial j(r) + \lambda h(x, r, \mathbf{p})$ with a Carathéodory function h from $\Omega \times \mathbb{R} \times \mathbb{R}^N$ into \mathbb{R} . This problem could not be reduced into any gradient systems (cf. [34], [33], [31]); however, (1.5) can be regarded as a perturbation problem of a gradient system, and it also falls within our abstract setting.

This paper is composed of six sections. In Section 2, we briefly review the theory of generalized semiflow. In Section 3, we discuss the existence of global (in time) solutions for (CP). Then we start our analysis of the long-time behavior of solutions for (CP). We first define the set \mathcal{G} of all solutions for (CP) and prove that it forms a generalized semiflow in a metric space X in Section 4. Moreover, we verify the existence of a global attractor for the generalized semiflow \mathcal{G} in Section 5. Applications of the preceding results to generalized Allen-Cahn equations are given in Section 6.

Notation. Let I be a section of \mathbb{R} and let E be a set. We then denote by $AC(I; E)$ (respectively, $AC(I)$) the set of all E -valued (respectively, real-valued) absolutely continuous functions defined on I . Moreover, C stands for a non-negative constant independent of the elements of the corresponding space and set and may vary from line to line.

2 Theory of generalized semiflow

The notion of generalized semiflow was first introduced by J.M. Ball [8]. He also defined global attractors for generalized semiflows and provided a criterion of the existence of global attractors. We first recall the definition of generalized semiflow.

DEFINITION 2.1. *Let X be a metric space with metric $d_X = d_X(\cdot, \cdot)$. A family \mathcal{G} of maps $\varphi : [0, \infty) \rightarrow X$ is said to be a generalized semiflow in X , if the following four conditions are all satisfied:*

(H1) (Existence) *for each $x \in X$ there exists $\varphi \in \mathcal{G}$ such that $\varphi(0) = x$;*

(H2) (Translation invariance) if $\varphi \in \mathcal{G}$ and $\tau \geq 0$, then the map φ^τ also belongs to \mathcal{G} , where $\varphi^\tau(t) := \varphi(t + \tau)$ for $t \in [0, \infty)$;

(H3) (Concatenation invariance) if $\varphi_1, \varphi_2 \in \mathcal{G}$ and $\varphi_2(0) = \varphi_1(\tau)$ at some $\tau \geq 0$, then the map ψ , the concatenation of φ_1 and φ_2 at τ , defined by

$$\psi(t) := \begin{cases} \varphi_1(t) & \text{if } t \in [0, \tau], \\ \varphi_2(t - \tau) & \text{if } t \in (\tau, \infty) \end{cases}$$

also belongs to \mathcal{G} ;

(H4) (Upper semicontinuity) if $\varphi_n \in \mathcal{G}$, $x \in X$ and $\varphi_n(0) \rightarrow x$ in X , then there exist a subsequence (n') of (n) and $\varphi \in \mathcal{G}$ such that $\varphi_{n'}(t) \rightarrow \varphi(t)$ for each $t \in [0, \infty)$.

Let \mathcal{G} be a generalized semiflow in a metric space X . We define a mapping $T(t) : 2^X \rightarrow 2^X$ by

$$T(t)E := \{\varphi(t); \varphi \in \mathcal{G} \text{ and } \varphi(0) \in E\} \quad \text{for } E \subset X \quad (2.1)$$

for each $t \geq 0$. Then one can check from (H1)–(H3) that $(T(t))_{t \geq 0}$ satisfies the semi-group properties, that is, (i) $T(0)$ is the identity mapping in 2^X ; (ii) $T(t)T(s) = T(t + s)$ for all $t, s \geq 0$.

Moreover, global attractors for generalized semiflows are defined as follows.

DEFINITION 2.2. Let \mathcal{G} be a generalized semiflow in a metric space X and let $(T(t))_{t \geq 0}$ be the family of mappings defined as in (2.1). A set $\mathcal{A} \subset X$ is said to be a global attractor for the generalized semiflow \mathcal{G} if the following (i)–(iii) hold.

- (i) \mathcal{A} is compact in X ;
- (ii) \mathcal{A} is invariant under $T(t)$, i.e., $T(t)\mathcal{A} = \mathcal{A}$, for all $t \geq 0$;
- (iii) \mathcal{A} attracts any bounded subsets B of X by $(T(t))_{t \geq 0}$, i.e.,

$$\lim_{t \rightarrow \infty} \text{dist}(T(t)B, \mathcal{A}) = 0,$$

where $\text{dist}(\cdot, \cdot)$ is defined by

$$\text{dist}(A, B) := \sup_{a \in A} \inf_{b \in B} d_X(a, b) \quad \text{for } A, B \subset X.$$

As in the standard theory of dynamical systems for semi-group operators, one can also introduce the notion of ω -limit set.

DEFINITION 2.3. Let \mathcal{G} be a generalized semiflow in a metric space X . For $E \subset X$, the ω -limit set of E for \mathcal{G} is given as follows.

$$\omega(E) := \left\{ x \in X; \text{ there exist sequences } (\varphi_n) \text{ in } \mathcal{G} \text{ and } (t_n) \text{ on } [0, \infty) \right. \\ \left. \text{such that } \varphi_n(0) \text{ is bounded and belongs to } E \text{ for } n \in \mathbb{N}, \right. \\ \left. t_n \rightarrow \infty \text{ and } \varphi_n(t_n) \rightarrow x \text{ in } X \right\}.$$

In order to prove the existence of global attractors for generalized semiflows, we employ the following theorem due to J.M. Ball [8].

THEOREM 2.4 (J.M. Ball [8]). A generalized semiflow \mathcal{G} in a metric space X has a global attractor \mathcal{A} if and only if the following two conditions are satisfied.

(i) \mathcal{G} is point dissipative, that is,

$$\exists a \text{ bounded set } B \subset X, \forall \varphi \in \mathcal{G}, \exists \tau = \tau(\varphi) \geq 0, \forall t \geq \tau, \varphi(t) \in B;$$

(ii) \mathcal{G} is asymptotically compact, that is, for any sequences (φ_n) in \mathcal{G} and (t_n) on $[0, \infty)$, if $(\varphi_n(0))$ is bounded in X and $t_n \rightarrow \infty$, then $(\varphi_n(t_n))$ is precompact in X .

Moreover, \mathcal{A} is the unique global attractor for \mathcal{G} and given by

$$\mathcal{A} = \bigcup \{ \omega(B); B \text{ is a bounded subset of } X \} =: \omega(X).$$

Furthermore, \mathcal{A} is the maximal compact invariant subset of X under the family of mappings $(T(t))_{t \geq 0}$.

The following proposition gives a sufficient condition for the asymptotic compactness of generalized semiflows.

PROPOSITION 2.5 (J.M. Ball [8]). Let \mathcal{G} be a generalized semiflow in a metric space X . If \mathcal{G} satisfies the following conditions:

(i) \mathcal{G} is eventually bounded, that is, for any bounded set $D \subset X$, there exist $\tau = \tau(D) \geq 0$ and a bounded set $B = B(D) \subset X$ such that

$$\bigcup_{t \geq \tau} T(t)D \subset B,$$

(ii) \mathcal{G} is compact, that is, for any sequence (u_n) in \mathcal{G} , if $(u_n(0))$ is bounded in X , then there exists a subsequence (n') of (n) such that $(u_{n'}(t))$ is convergent in X for all $t > 0$,

then \mathcal{G} is asymptotically compact.

3 Global existence

In this section, we discuss the existence of global (in time) solutions for (CP). Let us recall our basic setting again: V and V^* are a real reflexive Banach space and its dual space, respectively, and H is a Hilbert space satisfying (1.1). In addition, we always assume that φ and ψ are proper (i.e., $\varphi \not\equiv \infty$, $\psi \not\equiv \infty$), lower semicontinuous and convex functions from V into $(-\infty, \infty]$ (see also (ii) of Remark 3.5). The subdifferential operator $\partial_V \varphi : V \rightarrow 2^{V^*}$ of φ is defined by

$$\partial_V \varphi(u) := \{\xi \in V^*; \varphi(v) - \varphi(u) \geq \langle \xi, v - u \rangle \text{ for all } v \in D(\varphi)\},$$

where $D(\varphi) := \{u \in V; \varphi(u) < \infty\}$ is the effective domain of φ , with the domain $D(\partial_V \varphi) := \{u \in D(\varphi); \partial_V \varphi(u) \neq \emptyset\}$. Moreover, $\partial_V \psi$ is also defined in a similar way. It is well known that every subdifferential operator is maximal monotone in $V \times V^*$ (see [15], [10], [13] for more details).

The existence of solutions for (CP) has already been studied in more general settings by the author [3]. Throughout this paper we are concerned with strong solutions for (CP) given as follows.

DEFINITION 3.1. *For $T \in (0, \infty)$, a function $u \in AC([0, T]; V)$ is said to be a strong solution of (CP) on $[0, T]$, if the following conditions are satisfied:*

- (i) $u(0) = u_0$,
- (ii) *there exists a negligible set $N \subset (0, T)$, i.e., the Lebesgue measure of N is zero, such that $u(t) \in D(\partial_V \varphi)$ and $u'(t) \in D(\partial_V \psi)$ for all $t \in [0, T] \setminus N$, and moreover, there exist sections $\eta(t) \in \partial_V \psi(u'(t))$, $\xi(t) \in \partial_V \varphi(u(t))$ and $g(t) \in B(u(t))$ such that*

$$\eta(t) + \xi(t) + \lambda g(t) = f \text{ in } V^* \text{ for all } t \in [0, T] \setminus N, \quad (3.1)$$

- (iii) $u(t) \in D(\varphi)$ for all $t \in [0, T]$, and the function $t \mapsto \varphi(u(t))$ is absolutely continuous on $[0, T]$.

Furthermore, for $T \in (0, \infty]$, a function $u \in AC([0, T]; V)$ is said to be a strong solution of (CP) on $[0, T)$, if u is a strong solution of (CP) on $[0, S]$ for every $S \in (0, T)$.

In order to discuss the existence of solutions for (CP), we give basic assumptions with parameters $p \in (1, \infty)$, $T > 0$, $\varepsilon > 0$ in the following. Here and thereafter, p' denotes the Hölder conjugate of $p \in (1, \infty)$, i.e., $p' := p/(p - 1) \in (1, \infty)$, and moreover, the graph of an operator A is denoted by A again, i.e., $[u, \xi] \in A$ means that $u \in D(A)$ and $\xi \in A(u)$.

(A1) There exist constants $C_1 > 0$, $C_2 \geq 0$ such that

$$C_1|u|_V^p \leq \psi(u) + C_2 \quad \text{for all } u \in D(\psi).$$

(A2) There exist constants $C_3, C_4 \geq 0$ such that

$$|\eta|_{V^*}^{p'} \leq C_3\psi(u) + C_4 \quad \text{for all } [u, \eta] \in \partial_V \psi.$$

(Φ1) There exist a reflexive Banach space X_0 and a non-decreasing function ℓ_1 in \mathbb{R} such that X_0 is compactly embedded in V and

$$|u|_{X_0} \leq \ell_1(|u|_H + \varphi(u)) \quad \text{for all } u \in D(\partial_V \varphi).$$

(B1) $_\varepsilon$ $D(\partial_V \varphi) \subset D(B)$. There exists a constant $c_\varepsilon \geq 0$ such that

$$|g|_{V^*}^{p'} \leq \varepsilon|\xi|_{V^*}^\sigma + c_\varepsilon(|\varphi(u)| + |u|_V^p + 1) \quad \text{with } \sigma := \min\{2, p'\}$$

for all $u \in D(\partial_V \varphi)$, $g \in B(u)$ and $\xi \in \partial_V \varphi(u)$.

(B2) Let $S \in (0, T]$ and let (u_n) and (ξ_n) be sequences in $C([0, S]; V)$ and $L^\sigma(0, S; V^*)$ with $\sigma := \min\{2, p'\}$, respectively, such that $u_n \rightarrow u$ strongly in $C([0, S]; V)$, $[u_n(t), \xi_n(t)] \in \partial_V \varphi$ for a.e. $t \in (0, S)$, and

$$\sup_{t \in [0, S]} |\varphi(u_n(t))| + \int_0^S |u'_n(t)|_H^p dt + \int_0^S |\xi_n(t)|_{V^*}^\sigma dt$$

is bounded for all $n \in \mathbb{N}$,

and let (g_n) be a sequence in $L^{p'}(0, S; V^*)$ such that $g_n(t) \in B(u_n(t))$ for a.e. $t \in (0, S)$ and $g_n \rightarrow g$ weakly in $L^{p'}(0, S; V^*)$. Then (g_n) is precompact in $L^{p'}(0, S; V^*)$ and $g(t) \in B(u(t))$ for a.e. $t \in (0, S)$.

(B3) Let $S \in (0, T]$ and $u \in C([0, S]; V) \cap W^{1,p}(0, S; H)$ be such that $\sup_{t \in [0, S]} |\varphi(u(t))| < \infty$ and suppose that there exists $\xi \in L^{p'}(0, S; V^*)$ such that $\xi(t) \in \partial_V \varphi(u(t))$ for a.e. $t \in (0, S)$. Then there exists a V^* -valued strongly measurable function g such that $g(t) \in B(u(t))$ for a.e. $t \in (0, S)$. Moreover, the set $B(u)$ is convex for all $u \in D(B)$.

Here let us give a remark on assumptions for the non-monotone multivalued operator $B : V \rightarrow V^*$.

REMARK 3.2. (i) Condition (B1) $_\varepsilon$ provides some growth condition for $B : V \rightarrow V^*$ with a constant $\varepsilon > 0$. Condition (B2) means that the operator $\mathcal{B} : u \mapsto B(u(\cdot))$ is compact and closed in the sense of multivalued operators. Moreover, we also assume (B3) so that the set $\mathcal{B}(u)$ lies on a proper Bochner-Lebesgue space for u sufficiently regular in time and it is convex.

- (ii) In case B is single-valued, one can replace (B2) by simpler conditions. The following condition is much simpler, but somewhat restrictive in view of applications to PDEs.

(B2)₁ B is locally Lipschitz continuous from V into V^* .

We also give a milder one.

(B2)₂ For each $S \in (0, T)$, the operator $\mathcal{B} : u \mapsto B(u(\cdot))$ is continuous and compact from $L^\infty(0, S; X_0) \cap W^{1,p}(0, S; H)$ into $L^{p'}(0, S; V^*)$,

where X_0 is the Banach space appeared in $(\Phi 1)$. Each of (B2)₁ and (B2)₂ implies (B2) under (B1) _{ε} and $(\Phi 1)$. Moreover, (B2)₂ can be also derived from a pointwise condition:

(B2)₃ There exists a Banach space W such that X_0 is compactly embedded in W and W is continuously embedded in H . Moreover, it holds that

$$|B(u) - B(v)|_{V^*} \leq \ell_2(|u|_{X_0})\ell_3(|u - v|_W) \quad \text{for all } u, v \in X_0$$

with non-decreasing functions ℓ_2, ℓ_3 in \mathbb{R} satisfying $\ell_3(s) \rightarrow 0$ as $s \rightarrow +0$.

Indeed, if (u_n) is bounded in $L^\infty(0, S; X_0) \cap W^{1,p}(0, S; H)$, then by Aubin-Lions's compactness theorem, up to a subsequence, one can ensure that $u_n \rightarrow u$ strongly in $C([0, S]; W)$. Thus $B(u_n(\cdot)) \rightarrow B(u(\cdot))$ strongly in $L^\infty(0, S; V^*)$.

- (iii) If B is single-valued and either (B2)₁ or (B2)₂ holds, then (B3) is automatically verified.

REMARK 3.3. Assumption (A2) implies that

(A2)' There exists a constant $C_5 \geq 0$ such that

$$\frac{1}{2}\psi(u) \leq \langle \eta, u \rangle + C_5 \quad \text{for all } [u, \eta] \in \partial_V \psi.$$

For its proof, we refer the reader to Proposition 3.2 of [3].

Now, applying the abstract theory developed in [3], we can assure that

THEOREM 3.4 (Global existence). *Let $p \in (1, \infty)$, $T > 0$ and $\lambda \in [0, 1]$ be fixed. Suppose that (A1), (A2), (Φ 1), (B1) $_{\varepsilon}$ –(B3) are all satisfied with a sufficiently small $\varepsilon > 0$ (the smallness of ε is determined only by p , C_1 , C_3 and $C_H := \sup\{|u|_H/|u|_V; u \in V \setminus \{0\}\} > 0$). Then, for all $f \in V^*$ and $u_0 \in D(\varphi)$, there exists at least one strong solution $u \in W^{1,p}(0, T; V)$ on $[0, T]$ such that*

$$\eta, \xi, g \in L^{p'}(0, T; V^*),$$

where $\eta(\cdot)$, $\xi(\cdot)$ and $g(\cdot)$ are the sections of $\partial_V \psi(u'(\cdot))$, $\partial_V \varphi(u(\cdot))$ and $B(u(\cdot))$, respectively, as in (3.1).

We close this section with the following remark.

REMARK 3.5. (i) As in §6.1 of [3], we have:

PROPOSITION 3.6. *Let $p \in (1, \infty)$ and $T > 0$ be given. Assume that (A2) and (B1) $_{\varepsilon}$ hold with an arbitrary constant $\varepsilon \in (0, 4^{1-p'})$. Let $\lambda \in [0, 1]$ and let u be a strong solution of (CP) on $[0, T]$ and let $g(\cdot)$ denote the section of $B(u(\cdot))$ as in (3.1). Then there exist constants $\gamma_{\varepsilon} = \gamma_{\varepsilon}(p, C_3) \geq 0$ and $M_{\varepsilon} = M_{\varepsilon}(p, c_{\varepsilon}) \geq 0$ such that*

$$|g(t)|_{V^*}^{p'} \leq M_{\varepsilon} \left(|f|_{V^*}^{p'} + C_4 + |\varphi(u(t))| + |u(t)|_V^p + 1 \right) + \gamma_{\varepsilon} \psi(u'(t))$$

for a.e. $t \in (0, T)$. Here γ_{ε} depends on ε as follows:

$$\gamma_{\varepsilon} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow +0.$$

Proof. By (B1) $_{\varepsilon}$, we have

$$\begin{aligned} |g(t)|_{V^*}^{p'} &\leq \varepsilon |\xi(t)|_{V^*}^{p'} + c_{\varepsilon} (|\varphi(u(t))| + |u(t)|_V^p + 1) \\ &\leq 4^{p'-1} \varepsilon \left(|f|_{V^*}^{p'} + |\eta(t)|_{V^*}^{p'} + |g(t)|_{V^*}^{p'} + 1 \right) \\ &\quad + c_{\varepsilon} (|\varphi(u(t))| + |u(t)|_V^p + 1), \end{aligned}$$

which together with (A2) implies

$$\begin{aligned} (1 - 4^{p'-1} \varepsilon) |g(t)|_{V^*}^{p'} &\leq 4^{p'-1} \varepsilon \left(|f|_{V^*}^{p'} + C_4 + 1 \right) + 4^{p'-1} \varepsilon C_3 \psi(u'(t)) \\ &\quad + c_{\varepsilon} (|\varphi(u(t))| + |u(t)|_V^p + 1). \end{aligned}$$

By setting $\gamma_{\varepsilon} := 4^{p'-1} \varepsilon C_3 / (1 - 4^{p'-1} \varepsilon) > 0$, we can derive our desired result. \square

(ii) We can assume $\varphi \geq 0$ and $\psi \geq 0$ without any loss of generality (see Remark 3.8 of [3]). In the rest of this paper, we always assume so.

4 Formation of a generalized semiflow

Our analysis of the large-time behavior of solutions for (CP) is based on the theory of generalized semiflow briefly reviewed in §2. Let $X := D(\varphi)$ be a metric space equipped with the distance $d_X(\cdot, \cdot)$ defined by

$$d_X(u, v) := |u - v|_V + |\varphi(u) - \varphi(v)| \quad \text{for } u, v \in X$$

and set

$$\mathcal{G} := \{u \in AC([0, \infty); X); u \text{ is a strong solution of (CP) on } [0, \infty) \\ \text{with some } u_0 \in X\}.$$

Then our result of this section reads,

THEOREM 4.1. *Let $p \in (1, \infty)$ and $\lambda \in [0, 1]$ be fixed. Assume that (A1), (A2), (Φ 1), (B1) $_{\varepsilon}$ –(B3) are satisfied with a sufficiently small $\varepsilon > 0$ for any $T > 0$ (the smallness of ε is determined only by p, C_1, C_3 and C_H). Then \mathcal{G} is a generalized semiflow in X .*

Proof. It suffices to check the four conditions (H1)–(H4) (see Definition 2.1). Since (H2) and (H3) follow immediately, we give proofs only for (H1), (H4).

Proof of (H1). Let $u_0 \in X$ be given. Then due to Theorem 3.4, for any $T > 0$, the Cauchy problem (CP) admits at least one strong solution $u_1 \in W^{1,p}(0, T; V)$. Moreover, since $u_1(T)$ also belongs to $D(\varphi)$, we can also obtain a strong solution $u_2 \in W^{1,p}(0, T; V)$ of (CP) with u_0 replaced by $u_1(T)$. Iterating the argument above, we can construct a sequence $(u_k)_{k \in \mathbb{N}}$ in $W^{1,p}(0, T; V)$ such that u_{k+1} becomes a strong solution of (CP) with the initial condition $u_{k+1}(0) = u_k(T)$ for each $k \in \mathbb{N}$.

Furthermore, define $u : [0, \infty) \rightarrow V$ by

$$u(t) := u_k(t - (k - 1)T) \quad \text{if } t \in [(k - 1)T, kT].$$

Then $u \in W_{loc}^{1,p}([0, \infty); V)$ and $\varphi(u(\cdot)) \in AC([0, \infty))$, and moreover, the restriction of u on $[0, S]$ becomes a strong solution of (CP) for any $S > 0$. Therefore u is a strong solution of (CP) on $[0, \infty)$, which implies that $u \in \mathcal{G}$ and $u(0) = u_0$. Thus (H1) follows.

Proof of (H4). Let $u_n \in \mathcal{G}$ and $v \in X$ be such that $u_n(0) \rightarrow v$ in X . Then multiplying (1.2) with $u = u_n$ by $u'_n(t)$, we have

$$\langle \eta_n(t), u'_n(t) \rangle + \langle \xi_n(t), u'_n(t) \rangle + \lambda \langle g_n(t), u'_n(t) \rangle = \langle f, u'_n(t) \rangle$$

with sections $\eta_n(t) \in \partial_V \psi(u'_n(t))$, $\xi_n(t) \in \partial_V \varphi(u_n(t))$, $g_n(t) \in B(u_n(t))$ for a.e. $t \in (0, \infty)$. We then derive from (A2)' and the chain rule for subdifferentials that

$$\begin{aligned} \frac{1}{2} \psi(u'_n(t)) - C_5 + \frac{d}{dt} \varphi(u_n(t)) &= \langle f - \lambda g_n(t), u'_n(t) \rangle \\ &\leq (|f|_{V^*} + \lambda |g_n(t)|_{V^*}) |u'_n(t)|_V. \end{aligned}$$

Furthermore, by Young's inequality and (A1),

$$\begin{aligned} \frac{1}{4} \psi(u'_n(t)) + \frac{d}{dt} \varphi(u_n(t)) \\ \leq M_1 \left(|f|_{V^*}^{p'} + \lambda^{p'} |g_n(t)|_{V^*}^{p'} + C_2 + C_5 + 1 \right) \end{aligned} \quad (4.1)$$

with a constant M_1 depending only on p and C_1 . By Proposition 3.6, we can choose $\varepsilon > 0$ so small that $\gamma_\varepsilon = \gamma_\varepsilon(p, C_3) \leq (8M_1)^{-1}$ (hence, the smallness of ε is determined by p , C_1 and C_3), and then,

$$\begin{aligned} |g_n(t)|_{V^*}^{p'} &\leq M_\varepsilon \left(|f|_{V^*}^{p'} + C_4 + \varphi(u_n(t)) + |u_n(t)|_V^p + 1 \right) \\ &\quad + \frac{1}{8M_1} \psi(u'_n(t)) \end{aligned} \quad (4.2)$$

for a.e. $t \in (0, \infty)$. Then it follows from (4.1) and (4.2) that

$$\begin{aligned} \frac{1}{16} \psi(u'_n(t)) + \frac{C_1}{16} |u'_n(t)|_V^p + \frac{d}{dt} \varphi(u_n(t)) \\ \leq C \left(|f|_{V^*}^{p'} + 1 \right) + \lambda^{p'} M_2 \left(\varphi(u_n(t)) + |u_n(t)|_V^p \right), \end{aligned} \quad (4.3)$$

where $M_2 := M_1 M_\varepsilon$ is independent of λ (this fact will be used in §5).

Here, for each $\delta > 0$, we can take a constant $C_{\delta,p}$ depending only on δ and p such that

$$\frac{d}{dt} |u_n(t)|_V^p \leq \delta |u'_n(t)|_V^p + C_{\delta,p} |u_n(t)|_V^p.$$

Put $\delta = C_1/16$. Therefore

$$\begin{aligned} \frac{1}{16} \psi(u'_n(t)) + \frac{d}{dt} \left(\varphi(u_n(t)) + |u_n(t)|_V^p \right) \\ \leq C \left(|f|_{V^*}^{p'} + 1 \right) + C \left(\varphi(u_n(t)) + |u_n(t)|_V^p \right) \end{aligned} \quad (4.4)$$

for a.e. $t \in (0, \infty)$. Hence integrate both sides over $(0, t)$ to get

$$\begin{aligned} & \frac{1}{16} \int_0^t \psi(u'_n(\tau)) d\tau + \varphi(u_n(t)) + |u_n(t)|_V^p \\ & \leq \varphi(u_0) + |u_0|_V^p + Ct \left(|f|_{V^*}^{p'} + 1 \right) \\ & \quad + C \int_0^t \left(\varphi(u_n(\tau)) + |u_n(\tau)|_V^p \right) d\tau \end{aligned}$$

for all $t \geq 0$. Thus Gronwall's inequality yields

$$\begin{aligned} & \sup_{t \in [0, T]} \left(\varphi(u_n(t)) + |u_n(t)|_V^p \right) \\ & \leq \left\{ \varphi(u_0) + |u_0|_V^p + CT \left(|f|_{V^*}^{p'} + 1 \right) \right\} e^{CT} \end{aligned} \quad (4.5)$$

with an arbitrary positive number $T > 0$. Hence $(\Phi 1)$ implies that $(u_n(t))_{n \in \mathbb{N}}$ is precompact in V for each $t \in [0, T]$. Furthermore, we can also deduce that

$$\int_0^T \psi(u'_n(t)) dt \leq C_T, \quad (4.6)$$

$$\int_0^T |u'_n(t)|_V^p dt \leq C_T \quad (4.7)$$

with a constant $C_T \geq 0$. Here and henceforth, C_T denotes a non-negative constant independent of n and t but possibly depending on T and may vary from line to line.

We particularly put $T = 1$ in (4.5) and (4.7). Then by virtue of Ascoli's compactness theorem, (u_n) becomes precompact in $C([0, 1]; V)$. Hence there exist a subsequence $(n_k^{(1)})$ of (n) and a function $u^{(1)} \in C([0, 1]; V)$ such that

$$u_{n_k^{(1)}} \rightarrow u^{(1)} \quad \text{strongly in } C([0, 1]; V).$$

Moreover, from the arbitrariness of T in (4.5) and (4.7), we can iteratively take a function $u^{(m+1)} \in C([0, m+1]; V)$ and a subsequence $(n_k^{(m+1)})$ of $(n_k^{(m)})$ such that

$$u_{n_k^{(m+1)}} \rightarrow u^{(m+1)} \quad \text{strongly in } C([0, m+1]; V)$$

for each $m \in \mathbb{N}$. Here we remark that the restriction of $u^{(m+1)}$ on $[0, m]$ coincides with $u^{(m)}$ for every $m \in \mathbb{N}$. Furthermore, let (n_k) be the diagonal part of the double sequence $(n_k^{(m)})_{k, m \in \mathbb{N}}$, that is, $n_k := n_k^{(k)}$. Then we obtain

$$u_{n_k}(t) \rightarrow u(t) \quad \text{strongly in } V \quad \text{for each } t \in [0, \infty), \quad (4.8)$$

where $u \in C([0, \infty); V)$ is given by

$$u(t) = u^{(m)}(t) \quad \text{if } t \in [0, m] \text{ for some } m \in \mathbb{N}.$$

We next prove that $u \in \mathcal{G}$, that is, u is a strong solution of (CP) on $[0, \infty)$. Recall (4.5)–(4.7) with an arbitrary $T > 0$ and use (A2), (4.2) and (3.1) to get

$$\int_0^T |\eta_n(t)|_{V^*}^{p'} dt \leq C_T, \quad (4.9)$$

$$\int_0^T |g_n(t)|_{V^*}^{p'} dt \leq C_T, \quad (4.10)$$

$$\int_0^T |\xi_n(t)|_{V^*}^{p'} dt \leq C_T \quad (4.11)$$

for all $n \in \mathbb{N}$. Hence we can take a subsequence of (n_k) , which will be denoted by the same letter, such that

$$u_{n_k} \rightarrow u \quad \text{weakly in } W^{1,p}(0, T; V), \quad (4.12)$$

$$\text{strongly in } C([0, T]; V), \quad (4.13)$$

$$\eta_{n_k} \rightarrow \eta \quad \text{weakly in } L^{p'}(0, T; V^*), \quad (4.14)$$

$$g_{n_k} \rightarrow g \quad \text{weakly in } L^{p'}(0, T; V^*), \quad (4.15)$$

$$\xi_{n_k} \rightarrow \xi \quad \text{weakly in } L^{p'}(0, T; V^*) \quad (4.16)$$

with some $\eta, g, \xi \in L^{p'}(0, T; V^*)$. Therefore from the demiclosedness of sub-differential operators we can derive that $\xi(t) \in \partial_V \varphi(u(t))$ for a.e. $t \in (0, T)$, and moreover, (B2) implies that $g(t) \in B(u(t))$ for a.e. $t \in (0, T)$ and

$$g_{n_k} \rightarrow g \quad \text{strongly in } L^{p'}(0, T; V^*). \quad (4.17)$$

Furthermore, multiply $\eta_{n_k}(t)$ by $u'_{n_k}(t)$ and integrate this over $(0, T)$. It then follows that

$$\begin{aligned} & \int_0^T \langle \eta_{n_k}(t), u'_{n_k}(t) \rangle dt \\ &= -\varphi(u_{n_k}(T)) + \varphi(u_{n_k}(0)) + \int_0^T \langle f - \lambda g_{n_k}(t), u'_{n_k}(t) \rangle dt. \end{aligned}$$

Therefore we get, by (4.12), (4.13) and (4.17),

$$\begin{aligned} & \limsup_{n_k \rightarrow \infty} \int_0^T \langle \eta_{n_k}(t), u'_{n_k}(t) \rangle dt \\ & \leq -\varphi(u(T)) + \varphi(v) + \int_0^T \langle f - \lambda g(t), u'(t) \rangle dt. \end{aligned} \quad (4.18)$$

Here we also used the fact that $u_n(0) \rightarrow v$ in X , in particular, $\varphi(u_n(0)) \rightarrow \varphi(v)$. Thus by Proposition 1.1 of [10, Chap. II] (see also [15]), we obtain $\eta(t) \in \partial_V \psi(u'(t))$ for a.e. $t \in (0, T)$, which implies that u is a strong solution of (CP) with $u_0 = v$ on $[0, T]$. From the arbitrariness of T , we can also verify that u is a strong solution of (CP) on $[0, \infty)$. Hence u belongs to \mathcal{G} .

Finally, we show that $\varphi(u_{n_k}(t)) \rightarrow \varphi(u(t))$ for every $t \in [0, \infty)$, by taking a subsequence of (n_k) independent of t . To do so, we first derive the convergence for a.e. $t \in (0, \infty)$.

LEMMA 4.2. *It follows that*

$$\liminf_{n_k \rightarrow \infty} \varphi(u_{n_k}(t)) = \varphi(u(t)) \quad \text{for a.e. } t \in (0, \infty).$$

Proof. Let $T > 0$ be arbitrarily given. By the definition of subdifferentials, we see

$$\varphi(u_n(t)) \leq \varphi(u(t)) + |\xi_n(t)|_{V^*} |u_n(t) - u(t)|_V.$$

Now, by (4.11), Fatou's lemma ensures that

$$p(\cdot) := \liminf_{n \rightarrow \infty} |\xi_n(\cdot)|_{V^*}^{p'} \in L^1(0, T).$$

Hence $p(t) < \infty$ for a.e. $t \in (0, T)$. Thus we get, by (4.8),

$$\begin{aligned} \liminf_{n_k \rightarrow \infty} \varphi(u_{n_k}(t)) &\leq \varphi(u(t)) + \liminf_{n_k \rightarrow \infty} (|\xi_{n_k}(t)|_{V^*} |u_{n_k}(t) - u(t)|_V) \\ &\leq \varphi(u(t)) + p(t) \left(\lim_{n_k \rightarrow \infty} |u_{n_k}(t) - u(t)|_V \right) \\ &= \varphi(u(t)) \quad \text{for a.e. } t \in (0, T). \end{aligned}$$

Combining this fact with the lower semicontinuity of φ , we can prove this lemma. \square

Continuation of proof of (H4). We next exhibit the convergence of $\varphi(u_{n_k}(t))$ at every $t \in [0, \infty)$. Recalling (4.3), we find

$$\frac{d\zeta_n}{dt}(t) \leq 0 \quad \text{for a.e. } t \in (0, \infty),$$

where ζ_n is an absolutely continuous function from $[0, \infty)$ into \mathbb{R} given by

$$\zeta_n(t) := \varphi(u_n(t)) - Ct \left(|f|_{V^*}^{p'} + 1 \right) - \lambda^{p'} M_2 \int_0^t \left(\varphi(u_n(\tau)) + |u_n(\tau)|_V^p \right) d\tau$$

for $t \in [0, \infty)$. Hence ζ_n is non-increasing on $[0, \infty)$. We further exploit the following lemma (see Lemma 3.3.3 of [4] for more details).

LEMMA 4.3 (Helly). *If f_n is a non-increasing function on $[0, \infty)$, then there exist a non-increasing function $g : [0, \infty) \rightarrow [-\infty, \infty]$ and a subsequence of (n') of (n) independent of t such that*

$$\lim_{n' \rightarrow \infty} f_{n'}(t) = g(t) \quad \text{for all } t \in [0, \infty).$$

Continuation of proof of (H4). We apply Lemma 4.3 to our situation with $f_n = \zeta_n$ and write ϕ for the non-increasing function g . Then

$$\lim_{n_k \rightarrow \infty} \zeta_{n_k}(t) = \phi(t) \quad \text{for all } t \in [0, \infty).$$

It remains only to reveal the representation of ϕ .

LEMMA 4.4. *For each $t \in [0, \infty)$, it follows that*

$$\phi(t) = \varphi(u(t)) - Ct \left(|f|_{V^*}^{p'} + 1 \right) - \lambda^{p'} M_2 \int_0^t \left(\varphi(u(\tau)) + |u(\tau)|_V^p \right) d\tau.$$

Proof. Let $T > 0$ be fixed. We can then derive that

$$\int_0^t \varphi(u_{n_k}(\tau)) d\tau \rightarrow \int_0^t \varphi(u(\tau)) d\tau \quad \text{for all } t \in [0, T]$$

from the definition of subdifferentials together with (4.13) and (4.16). Hence the definition of ζ_n and Lemma 4.2 yield

$$\liminf_{n_k \rightarrow \infty} \zeta_{n_k}(t) = \zeta(t) \quad \text{for a.e. } t \in (0, T),$$

where $\zeta \in AC([0, \infty))$ is given by

$$\zeta(t) := \varphi(u(t)) - Ct \left(|f|_{V^*}^{p'} + 1 \right) - \lambda^{p'} M_2 \int_0^t \left(\varphi(u(\tau)) + |u(\tau)|_V^p \right) d\tau.$$

Therefore we can obtain $\phi(t) = \zeta(t)$ for a.e. $t \in (0, \infty)$ from the arbitrariness of $T > 0$.

Furthermore, since ζ is continuous and ϕ is non-increasing on $[0, \infty)$, we claim that

$$\phi(t) = \zeta(t) \quad \text{for all } t \in [0, \infty).$$

Indeed, in case $t \in (0, \infty)$, since $\zeta = \phi$ a.e. in $(0, \infty)$, for each $n \in \mathbb{N}$ there exist $t_n \in (t - 1/(2n), t) \cap [0, \infty)$ and $s_n \in (t, t + 1/(2n))$ such that $\phi = \zeta$ at t_n and s_n . Since ϕ is non-increasing, we find that $\phi(t_n) \geq \phi(t) \geq \phi(s_n)$. Hence letting $n \rightarrow \infty$, we deduce that $\phi(t) = \zeta(t)$, by the continuity of ζ . In case $t = 0$, it follows that $\phi(0) = \lim_{n_k \rightarrow \infty} \zeta_{n_k}(0) = \lim_{n_k \rightarrow \infty} \varphi(u_{n_k}(0)) = \varphi(v) = \zeta(0)$ from the fact that $\varphi(u_n(0)) \rightarrow \varphi(v)$. \square

Continuation of proof of (H4). From Lemma 4.4, we conclude that $u_{n_k}(t) \rightarrow u(t)$ in X for each $t \in [0, \infty)$. Thus (H4) follows. \square

5 Existence of global attractors

In this section, we prove the existence of global attractors for the generalized semiflow \mathcal{G} . To this end, we introduce the following structure condition, which plays an essential role in our proof (see also Remark 5.5).

(S1) For all $\lambda \in [0, \bar{\lambda}]$ with some $\bar{\lambda} \in (0, 1]$, there exist constants $\alpha > 0$ independent of λ and $C_6 = C_6(\lambda) \geq 0$ (possibly depending on λ) such that

$$\alpha \left(\varphi(u) + |u|_V^p \right) \leq \langle \xi + \lambda g, u \rangle + C_6$$

for all $u \in D(\partial_V \varphi) \cap D(B)$, $\xi \in \partial_V \varphi(u)$ and $g \in B(u)$.

Our result reads,

THEOREM 5.1. *Let $p \in (1, \infty)$ be fixed and assume that (A1), (A2), ($\Phi 1$) and (B1) $_\varepsilon$ –(B3) are satisfied with a sufficiently small $\varepsilon > 0$ (the smallness of ε is determined only by p , C_1 , C_3 and C_H) for any $T > 0$ such that every constant and function appeared in the assumptions are independent of λ . Moreover, suppose that (S1) is satisfied. Then there exists $\lambda_0 \in (0, \bar{\lambda}]$ such that if $\lambda \in [0, \lambda_0]$ then the generalized semiflow \mathcal{G} has a unique global attractor \mathcal{A} , which is given by*

$$\mathcal{A} := \bigcup \{ \omega(B); B \text{ is a bounded subset of } X \} = \omega(X).$$

Furthermore, \mathcal{A} is the maximal compact invariant subset of X .

In order to prove this theorem, thanks to Theorem 2.4 and Proposition 2.5, it suffices to prove that the generalized semiflow \mathcal{G} is point dissipative, eventually bounded and compact. To do so, we first establish a dissipative estimate for $u \in \mathcal{G}$ in X .

LEMMA 5.2. *Under the same assumptions as in Theorem 5.1, there exist constants $\lambda_0 \in (0, \bar{\lambda}]$, $R \geq 0$ and an increasing function $T_0(\cdot)$ on $[0, \infty)$ such that if $\lambda \in [0, \lambda_0]$, then*

$$\begin{aligned} \varphi(u(t)) + |u(t)|_V^p &\leq R && \text{for all } u_0 \in X, u \in \mathcal{G} \text{ satisfying } u(0) = u_0 \\ &&& \text{and } t \geq T_0(\varphi(u_0) + |u_0|_V^p). \end{aligned} \quad (5.1)$$

Proof. Let $u_0 \in X$ and let $u \in \mathcal{G}$ be such that $u(0) = u_0$. By (S1) and (3.1), we find that

$$\begin{aligned} \alpha \left(\varphi(u(t)) + |u(t)|_V^p \right) &\leq \langle \xi(t) + \lambda g(t), u(t) \rangle + C_6 \\ &= \langle f - \eta(t), u(t) \rangle + C_6 \\ &\leq C \left(|f|_{V^*}^{p'} + |\eta(t)|_{V^*}^{p'} + 1 \right) + \frac{\alpha}{2} |u(t)|_V^p \end{aligned}$$

with some sections $\xi(t) \in \partial_V \varphi(u(t))$, $g(t) \in B(u(t))$, $\eta(t) \in \partial_V \psi(u'(t))$ for a.e. $t \in (0, \infty)$. Thus by (A2) it follows that

$$\frac{\alpha}{2} \left(\varphi(u(t)) + |u(t)|_V^p \right) \leq C \left(|f|_{V^*}^{p'} + 1 \right) + C_7 \psi(u'(t)) \quad (5.2)$$

for a.e. $t \in (0, \infty)$ with some constant $C_7 = C_7(p, \alpha, C_3) \geq 0$ independent of λ .

Recall (4.3) with u_n replaced by u and note that

$$\frac{C_1}{16} |u'(t)|_V^p \geq \sigma |u'(t)|_V^p \geq \sigma \frac{d}{dt} |u(t)|_V^p - \frac{p}{p'} \sigma |u(t)|_V^p,$$

where

$$\sigma := \min \left\{ \frac{C_1}{16}, \frac{\alpha p'}{64pC_7}, 1 \right\} \in (0, 1].$$

Then it follows that

$$\begin{aligned} & \frac{1}{16} \psi(u'(t)) + \frac{d}{dt} \left(\varphi(u(t)) + \sigma |u(t)|_V^p \right) \\ & \leq C \left(|f|_{V^*}^{p'} + 1 \right) + \left(\lambda^{p'} M_2 + \frac{p}{p'} \sigma \right) \left(\varphi(u(t)) + |u(t)|_V^p \right) \end{aligned} \quad (5.3)$$

for a.e. $t \in (0, \infty)$. Multiplying (5.2) by $(16C_7)^{-1}$ and adding this to (5.3), we derive

$$\begin{aligned} \frac{d}{dt} \left(\varphi(u(t)) + \sigma |u(t)|_V^p \right) + \left(\frac{\alpha}{32C_7} - \lambda^{p'} M_2 - \frac{p}{p'} \sigma \right) \left(\varphi(u(t)) + |u(t)|_V^p \right) \\ \leq C \left(|f|_{V^*}^{p'} + 1 \right) \end{aligned}$$

for a.e. $t \in (0, \infty)$. Set $\phi(t) := \varphi(u(t)) + \sigma |u(t)|_V^p$. If $\lambda > 0$ is so small that

$$\beta := \frac{\alpha}{32C_7} - \lambda^{p'} M_2 - \frac{p}{p'} \sigma \geq \frac{\alpha}{64C_7} - \lambda^{p'} M_2 > 0,$$

then

$$\frac{d\phi}{dt}(t) + \beta \phi(t) \leq F := C \left(|f|_{V^*}^{p'} + 1 \right) \quad \text{for a.e. } t \in (0, \infty). \quad (5.4)$$

By using standard techniques for differential inequalities, we have

$$\phi(t) \leq \frac{F}{\beta} + \phi(0) e^{-\beta t} \quad \text{for all } t \in [0, \infty),$$

in particular,

$$\phi(t) \leq \frac{F}{\beta} + 1 \quad \text{for all } t \geq \log(\phi(0) + 1)/\beta.$$

Thus, by putting $R := F/(\sigma\beta) + 1/\sigma$ and $T_0(\cdot) := \log(\cdot + 1)/\beta$, we obtain (5.1). \square

The dissipative estimate obtained above implies the point dissipativity and the eventual boundedness of \mathcal{G} .

LEMMA 5.3. *Under the same assumptions as in Theorem 5.1, the following (i) and (ii) are satisfied.*

(i) \mathcal{G} is point dissipative.

(ii) \mathcal{G} is eventually bounded.

Proof. Let $R \geq 0$ and $T_0(\cdot)$ be the constant and the increasing function given by Lemma 5.2 respectively. Moreover, we write

$$B_r := \{v \in X; \varphi(v) + |v|_V^p \leq r\} \quad \text{for } r > 0.$$

Proof of (i). Put $B := B_R$. Let $u \in \mathcal{G}$ and set $\tau := T_0(\varphi(u(0)) + |u(0)|_V^p)$. Then by Lemma 5.2, we can deduce that $u(t) \in B$ for all $t \geq \tau$.

Proof of (ii). Let D be a bounded set in X . Then we can take $R_1 \in (0, \infty)$ such that $D \subset B_{R_1}$. Moreover, put $\tau := T_0(R_1)$ and $B := B_R$. Then by Lemma 5.2, for any $u \in \mathcal{G}$ with $u(0) \in D \subset B_{R_1}$, it follows that $u(t) \in B$ for all $t \geq \tau$. \square

Concerning the compactness of \mathcal{G} , we have:

LEMMA 5.4. *Under the same assumptions as in Theorem 5.1, \mathcal{G} is compact.*

Proof. Since $u_n(0)$ is bounded in X , i.e., $|u_n(0)|_V + \varphi(u_n(0)) \leq C$ with some constant C independent of n , the estimates (4.5)–(4.7), (4.9)–(4.11) and the convergences (4.8), (4.12)–(4.17) are established with an arbitrary $T > 0$ as in the proof of (H4) (see the proof of Theorem 4.1). Moreover, we can also verify that there exist a subsequence (n_k) of (n) and $u \in C([0, \infty); X)$ such that

$$\varphi(u_{n_k}(t)) \rightarrow \varphi(u(t)) \quad \text{for all } t \in (0, \infty). \quad (5.5)$$

Thus $u_{n_k}(t) \rightarrow u(t)$ in X for each positive t . This completes our proof. \square

Proof of Theorem 5.1. Thanks to Theorem 2.4 and Proposition 2.5, we can immediately derive the conclusion of Theorem 5.1 from Lemmas 5.2–5.4. \square

REMARK 5.5. Theorem 5.1 treats a critical case that the perturbation $B(u)$ of (CP) almost balances with the leading term $\partial_V \varphi(u)$ in a sense and the domination will be determined by the parameter $\lambda > 0$. Indeed, the following concrete PDE problem can be reduced into (CP) concerned in Theorem 5.1:

$$u_t - \Delta u = \lambda u \quad \text{in } \Omega \times (0, \infty), \quad u = 0 \quad \text{on } \partial\Omega \times (0, \infty), \quad u(\cdot, 0) = u_0 \quad \text{in } \Omega,$$

where Ω is a bounded domain of \mathbb{R}^N , by putting $X_0 = X = H_0^1(\Omega)$, $V = H = V^* = L^2(\Omega)$, $\partial_V \psi(u) = u$, $\partial_V \varphi(u) = -\Delta u$ and $B(u) = -u$. Then the unique solution $u = u(x, t)$ of (CP) for some data is unbounded as $t \rightarrow \infty$ (hence there is no bounded absorbing set) for the case $\lambda > \lambda_1$, where $\lambda_1 = \lambda_1(\Omega)$ denotes the principal eigen-value of $-\Delta$ with the homogeneous Dirichlet boundary condition, and every solution converges to the origin for the case $\lambda < \lambda_1$. Moreover, we also note that the inequality of (S1) holds only for $\lambda < \lambda_1$.

Let us next consider the case that the perturbation $B(u)$ is completely dominated by the leading term $\partial_V \varphi(u)$. Then we can remove the constraint on the smallness of $\lambda > 0$ in Theorem 5.1. To do so, we use the following condition instead of $(B1)_\varepsilon$.

(B1) $D(\partial_V \varphi) \subset D(B)$. For any $\varepsilon > 0$, there exist constants $r_\varepsilon \in [0, 1)$ and $c_\varepsilon \geq 0$ such that

$$|g|_{V^*}^{p'} \leq \varepsilon |\xi|_{V^*}^\sigma + c_\varepsilon \{(|\varphi(u)| + |u|_V^p)^{r_\varepsilon} + 1\} \quad \text{with } \sigma := \min\{2, p'\}$$

for all $u \in D(\partial_V \varphi)$, $g \in B(u)$ and $\xi \in \partial_V \varphi(u)$.

More precisely, we have:

COROLLARY 5.6. *Let $p \in (1, \infty)$ be fixed and assume that (A1), (A2), (Φ 1) and (B1)–(B3) for any $T > 0$. Moreover, suppose that (S1) is satisfied with $\bar{\lambda} = 1$. Then the same conclusion as in Theorem 5.1 holds true for any $\lambda \in [0, 1]$.*

Proof. The smallness of λ is used only in the proof of Lemma 5.2. So it suffices to prove this lemma without the smallness by assuming (B1) instead of $(B1)_\varepsilon$. Let $u \in \mathcal{G}$. Then as in Proposition 3.6, it follows from (B1) that there exists a constant $\gamma_\varepsilon > 0$ such that $\gamma_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$ and

$$|g(t)|_{V^*}^{p'} \leq M_\varepsilon \left\{ |f|_{V^*}^{p'} + C_4 + \left(\varphi(u(t)) + |u(t)|_V^p \right)^{r_\varepsilon} + 1 \right\} + \gamma_\varepsilon \psi(u'(t)) \quad (5.6)$$

for a.e. $t \in [0, \infty)$. Hence as in the proof of Lemma 5.2, by choosing $\varepsilon > 0$ and $\sigma > 0$ small enough,

$$\begin{aligned} & \frac{1}{16} \psi(u'(t)) + \frac{d}{dt} \left(\varphi(u(t)) + \sigma |u(t)|_V^p \right) \\ & \leq C \left(|f|_{V^*}^{p'} + 1 \right) + \lambda^{p'} M_2 \left(\varphi(u(t)) + |u(t)|_V^p \right)^{r_\varepsilon} + \frac{p}{p'} \sigma |u(t)|_V^p \quad (5.7) \end{aligned}$$

for a.e. $t \in (0, \infty)$. Since $r_\varepsilon \in (0, 1)$, by Young's inequality, for any $\mu > 0$ we can take a constant $d_\mu > 0$ such that

$$\left(|\varphi(u)| + |u|_V^p\right)^{r_\varepsilon} \leq \mu \left(|\varphi(u)| + |u|_V^p\right) + d_\mu.$$

Here we particularly choose $\mu > 0$ so small that

$$\beta := \frac{\alpha}{32C_7} - \lambda^{p'} \mu M_2 - \frac{p}{p'} \sigma \geq \frac{\alpha}{64C_7} - \lambda^{p'} \mu M_2 > 0.$$

Then we obtain (5.4) from (5.7) and the rest of proof runs as before. \square

6 Applications to generalized Allen-Cahn equations

Gurtin [23] proposed a generalized Allen-Cahn equation, which describes the evolution of an order parameter $u = u(x, t)$, of the form

$$\rho(u, \nabla u, u_t) u_t = \operatorname{div} \left(\partial_{\mathbf{p}} \hat{\psi}(u, \nabla u) \right) - \partial_r \hat{\psi}(u, \nabla u) + f, \quad (6.1)$$

where $\rho = \rho(r, \mathbf{p}, s) \geq 0$ is a kinetic modulus, $\hat{\psi} = \hat{\psi}(r, \mathbf{p})$ denotes a free energy density with its derivatives $\partial_r \hat{\psi}$ and $\partial_{\mathbf{p}} \hat{\psi}$ in r and \mathbf{p} , respectively, and f is an external microforce, by taking account of a balance law for microforces as well as constitutive relations consistent with the second law of thermodynamics. As a simple instance of the free energy density $\hat{\psi}$, we often take

$$\hat{\psi}(r, \mathbf{p}) = \frac{1}{2} |\mathbf{p}|^2 + W(r)$$

with a double-well potential $W(r)$ (e.g., $W(r) = (r^2 - 1)^2$). Moreover, if $\rho \equiv 1$, then (6.1) coincides with the usual Allen-Cahn equation,

$$u_t - \Delta u + \partial_r W(u) = f.$$

In §6.1, we particularly deal with a generalized Allen-Cahn equation of quasilinear type such as

$$\left. \begin{aligned} \alpha(u_t) - \Delta_m u + \partial_r W(x, u) &\ni f && \text{in } \Omega \times (0, \infty), \\ u &= 0 && \text{on } \partial\Omega \times (0, \infty), \\ u(\cdot, 0) &= u_0 && \text{in } \Omega, \end{aligned} \right\} \quad (6.2)$$

where $u_0, f : \Omega \rightarrow \mathbb{R}$ are given, $\alpha(r) = |r|^{p-2}r$ with $p \geq 2$, and Δ_m stands for the so-called m -Laplace operator given by

$$\Delta_m u(x) = \nabla \cdot (|\nabla u(x)|^{m-2} \nabla u(x)), \quad 1 < m < \infty.$$

Here we set

$$\partial_r W(x, r) = \partial j(r) + \lambda g(x, r)$$

with the subdifferential ∂j of a proper, lower semicontinuous convex function $j : \mathbb{R} \rightarrow [0, +\infty]$, $\lambda \in [0, 1]$ and a Carathéodory function $g : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ possibly non-monotone in the second variable. Then $\partial_r W$ can be regarded as the derivative (in a proper sense) of a potential $W = W(x, r) : \Omega \times \mathbb{R} \rightarrow (-\infty, +\infty]$ given by

$$W(x, r) := j(r) + \lambda \int_0^r g(x, \rho) d\rho \quad \text{for } x \in \Omega, r \in \mathbb{R}. \quad (6.3)$$

As a typical example of j , we take a power function, i.e., $j(r) = |r|^\sigma$, or the indicator function of the unit ball, i.e., $j(r) = 0$ if $|r| \leq 1$; $j(r) = \infty$ if $|r| > 1$. Here one can easily find that (6.2) is reduced into Gurtin's generalized Allen-Cahn equation (6.1) with a spatially inhomogeneous free energy density $\hat{\psi} = \hat{\psi}(x, r, \mathbf{p})$ by setting

$$\hat{\psi}(x, r, \mathbf{p}) = \frac{1}{m} |\mathbf{p}|^m + W(x, r) \quad \text{and} \quad \rho(r, \mathbf{p}, s) = |s|^{p-2}.$$

In §6.2 we also consider the following equation as a further generalized form of semilinear Allen-Cahn equations.

$$\left. \begin{array}{ll} \alpha(u_t) - \Delta u + N(x, u, \nabla u) = f & \text{in } \Omega \times (0, \infty), \\ u = 0 & \text{on } \partial\Omega \times (0, \infty), \\ u(\cdot, 0) = u_0 & \text{in } \Omega, \end{array} \right\} \quad (6.4)$$

where $N = N(x, r, \mathbf{p})$ is written as follows

$$N(x, r, \mathbf{p}) = \partial j(r) + \lambda h(x, r, \mathbf{p})$$

with a Carathéodory function $h : \Omega \times \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$, and ∂j is single-valued. We emphasize that this problem may not be written as a (generalized) gradient system such as (6.1), since the nonlinear term N depends on the gradient of u . However, this problem can be regarded as a perturbation problem of (6.1).

We shall apply the preceding abstract theory to (6.2) and (6.4) to prove the existence of global attractors in Ball's sense. Throughout this section, let Ω be a bounded domain of \mathbb{R}^N with smooth (e.g., C^2 -class) boundary $\partial\Omega$.

6.1 Quasilinear Allen-Cahn equations

The aim of this subsection is to discuss the existence of global (in time) solutions for (6.2) and their asymptotic behavior. We assume that the mappings $\alpha : \mathbb{R} \rightarrow \mathbb{R}$ and $g : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ satisfy

- (a1) there exists $p \in [2, \infty)$ such that $\alpha(r) = |r|^{p-2}r$ for $r \in \mathbb{R}$;
- (a2) $j = j(r)$ is a proper lower semicontinuous convex function from \mathbb{R} into $[0, \infty]$ such that $\partial j(0) \ni 0$; there exist constants $\sigma > 1$, $C_8 > 0$ and $C_9 \geq 0$ such that

$$C_8|r|^\sigma \leq j(r) + C_9 \quad \text{for all } r \in \mathbb{R};$$

- (a3) $g = g(x, r)$ is a Carathéodory function, i.e., measurable in x and continuous in r . Moreover, there exist $q \geq 1 + 1/p'$, $C_{10} \geq 0$ and $a_1 \in L^1(\Omega)$ such that

$$|g(x, r)|^{p'} \leq C_{10}|r|^{p'(q-1)} + a_1(x)$$

for a.e. $x \in \Omega$ and all $r \in \mathbb{R}$.

Our basic assumption on the exponent p is as follows:

$$2 \leq p < \max\{m^*, \sigma\}, \tag{6.5}$$

where m^* denotes the Sobolev critical exponent, that is,

$$m^* := \frac{Nm}{(N-m)_+} = \begin{cases} \frac{Nm}{N-m} & \text{if } m < N, \\ \infty & \text{if } m \geq N. \end{cases}$$

In order to reduce (6.2) into an abstract Cauchy problem, we set $V = L^p(\Omega)$, $V^* = L^{p'}(\Omega)$ and $H = L^2(\Omega)$. Then (1.1) follows. Moreover, define the functional $\psi : V \rightarrow [0, \infty)$ by

$$\psi(u) := \frac{1}{p} \int_{\Omega} |u(x)|^p dx \quad \text{for } u \in V. \tag{6.6}$$

Then $D(\psi) = D(\partial_V \psi) = V$ and $\partial_V \psi(u)$ coincides with $\alpha(u(\cdot))$ in V^* . Define $\varphi_1, \varphi_2 : V \rightarrow [0, \infty]$ by

$$\varphi_1(u) := \begin{cases} \frac{1}{m} \int_{\Omega} |\nabla u(x)|^m dx & \text{if } u \in W_0^{1,m}(\Omega), \\ \infty & \text{otherwise} \end{cases}$$

and

$$\varphi_2(u) := \begin{cases} \int_{\Omega} j(u(x))dx & \text{if } j(u(\cdot)) \in L^1(\Omega), \\ \infty & \text{otherwise.} \end{cases}$$

It is easily seen that $\partial_V \varphi_1(u) = -\Delta_m u$ with $u|_{\partial\Omega} = 0$ and $\partial_V \varphi_2(u) = \partial j(u(\cdot))$. We further set $\varphi : V \rightarrow [0, \infty]$ by

$$\varphi(u) := \varphi_1(u) + \varphi_2(u) \quad (6.7)$$

with the effective domain $D(\varphi) = D(\varphi_1) \cap D(\varphi_2) = \{u \in W_0^{1,m}(\Omega); j(u(\cdot)) \in L^1(\Omega)\}$. Then since $\partial j(0) \ni 0$, as in Corollary 16 of [14], we observe

$$\varphi_1(J_{\mu}^H u) \leq \varphi_1(u) \quad \text{for } u \in D(\varphi_1) \text{ and } \mu > 0,$$

where J_{μ}^H denotes the resolvent of the subdifferential for the extension of φ_2 onto H by infinity. Then we can verify that $\partial_V \varphi_1 + \partial_V \varphi_2$ is maximal monotone; hence, $\partial_V \varphi(u) = \partial_V \varphi_1(u) + \partial_V \varphi_2(u)$ coincides with $-\Delta_m u + \partial j(u(\cdot))$ equipped with $u|_{\partial\Omega} = 0$ in V^* (see [2]). Furthermore, we define the mapping

$$B : V \rightarrow V^*; u \mapsto g(\cdot, u(\cdot)).$$

Then (6.2) is rewritten into (CP) with the functionals ψ, φ and the mapping B defined above.

In the rest of this paper, a function $u : \Omega \times (0, \infty) \rightarrow \mathbb{R}$ is said to be an L^p -solution of (6.2) on $[0, \infty)$, if u is a strong solution on $[0, \infty)$ of (CP) with ψ, φ, B defined above. Applying the preceding abstract theory to (6.2), we can prove the following theorems.

THEOREM 6.1. *Let Ω be a bounded domain of \mathbb{R}^N with smooth boundary $\partial\Omega$. Suppose that (a1)–(a3) and (6.5) are satisfied, and*

$$p'(q-1) \leq \max\{m, p\} \quad \text{or} \quad \begin{cases} p'(q-1) \leq \sigma & \text{if } \sigma < m^*, \\ p'(q-1) < \sigma & \text{if } \sigma \geq m^*. \end{cases} \quad (6.8)$$

Then for any $\lambda \in [0, 1]$, $f \in L^{p'}(\Omega)$, $u_0 \in X := \{v \in W_0^{1,m}(\Omega); j(v(\cdot)) \in L^1(\Omega)\}$, the initial-boundary value problem (6.2) admits at least one L^p -solution on $[0, \infty)$. Moreover, the set

$$\mathcal{G} := \{u : [0, \infty) \rightarrow L^p(\Omega); u \text{ is an } L^p\text{-solution of (6.2) on } [0, \infty) \\ \text{with some } u_0 \in X\}$$

forms a generalized semiflow in X with the distance $d_X(\cdot, \cdot)$ given by

$$d_X(u, v) := |\varphi(u) - \varphi(v)| + |u - v|_{L^p(\Omega)} \quad \text{for } u \in X,$$

where φ is given by (6.7).

REMARK 6.2. We can derive from the assumptions (6.5) and (6.8) that

- (i) $p'(q-1) \leq \max\{m, p, \sigma\}$,
- (ii) $X_0 := W_0^{1,m}(\Omega) \cap L^\sigma(\Omega)$ is compactly embedded both in $L^p(\Omega)$ and in $L^{p'(q-1)}(\Omega)$.

Indeed, (i) follows immediately from (6.8). Moreover, applying the Rellich-Kondrachov compactness theorem and interpolation inequalities for Lebesgue spaces, we assure by (6.5) that X_0 is compactly embedded in $L^p(\Omega)$. Furthermore, (ii) is similarly derived from (6.5) and (6.8). These two facts play a crucial role in our proofs and we can replace (6.5) and (6.8) by (i) and (ii) in Theorems 6.1 and 6.3.

Proof. Conditions (A1) and (A2) follow immediately from (a1) and (6.6). Set $X_0 := W_0^{1,m}(\Omega) \cap L^\sigma(\Omega)$ with the norm $|\cdot|_{X_0} := (|\nabla \cdot|_{L^m(\Omega)}^2 + |\cdot|_{L^\sigma(\Omega)}^2)^{1/2}$. Then we get, by (a2)

$$|u|_{X_0}^{\min\{m,\sigma\}} \leq C(\varphi(u) + 1) \quad \text{for all } u \in D(\varphi)$$

with some constant $C > 0$. Moreover, by (ii) of Remark 6.2, X_0 is compactly embedded in V . Hence $(\Phi 1)$ holds. As for $(B1)_\varepsilon$, we note by (a3) that

$$|B(u)|_{V^*}^{p'} \leq C_{10} \int_{\Omega} |u(x)|^{p'(q-1)} dx + \int_{\Omega} a_1(x) dx \quad \text{for } u \in L^{p'(q-1)}(\Omega). \quad (6.9)$$

Here by (ii) of Remark 6.2, we have $D(\varphi) \subset X_0 \subset L^{p'(q-1)}(\Omega) \subset D(B)$, and moreover, by (a2)

$$\int_{\Omega} |u(x)|^{p'(q-1)} dx \leq C(\varphi(u) + |u|_V^p + 1) \quad \text{for } u \in D(\varphi),$$

which together with (6.9) implies $(B1)_\varepsilon$. Furthermore, by (6.9), the mapping $u \mapsto B(u(\cdot))$ becomes continuous from $L^{p'(q-1)}(0, T; L^{p'(q-1)}(\Omega))$ into $L^p(0, T; V^*)$ for any $T > 0$ (see Theorem 1.43 of [39]), in particular, (B3) holds. Recall that X_0 is compactly embedded in $L^{p'(q-1)}(\Omega)$ by (ii) of Remark 6.2. Hence we also obtain (B2), since the sequence (u_n) of (B2) becomes precompact in $C([0, S]; L^{p'(q-1)}(\Omega))$ by Aubin-Lions's compactness theorem. Finally, applying Theorems 3.4 and 4.1, we can obtain our desired conclusion. \square

THEOREM 6.3. Let Ω be a bounded domain of \mathbb{R}^N with smooth boundary $\partial\Omega$ and let $f \in L^{p'}(\Omega)$. In addition to (a1)–(a3), (6.5) and (6.8), suppose that

$$p \leq \max\{m, \sigma\} \quad \text{and} \quad \begin{cases} \lambda \text{ is arbitrary} & \text{if } p'(q-1) < \max\{m, \sigma\}, \\ \lambda \text{ is small} & \text{otherwise.} \end{cases} \quad (6.10)$$

Then the generalized semiflow \mathcal{G} has a global attractor \mathcal{A} in X .

Proof. We can derive from (a3), (i) of Remark 6.2 and (6.10) that

$$\begin{aligned}
& \langle \partial_V \varphi(u) + \lambda B(u), u \rangle \\
& \geq \int_{\Omega} |\nabla u(x)|^m dx + \int_{\Omega} (j(u(x)) - j(0)) dx + \lambda \int_{\Omega} g(x, u(x)) u(x) dx \\
& \geq \int_{\Omega} |\nabla u(x)|^m dx + C_8 \int_{\Omega} |u(x)|^\sigma dx - (C_9 + |j(0)|) |\Omega| \\
& \quad - \lambda \int_{\Omega} \left(C_{10} |u(x)|^{p'(q-1)} + a_1(x) \right)^{1/p'} |u(x)| dx \\
& \geq \frac{1}{2} \int_{\Omega} |\nabla u(x)|^m dx + \frac{C_8}{2} \int_{\Omega} |u(x)|^\sigma dx - M_\lambda
\end{aligned}$$

with some constant M_λ depending on λ , by assuming that $\lambda > 0$ is small enough only in case $\max\{m, \sigma\} = q$. Hence (S1) follows from the fact that $p \leq \max\{m, \sigma\}$. Consequently, Theorem 5.1 proves the existence of a global attractor with a sufficiently small $\lambda > 0$.

In addition, assuming $p'(q-1) < \max\{m, \sigma\}$ (then, it is not true that $\max\{m, \sigma\} = q$), we observe

$$\int_{\Omega} |u(x)|^{p'(q-1)} dx \leq C (\varphi(u) + |u|_V^p + 1)^r \quad \text{for } u \in D(\varphi)$$

with some $r \in (0, 1)$. Hence (B1) follows. Therefore Corollary 5.6 ensures the existence of a global attractor for any $\lambda \in [0, 1]$. \square

REMARK 6.4. (i) Our arguments described above are still valid even if (a1) is relaxed into

(a1)' there exist $p \in [2, \infty)$ and a lower semicontinuous convex function $A : \mathbb{R} \rightarrow [0, \infty)$ such that $\partial A = \alpha$ and

$$C_{11}(|r|^p - 1) \leq A(r) \text{ and } |\eta|^{p'} \leq C_{12}(A(r) + 1) \text{ for all } [r, \eta] \in \alpha$$

with some constants $C_{11}, C_{12} \geq 0$.

Furthermore, we can also treat the case where $-\Delta_m u(x, t)$ is replaced by $-\operatorname{div} \mathbf{a}(x, \nabla u(x, t))$ and $\mathbf{a} = \mathbf{a}(x, \mathbf{p})$ is a function from $\Omega \times \mathbb{R}^N$ into \mathbb{R}^N such that \mathbf{a} is continuous and maximal monotone in \mathbf{p} and measurable in x , by imposing appropriate growth conditions on \mathbf{a} in \mathbf{p} (see [3] for more details).

(ii) If we have a boundedness condition of ∂j , we could weaken the assumptions on q , the growth order of the perturbation. Let us consider a simple case that $j(r) = (1/\sigma)|r|^\sigma$ with $\sigma > 1 + 1/p'$. Then $\partial j(r) = |r|^{\sigma-2}r$,

which implies $|\partial j(r)| = |r|^{\sigma-1}$. Here we note that

$$|u|_{L^{p'(\sigma-1)}(\Omega)}^{\sigma-1} = |\partial j(u(\cdot))|_{L^{p'}(\Omega)} \leq |-\Delta_m u + \partial j(u(\cdot))|_{L^{p'}(\Omega)}$$

for $u \in W_0^{1,m}(\Omega) \cap L^{p'(q-1)}(\Omega)$ (see [2]). Hence in case $q < \sigma$, for any $\varepsilon > 0$, we can choose $C_\varepsilon > 0$ such that

$$|u|_{L^{p'(q-1)}(\Omega)}^{p'(q-1)} \leq \varepsilon |u|_{L^{p'(\sigma-1)}(\Omega)}^{p'(\sigma-1)} + C_\varepsilon \leq \varepsilon |\partial_V \varphi(u)|_{V^*}^{p'} + C_\varepsilon.$$

Combining this fact with (6.9), we obtain (B1). Moreover, under the same assumption, $D := W_0^{1,m}(\Omega) \cap L^\sigma(\Omega) \cap L^{p'(\sigma-1)}(\Omega)$ is compactly embedded in $L^{p'(q-1)}(\Omega)$; hence the sequence (g_n) of (B2) becomes pre-compact in $L^{p'}(0, S; V^*)$, since $D(\partial_V \varphi) \subset D$. Thus we can prove Theorem 6.1 with $q < \sigma$ instead of (6.8). Furthermore, as in Theorem 6.3, we can also verify the existence of a global attractor for any $\lambda \in [0, 1]$ under $q < \sigma$ instead of (6.8) by assuming a structure condition such as

$$g(x, r)r \geq -C(|r|^q + 1) \quad \text{for a.e. } x \in \Omega \text{ and } r \in \mathbb{R}.$$

6.2 Semilinear Allen-Cahn equations with perturbations involving gradients

We next treat (6.4). Let us assume

- (a2)' ∂j is single-valued, and (a2) holds;
- (a4) The function $h = h(x, r, \mathbf{p})$ is a Carathéodory function in $\Omega \times \mathbb{R} \times \mathbb{R}^N$ (i.e., measurable in x and continuous in (r, \mathbf{p})). Moreover, there exist constants $q_1, q_2 \geq 1 + 1/p'$, $C_{13} \geq 0$ and $a_2 \in L^1(\Omega)$ such that

$$|h(x, r, \mathbf{p})|^{p'} \leq C_{13}(|r|^{p'(q_1-1)} + |\mathbf{p}|^{p'(q_2-1)}) + a_2(x)$$

for a.e. $x \in \Omega$ and all $(r, \mathbf{p}) \in \mathbb{R} \times \mathbb{R}^N$.

Furthermore, we use the following basic assumption on the exponent p ,

$$2 \leq p < \max\{2^*, \sigma\}. \tag{6.11}$$

Equation (6.4) has no longer any gradient structure, because the nonlinear term N depends on ∇u also. However, (6.4) can be transcribed into a (non-monotone) perturbation problem for a doubly nonlinear gradient system in the form (1.2). Indeed, we set $V = L^p(\Omega)$, $H = L^2(\Omega)$ and $V^* := L^{p'}(\Omega)$ as

well as functionals ψ and φ as in (6.6) and (6.7) with $m = 2$ respectively. Moreover, we put

$$B(u) := h(\cdot, u(\cdot), \nabla u(\cdot)) \quad \text{for } u \in V.$$

Then (6.4) is reduced into (CP). Hence L^p -solutions of (6.4) are also defined as in the case of (6.2). Now, exploiting Theorems 3.4 and 4.1, we can verify:

THEOREM 6.5. *Let Ω be a bounded domain of \mathbb{R}^N with smooth boundary $\partial\Omega$. Suppose that (a1), (a2)', (a4) and (6.11) are satisfied. In addition, assume that*

$$\text{either } q_1 \leq p \text{ or } \begin{cases} p'(q_1 - 1) \leq \sigma & \text{if } \sigma < \max\{2^*, (p')^\dagger\}, \\ p'(q_1 - 1) < \sigma & \text{if } \sigma \geq \max\{2^*, (p')^\dagger\} \end{cases} \text{ holds,} \quad (6.12)$$

and

$$\begin{cases} p'(q_2 - 1) \leq 2 & \text{if } 2 < (p')^*, \\ p'(q_2 - 1) < 2 & \text{if } 2 \geq (p')^*, \end{cases} \quad (6.13)$$

where $(p')^\dagger := Np'/(N - 2p')_+$ and $(p')^* := Np'/(N - p')_+$. Then for any $u_0 \in X := \{v \in H_0^1(\Omega); j(v(\cdot)) \in L^1(\Omega)\}$, $f \in L^{p'}(\Omega)$ and $\lambda \in [0, 1]$, the initial-boundary value problem (6.4) admits at least one L^p -solution on $[0, \infty)$. Moreover, the set

$$\mathcal{G} := \{u : [0, \infty) \rightarrow L^p(\Omega); u \text{ is an } L^p\text{-solution of (6.4) on } [0, \infty) \\ \text{with some } u_0 \in X\}$$

forms a generalized semiflow in X with the distance $d_X(\cdot, \cdot)$ given by

$$d_X(u, v) := |\varphi(u) - \varphi(v)| + |u - v|_{L^p(\Omega)} \quad \text{for } u \in X,$$

where φ is given by (6.7) with $m = 2$.

REMARK 6.6. As in Remark 6.2, assumptions (6.11), (6.12) and (6.13) yield in particular that

- (i) $p'(q_1 - 1) \leq \max\{p, \sigma\}$ and $p'(q_2 - 1) \leq 2$,
- (ii) $H_0^1(\Omega) \cap L^\sigma(\Omega)$ is compactly embedded in $L^p(\Omega)$, and furthermore, $W^{2,p'}(\Omega) \cap H_0^1(\Omega) \cap L^\sigma(\Omega)$ is compactly embedded in $L^{p'(q_1-1)}(\Omega) \cap W^{1,p'(q_2-1)}(\Omega)$.

We can prove Theorems 6.5 and 6.3 by using these two facts instead of (6.11), (6.12) and (6.13).

Proof. Due to Theorems 3.4 and 4.1, it suffices to check (A1), (A2), (Φ 1), (B 1) $_{\varepsilon}$ –(B 3). As in the proof of Theorem 6.1, we can derive (A1) and (A2) from (a1) and (6.6). Put $X_0 = H_0^1(\Omega) \cap L^\sigma(\Omega)$. Then (Φ 1) follows from the fact that $p < \max\{2^*, \sigma\}$ by (6.11). Here we observe by (i) of Remark 6.6 that $D(\varphi) \subset H_0^1(\Omega) \cap L^\sigma(\Omega) \subset L^{p'(q_1-1)}(\Omega) \cap W^{1,p'(q_2-1)}(\Omega) \subset D(B)$, and moreover,

$$|u|_{L^{p'(q_1-1)}(\Omega)}^{p'(q_1-1)} + |\nabla u|_{L^{p'(q_2-1)}(\Omega)}^{p'(q_2-1)} \leq C (\varphi(u) + |u|_V^p + 1) \quad (6.14)$$

for all $u \in D(\varphi)$. On the other hand, (a4) yields

$$|B(u)|_{V^*}^{p'} \leq C_{13} \left(|u|_{L^{p'(q_1-1)}(\Omega)}^{p'(q_1-1)} + |\nabla u|_{L^{p'(q_2-1)}(\Omega)}^{p'(q_2-1)} \right) + |a_2|_{L^1(\Omega)} \quad (6.15)$$

for $u \in L^{p'(q_1-1)}(\Omega) \cap W^{1,p'(q_2-1)}(\Omega)$. Thus (6.14) implies (B 1) $_{\varepsilon}$ with $\varepsilon = 0$. By Theorem 1.43 of [39] and (6.15), the function $\mathcal{B} : u \mapsto B(u(\cdot))$ is continuous from

$$\mathcal{X}_T := L^{p'(q_1-1)}(0, T; L^{p'(q_1-1)}(\Omega)) \cap L^{p'(q_2-1)}(0, T; W^{1,p'(q_2-1)}(\Omega))$$

into $L^{p'}(0, T; V^*)$, which particularly implies (B 3).

Finally, let us show (B 2). Due to Theorem 9.15 and Lemma 9.17 of [22], we can derive $D(\partial_V \varphi_1) = W^{2,p'}(\Omega) \cap H_0^1(\Omega)$ and

$$|u|_{W^{2,p'}(\Omega)} \leq C |\partial_V \varphi_1(u)|_{V^*} \quad \text{for all } u \in D(\partial_V \varphi_1).$$

Hence as in [2], we obtain

$$|u|_{W^{2,p'}(\Omega)} \leq C |\partial_V \varphi(u)|_{V^*} \quad \text{for all } u \in D(\partial_V \varphi). \quad (6.16)$$

We note that $D(\partial_V \varphi) \subset W^{2,p'}(\Omega) \cap H_0^1(\Omega) \cap L^\sigma(\Omega)$. By (ii) of Remark 6.6, it follows that $D(\partial_V \varphi)$ is compactly embedded in $L^{p'(q_1-1)}(\Omega) \cap W^{1,p'(q_2-1)}(\Omega)$. Hence by (6.16) and the Aubin-Lions-type compactness theorem (see, e.g., Theorem 5 of [37]), the sequence (u_n) of (B 2) is precompact in

$$L^{p'}(0, S; L^{p'(q_1-1)}(\Omega)) \cap L^{p'}(0, S; W^{1,p'(q_2-1)}(\Omega)).$$

Moreover, from (6.14) and the fact that

$$\sup_{t \in [0, S]} \left(\varphi(u_n(t)) + |u_n(t)|_V \right) \leq C,$$

the sequence (u_n) is bounded in $L^\infty(0, S; L^{p'(q_1-1)}(\Omega) \cap W^{1,p'(q_2-1)}(\Omega))$. Hence we can take a subsequence (n') of (n) such that

$$u_{n'} \rightarrow u \text{ strongly in } \mathcal{X}_S.$$

Therefore we infer that $\mathcal{B}(u_{n'}) \rightarrow \mathcal{B}(u)$ strongly in $L^{p'}(0, T; V^*)$ from the continuity of \mathcal{B} . Thus (B 2) holds. \square

Furthermore, the existence of global attractors for \mathcal{G} is also proved.

THEOREM 6.7. *Let Ω be a bounded domain of \mathbb{R}^N with smooth boundary $\partial\Omega$ and let $f \in L^{p'}(\Omega)$. In addition to (a1), (a2)', (a4), (6.11), (6.12) and (6.13), we suppose that*

$$p \leq \sigma$$

and

$$\begin{cases} \lambda \text{ is arbitrary} & \text{if } p'(q_1 - 1) < \sigma \text{ and } p'(q_2 - 1) < 2, \\ \lambda \text{ is small} & \text{otherwise.} \end{cases} \quad (6.17)$$

Then the generalized semiflow \mathcal{G} has a global attractor \mathcal{A} in X .

Proof. From the assumptions, it follows from (a2)' and (a4) that

$$\begin{aligned} & \langle \partial_V \varphi(u) + \lambda B(u), u \rangle \\ & \geq \int_{\Omega} |\nabla u(x)|^2 dx + \int_{\Omega} (j(u(x)) - j(0)) dx \\ & \quad - \lambda \int_{\Omega} |h(x, u(x), \nabla u(x))| |u(x)| dx \\ & \geq \int_{\Omega} |\nabla u(x)|^2 dx + C_8 \int_{\Omega} |u(x)|^\sigma dx - (C_9 + |j(0)|) |\Omega| \\ & \quad - \lambda \int_{\Omega} \left(C_{13} |u(x)|^{p'(q_1-1)} + C_{13} |\nabla u(x)|^{p'(q_2-1)} + a_2(x) \right)^{1/p'} |u(x)| dx \\ & \geq \frac{1}{2} \int_{\Omega} |\nabla u(x)|^2 dx + \frac{C_8}{2} \int_{\Omega} |u(x)|^\sigma dx - M_\lambda \end{aligned}$$

for $u \in D(\partial_V \varphi)$. From (i) of Remark 6.6 and (6.17), the last inequality follows, by assuming that $\lambda > 0$ is sufficiently small only in case

$$\text{either } q_1 = \sigma \text{ or } q_2 = \frac{2}{\sigma'} + 1. \quad (6.18)$$

Thus (S1) is satisfied, since $p \leq \sigma$.

In addition, assuming

$$p'(q_1 - 1) < \sigma \quad \text{and} \quad p'(q_2 - 1) < 2 \quad (6.19)$$

(then, (6.18) is not true), we have

$$|u|_{L^{p'(q_1-1)}(\Omega)}^{p'(q_1-1)} + |\nabla u|_{L^{p'(q_2-1)}(\Omega)}^{p'(q_2-1)} \leq C (\varphi(u) + |u|_V^p + 1)^r$$

with some $r \in (0, 1)$. Hence (B1) follows from (6.15).

Therefore Theorem 5.1 (respectively, Corollary 5.6) ensures that \mathcal{G} admits a global attractor in X for sufficiently small (respectively, arbitrary) $\lambda \in [0, 1]$ (respectively, if (6.19) holds). \square

Acknowledgements. The author expresses his sincere gratitude to Professors Tomomi Kojo and Tomoyuki Idogawa for fruitful discussions. Moreover, the author also thanks the anonymous referee for careful reading of this manuscript and for giving useful comments.

References

- [1] G. Akagi, On some doubly nonlinear parabolic equations, in: Current Advances in Nonlinear Analysis and Related Topics, GAKUTO Internat. Ser. Math. Sci. Appl., vol. 32, Gakkōtoshō, Tokyo, 2010, pp. 239–254.
- [2] G. Akagi, Maximality for the sum of two monotone operators in L^p -frameworks, submitted.
- [3] G. Akagi, Doubly nonlinear evolution equations with non-monotone perturbations in reflexive Banach spaces, submitted.
- [4] L. Ambrosio, N. Gigli, G. Savaré, *Gradient flows in metric spaces and in the space of probability measures*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, 2005.
- [5] T. Arai, On the existence of the solution for $\partial\varphi(u'(t)) + \partial\psi(u(t)) \ni f(t)$, *J. Fac. Sci. Univ. Tokyo Sec. IA Math.* **26** (1979), 75–96.
- [6] M. Aso, M. Frémond, N. Kenmochi, Quasi-variational evolution problems for irreversible phase change, in: Nonlinear partial differential equations and their applications, GAKUTO Internat. Ser. Math. Sci. Appl., vol. 20, Gakkōtoshō, Tokyo, 2004, pp. 517–525.
- [7] M. Aso, M. Frémond, N. Kenmochi, Phase change problems with temperature dependent constraints for the volume fraction velocities, *Nonlinear Anal.* **60** (2005), 1003–1023.
- [8] J.M. Ball, Continuity properties and global attractors of generalized semiflows and the Navier-Stokes equations, *J. Nonlinear Science* **7** (1997), 475–502.
- [9] J.M. Ball, Global attractors for damped semilinear wave equations, Partial differential equations and applications, *Discrete Contin. Dyn. Syst* **10** (2004), 31–52.

- [10] V. Barbu, *Nonlinear Semigroups and Differential Equations in Banach spaces*, Noordhoff, 1976.
- [11] V. Barbu, Existence theorems for a class of two point boundary problems, *J. Differential Equations* **17** (1975), 236–257.
- [12] A.V. Babin, M.I. Vishik, *Attractors of evolution equations*, Translated and revised from the 1989 Russian original by Babin, Studies in Mathematics and its Applications, 25. North-Holland Publishing Co., Amsterdam, 1992.
- [13] H. Brézis, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*, Math Studies, vol. 5 North-Holland, Amsterdam/New York, 1973.
- [14] H. Brézis, Monotonicity methods in Hilbert spaces and some applications to non-linear partial differential equations, in: E. Zarantonello (Eds.), Contributions to Nonlinear Functional Analysis, Academic Press, New York-London, 1971, pp. 101–156.
- [15] H. Brézis, M.G. Crandall, A. Pazy, Perturbations of nonlinear maximal monotone sets in Banach space, *Comm. Pure. Appl. Math.* **23** (1970), 123–144.
- [16] T. Caraballo, P. Marín-Rubio, J.C. Robinson, A comparison between two theories for multi-valued semiflows and their asymptotic behaviour, *Set-Valued Anal.* **11** (2003), 297–322.
- [17] V.V. Chepyzhov, M.I. Vishik, *Attractors for equations of mathematical physics*, American Mathematical Society Colloquium Publications, 49, American Mathematical Society, Providence, RI, 2002.
- [18] A. Cheskidov, Global attractors of evolutionary systems, *J. Dynam. Differential Equations* **21** (2009), 249–268.
- [19] A. Cheskidov, C. Foias, On global attractors of the 3D Navier-Stokes equations, *J. Differential Equations* **231** (2006), 714–754.
- [20] P. Colli, On some doubly nonlinear evolution equations in Banach spaces, *Japan J. Indust. Appl. Math.* **9** (1992), 181–203.
- [21] P. Colli, A. Visintin, On a class of doubly nonlinear evolution equations, *Comm. Partial Differential Equations* **15** (1990), 737–756.

- [22] D. Gilbarg, N.S. Trudinger, *Elliptic partial differential equations of second order*, Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001.
- [23] M.E. Gurtin, Generalized Ginzburg-Landau and Cahn-Hilliard equations based on a microforce balance, *Physica D* **92** (1996), 178–192.
- [24] J.K. Hale, *Asymptotic behavior of dissipative systems*, Mathematical Surveys and Monographs, 25, American Mathematical Society, Providence, RI, 1988.
- [25] O. Ladyzhenskaya, *Attractors for semigroups and evolution equations*, Cambridge University Press, Cambridge, 1991.
- [26] V.S. Melnik, J. Valero, On attractors of multivalued semi-flows and differential inclusions, *Set-Valued Anal.* **6** (1998), 83–111.
- [27] V.S. Melnik, J. Valero, On global attractors of multivalued semiprocesses and nonautonomous evolution inclusions, *Set-Valued Anal.* **8** (2000), 375–403.
- [28] A. Mielke, Evolution of rate-independent systems, in: Evolutionary equations, vol. II, Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam, 2005, pp. 461–559.
- [29] A. Mielke, F. Theil, On rate-independent hysteresis models, *NoDEA Nonlinear Differential Equations Appl.* **11** (2004), 151–189.
- [30] R. Rossi, A. Mielke, G. Savaré, A metric approach to a class of doubly nonlinear evolution equations and applications, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **7** (2008), 97–169.
- [31] R. Rossi, A. Segatti, U. Stefanelli, Attractors for gradient flows of non convex functionals and applications, *Arch. Ration. Mech. Anal.* **187** (2008), 91–135.
- [32] G. Schimperna, Global attractors for Cahn-Hilliard equations with non-constant mobility, *Nonlinearity* **20** (2007), 2365–2387.
- [33] G. Schimperna, A. Segatti, U. Stefanelli, Well-posedness and long-time behavior for a class of doubly nonlinear equations, *Discrete Contin. Dyn. Syst.* **18** (2007), 15–38.
- [34] A. Segatti, Global attractor for a class of doubly nonlinear abstract evolution equations, *Discrete Contin. Dyn. Syst.* **14** (2006), 801–820.

- [35] G.R. Sell, Differential equations without uniqueness and classical topological dynamics, *J. Differential Equations* **14** (1973), 42–56.
- [36] G.R. Sell, Global attractors for the three-dimensional Navier-Stokes equations, *J. Dynam. Differential Equations* **8** (1996), 1–33.
- [37] J. Simon, Compact sets in the space $L^p(0, T; B)$, *Ann. Math. Pura. Appl. (4)* **146** (1987), 65–96.
- [38] R. Temam, *Infinite-dimensional dynamical systems in mechanics and physics, Second edition*, Applied Mathematical Sciences, 68, Springer-Verlag, New York, 1997.
- [39] T. Roubíček, *Nonlinear partial differential equations with applications*, International Series of Numerical Mathematics, 153. Birkhäuser Verlag, Basel, 2005.
- [40] N. Yamazaki, Attractors for nonautonomous multivalued evolution systems generated by time-dependent subdifferentials, *Abstr. Appl. Anal.* **7** (2002), 453–473.
- [41] N. Yamazaki, Attractors of asymptotically periodic multivalued dynamical systems governed by time-dependent subdifferentials, *Electron. J. Differential Equations* (2004), no. 107, 22 pp.