



# Zonal spherical functions on the quantum homogeneous space $SU_q(n+1)/SU_q(n)$

Noumi, Masatoshi  
Yamada, Hirofumi  
Mimachi, Katsuhisa

---

(Citation)

Proceedings of the Japan Academy. Ser. A, Mathematical sciences, 65(6):169-171

(Issue Date)

1989-06

(Resource Type)

journal article

(Version)

Version of Record

(URL)

<https://hdl.handle.net/20.500.14094/90003238>



## 48. Zonal Spherical Functions on the Quantum Homogeneous Space $SU_q(n+1)/SU_q(n)$

By Masatoshi NOUMI,<sup>\*)</sup> Hirofumi YAMADA,<sup>\*\*)</sup>  
and Katsuhisa MIMACHI<sup>\*\*\*)</sup>

(Communicated by Shokichi IYANAGA, M. J. A., June 13, 1989)

In this note, we give an explicit expression to the zonal spherical functions on the quantum homogeneous space  $SU_q(n+1)/SU_q(n)$ . Details of the following arguments as well as the representation theory of the quantum group  $SU_q(n+1)$  will be presented in our forthcoming paper [3]. Throughout this note, we fix a non-zero real number  $q$ .

1. Following [4], we first make a brief review on the definition of the quantum groups  $SL_q(n+1; \mathbb{C})$  and its real form  $SU_q(n+1)$ .

The coordinate ring  $A(SL_q(n+1; \mathbb{C}))$  of  $SL_q(n+1; \mathbb{C})$  is the  $\mathbb{C}$ -algebra  $A = \mathbb{C}[x_{ij}; 0 \leq i, j \leq n]$  defined by the "canonical generators"  $x_{ij}$  ( $0 \leq i, j \leq n$ ) and the following fundamental relations:

$$(1.1) \quad x_{ik}x_{jk} = qx_{jk}x_{ik}, \quad x_{ki}x_{kj} = qx_{kj}x_{ki}$$

for  $0 \leq i < j \leq n$ ,  $0 \leq k \leq n$ ,

$$(1.2) \quad x_{il}x_{jk} = x_{jk}x_{il}, \quad x_{ik}x_{jl} - qx_{il}x_{jk} = x_{jl}x_{ik} - q^{-1}x_{jk}x_{il}$$

for  $0 \leq i < j \leq n$ ,  $0 \leq k < l \leq n$  and

$$(1.3) \quad \det_q = 1.$$

The symbol  $\det_q$  stands for the *quantum determinant*

$$(1.4) \quad \det_q = \sum_{\sigma \in S_{n+1}} (-q)^{l(\sigma)} x_{0\sigma(0)} x_{1\sigma(1)} \cdots x_{n\sigma(n)},$$

where  $S_{n+1}$  is the permutation group of the set  $\{0, 1, \dots, n\}$  and, for each  $\sigma \in S_{n+1}$ ,  $l(\sigma)$  denotes the number of pairs  $(i, j)$  with  $0 \leq i < j \leq n$  and  $\sigma(i) > \sigma(j)$ . This algebra  $A$  has the structure of a Hopf algebra, endowed with the *coproduct*  $\Delta: A \rightarrow A \otimes A$  and the *counit*  $\varepsilon: A \rightarrow \mathbb{C}$  satisfying

$$(1.5) \quad \Delta(x_{ij}) = \sum_{k=0}^n x_{ik} \otimes x_{kj} \quad \text{and} \quad \varepsilon(x_{ij}) = \delta_{ij} \quad \text{for } 0 \leq i, j \leq n.$$

Moreover, there exists a unique conjugate linear anti-homomorphism  $a \mapsto a^*: A \rightarrow A$  such that

$$(1.6) \quad x_{ji}^* = S(x_{ij}) \quad \text{for } 0 \leq i, j \leq n$$

with respect to the *antipode*  $S: A \rightarrow A$  of  $A$ . Together with this *\*-operation*, the Hopf algebra  $A = A(SL_q(n+1; \mathbb{C}))$  defines the *\*-Hopf algebra*  $A(SU_q(n+1))$ .

In what follows, we denote by  $G$  the quantum group  $SU_q(n+1)$  and by  $K$  the quantum subgroup  $SU_q(n)$  of  $G = SU_q(n+1)$ . Denote by  $y_{ij}$  ( $0 \leq i,$

<sup>\*)</sup> Department of Mathematics, Sophia University.

<sup>\*\*)</sup> Department of Mathematics, Tokyo Metropolitan University.

<sup>\*\*\*)</sup> Department of Mathematics, Nagoya University.

$j \leq n$ ) the canonical generators for the coordinate ring  $A(K)$ . Embedding of  $K$  into  $G$  is then specialized by the  $C$ -algebra epimorphism  $\pi_k: A(G) \rightarrow A(K)$  such that

$$(1.7) \quad \pi_K(x_{ij}) = y_{ij}, \quad \pi_K(x_{nn}) = 1 \quad \text{and} \quad \pi_K(x_{in}) = \pi_K(x_{nj}) = 0$$

for  $0 \leq i, j < n$ .

2. For a given dominant integral weight  $\lambda = \lambda_0 \varepsilon_0 + \cdots + \lambda_{n-1} \varepsilon_{n-1}$  ( $\lambda_0 \geq \cdots \geq \lambda_{n-1} \geq 0$ ), there exists a unique irreducible right  $A(G)$ -comodule  $V_\lambda$  with highest weight  $\lambda$ . We denote by  $\Lambda_k$  the fundamental weight  $\varepsilon_0 + \cdots + \varepsilon_{k-1}$  for  $1 \leq k \leq n$ . As a representation of  $K = SU_q(n)$ ,  $V_\lambda$  can be decomposed into irreducible components. It turns out that  $V_\lambda$  has a trivial representation of  $K$  as an irreducible component if and only if the highest weight  $\lambda$  is of the form  $\lambda = l\Lambda_1 + m\Lambda_n$  for some  $l, m \in \mathbb{N}$  and that the trivial representation may appear with multiplicity one. Such a representation  $V_\lambda$  is said to be of *class 1 relative to  $K$* .

If  $V_\lambda$  is of class 1, it can be decomposed into the form

$$(2.1) \quad V_\lambda = C v_0 \oplus V'_\lambda$$

as an  $A(K)$ -comodule, where  $v_0$  is a  $K$ -fixed vector of  $V_\lambda$ . Let  $\{v_1, \dots, v_{N-1}\}$  be a  $C$ -basis for  $V'_\lambda$  ( $N = \dim_C V_\lambda$ ) and define the matrix elements  $w_{ij}$  of the representation  $V_\lambda$  by

$$(2.2) \quad R_G(v_j) = \sum_{i=0}^{N-1} v_i \otimes w_{ij} \quad \text{for } 0 \leq j < N.$$

Here  $R_G: V_\lambda \rightarrow V_\lambda \otimes A(G)$  is the structure mapping of the right  $A(G)$ -comodule  $V_\lambda$ . Then the matrix element  $w_{00}$  does not depend on the choice of  $v_0, \dots, v_{N-1}$  and is *bi- $K$ -invariant* in the sense that

$$(2.3) \quad R_K(w_{00}) = w_{00} \otimes 1 \quad \text{and} \quad L_K(w_{00}) = 1 \otimes w_{00},$$

where

$$R_K = (id \otimes \pi_K) \circ \Delta \quad \text{and} \quad L_K = (\pi_K \otimes id) \circ \Delta.$$

We call  $w_{00}$  the *zonal spherical function* of  $V_\lambda$  relative to  $K$ .

3. We introduce the notation of quantum  $r$ -minor determinants. Let  $I$  and  $J$  be two subsets of  $\{0, 1, \dots, n\}$  with  $\#I = \#J = r$ . Arrange the elements of  $I$  and  $J$  in the increasing order:  $I = \{i_0, \dots, i_{r-1}\}$  ( $0 \leq i_0 < \cdots < i_{r-1} \leq n$ ) and  $J = \{j_0, \dots, j_{r-1}\}$  ( $0 \leq j_0 < \cdots < j_{r-1} \leq n$ ). We define the *quantum  $r$ -minor determinant*  $\xi_J^I$  by

$$(3.1) \quad \xi_J^I = \xi_{j_0 \dots j_{r-1}}^{i_0 \dots i_{r-1}} = \sum_{\sigma \in S_r} (-q)^{l(\sigma)} x_{i_0 j_{\sigma(0)}} x_{i_1 j_{\sigma(1)}} \cdots x_{i_{r-1} j_{\sigma(r-1)}}.$$

If  $I = \{0, 1, \dots, r-1\}$ , we use the abbreviation  $\xi_J = \xi_J^I$ .

To investigate spherical functions, we give a geometric realization of  $V_\lambda$  (cf. [3]).

Let  $\lambda = \lambda_{\mu_0} + \lambda_{\mu_1} + \cdots + \lambda_{\mu_{k-1}}$  ( $\mu_0 \geq \cdots \geq \mu_{k-1} > 0$ ) be a dominant integral weight. Let  $\mathbf{J} = (J_0, \dots, J_{k-1})$  be a sequence of non-empty subsets of  $\{0, 1, \dots, n\}$  with  $\#J_s = \mu_s$  for  $0 \leq s < k$ . We call  $\mathbf{J}$  a *column strict plane partition of shape  $\lambda$*  if the following conditions are satisfied:

$$(3.2) \quad \begin{cases} J_s = \{j_{0,s}, \dots, j_{\mu_s-1,s}\} \subset \{0, 1, \dots, n\}, \\ j_{r,s} < j_{r+1,s} \quad \text{and} \quad j_{r,s} \leq j_{r,s+1}. \end{cases}$$

The irreducible representation of  $V_\lambda$  can be realized as a right sub- $A(G)$ -

comodule as

$$(3.3) \quad V_\lambda = \sum_J C \xi_J \subset A(G),$$

where the sum is taken over the set of all column strict plane partitions of shape  $\lambda$  and

$$(3.4) \quad \xi_J = \xi_{J_0} \cdots \xi_{J_{k-1}} \quad \text{if } J = (J_0, \dots, J_{k-1}).$$

It is seen that these vectors  $\xi_J$  are linearly independent and that they form a  $C$ -basis for a right  $A(G)$ -comodule. Note that the product of minor determinants

$$(3.5) \quad \xi_{01 \dots \mu_0 - 1} \cdots \xi_{01 \dots \mu_{k-1} - 1}$$

gives the highest weight vector of  $V_\lambda$ . We remark that  $V_\lambda$  is identified with the vector space of all left relative  $B_-$ -invariants in  $A(SL_q(n+1; C))$  with respect to the character corresponding to  $\lambda$ , where  $B_-$  is the Borel subgroup "of lower triangular matrices" (see [3]).

If  $\lambda = l\lambda_1 + m\lambda_n$  ( $l, m \in N$ ), the spherical representation  $V_\lambda$  contains a  $K$ -fixed vector

$$(3.6) \quad v_0 = (\xi_n^0)^l (\xi_{01 \dots n-1}^{01 \dots n-1})^m.$$

By using algebraic properties of quantum minor determinants, we can determine explicitly the zonal spherical function of  $V_\lambda$  in terms of *basic hypergeometric series*  ${}_2\varphi_1$ :

$$(3.7) \quad {}_2\varphi_1 \left( \begin{matrix} a, b \\ c \end{matrix}; q, z \right) = \sum_{k=0}^{\infty} \frac{(a; q)_k (b; q)_k}{(c; q)_k (q; q)_k} z^k, \quad (a; q)_k = \prod_{j=0}^{k-1} (1 - aq^j).$$

**Theorem.** *Let  $V_\lambda$  be the representation of  $G = SU_q(n+1)$  with highest weight  $\lambda = l\lambda_1 + m\lambda_n$  ( $l, m \in N$ ). Then the zonal spherical function  $w = w_{v_0}$  of  $V_\lambda$  relative to  $K = SU_q(n)$  is expressed by a basic hypergeometric series in  $z = 1 - x_{nn} \xi_{01 \dots n-1}^{01 \dots n-1}$  as follows:*

$$(3.8) \quad w = (x_{nn})^{l-m} {}_2\varphi_1 \left( \begin{matrix} q^{-2m}, q^{2(l+n)} \\ q^{2n} \end{matrix}; q^2, q^2 z \right) \quad \text{if } l \geq m,$$

and

$$(3.9) \quad w = {}_2\varphi_1 \left( \begin{matrix} q^{-2l}, q^{2(m+n)} \\ q^{2n} \end{matrix}; q^2, q^2 z \right) (\xi_{01 \dots n-1}^{01 \dots n-1})^{m-l} \quad \text{if } l \leq m.$$

The above polynomials in  $z$  are so-called *little  $q$ -Jacobi polynomials*. As for *zonal* spherical functions, Theorem generalizes a result of [1, 2].

## References

- [1] T. Masuda *et al.*: Representations of quantum groups and a  $q$ -analogue of orthogonal polynomials. C. R. Acad. Sci. Paris, Série I, **307**, 559–564 (1988).
- [2] —: Representations of the quantum group  $SU_q(2)$  and the little  $q$ -Jacobi polynomials (preprint).
- [3] M. Noumi, H. Yamada, and K. Mimachi: Representations of the quantum group  $GL_q(n+1; C)$  (in preparation).
- [4] N. Yu. Reshetikhin, L. A. Takhtajan, and L. D. Faddeev: Quantization of Lie groups and Lie algebras (to appear in Algebra and Analysis) (in Russian).