



Diagnosis of desmoplastic small-round-cell tumor by cytogenetic analysis: a case report

Wakahashi, Senn ; Sudo, Tamotsu ; Ichida, Kotaro ; Sugita, Shintaro ; Hasegawa, Tadashi ; Nagao, Shoji ; Yamaguchi, Satoshi ; Sakuma, Toshik...

(Citation)

Clinical Case Reports, 4(5):520-523

(Issue Date)

2016-05

(Resource Type)

journal article

(Version)

Version of Record

(Rights)

©2016 The Authors. Clinical Case Reports published by John Wiley & Sons Ltd.
This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no...

(URL)

<https://hdl.handle.net/20.500.14094/90003437>



CASE REPORT

Diagnosis of desmoplastic small-round-cell tumor by cytogenetic analysis: a case report

Senn Wakahashi¹, Tamotsu Sudo^{2,3}, Kotaro Ichida¹, Shintaro Sugita⁴, Tadashi Hasegawa⁴, Shoji Nagao², Satoshi Yamaguchi², Toshiko Sakuma⁵ & Hideto Yamada¹

¹Department of Obstetrics and Gynecology, Kobe University Graduate School of Medicine, Kobe, Japan

²Department of Gynecologic Oncology, Hyogo Cancer Center, Akashi, Japan

³Section of Translational Research, Hyogo Cancer Center, Akashi, Japan

⁴Department of Surgical Pathology, Sapporo Medical University School of Medicine, Sapporo, Japan

⁵Department of Diagnostic Pathology, Hyogo Cancer Center, Akashi, Japan

Correspondence

Senn Wakahashi, Department of Obstetrics and Gynecology, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-Ku, Kobe, Hyogo 650-0017, Japan. Tel: +81-78-382-6005; Fax: +81-78-382-6019; E-mail: wsenn_0120@chic.ocn.ne.jp

Funding Information

No sources of funding were declared for this study.

Received: 6 December 2015; Accepted: 20 March 2016

Clinical Case Reports 2016; 4(5): 520–523

doi: 10.1002/ccr3.558

Key Clinical Message

We herein present atypical histologic and immunohistochemical features of DSRCT. The various differential diagnoses of DSRCT may occasionally generate confusion. Cytogenetic analysis may solve diagnostic dilemmas such as that in our case. Further studies are required to establish a standard treatment for DSRCT.

Keywords

Chromosomal translocation, cytogenetic analysis, diagnostic dilemma, desmoplastic small-round-cell tumor.

Introduction

Desmoplastic small-round-cell tumor (DSRCT) is a rare, highly aggressive intra-abdominal neoplasm with an extremely poor prognosis, first described in 1989 [1]. It frequently occurs in the abdominal cavity and pelvis with an undetermined histologic origin, and its incidence in males (>80%) and females differs [2–6]. In one comprehensive study, the typical immunohistochemical features were epithelial marker positivity (cytokeratin, 91%; epithelial membrane antigen [EMA], 88%), mesenchymal marker positivity (desmin, 91%; vimentin, 84%), and various results for neural antigens [6]. Atypical histologic and immunohistochemical features can pose a diagnostic dilemma. We herein report a case of epithelial marker – negative DSRCT, only a few cases of which have been reported worldwide. Cytogenetic profiles could provide useful diagnostic information in such cases.

Case Report

A 22-year-old woman (gravida 0, parity 0) presented with abdominal distension. A physical examination revealed a bulky abdominal tumor. Magnetic resonance imaging showed a mass in the pelvic cavity, and computed tomography showed a large, complex, heterogeneously enhancing mass invading the abdominal cavity and displacing the ovary, colon, and small intestine (Fig. 1). The serum CA-125 level was elevated at 285.3 U/mL (normal, <35 U/mL). The serum CA19-9, carcinoembryonic antigen, and alpha-fetoprotein levels were within the normal range.

Probe laparotomy only with tumor biopsy was performed to determine the diagnosis. During laparotomy, the mass appeared soft and immobile, nontender, and very vascular (Fig. 2). Immediately after laparotomy, the patient presented with acute progression of symptoms such as ascites and pelvic and abdominal tenderness. Early



Figure 1. Computed tomography shows a large heterogeneously enhancing mass invading the abdominal cavity.




Figure 2. Intraoperative image of the multilobulated mass.

therapeutic intervention, without waiting for pathological examination of a biopsy specimen, was necessary for this patient. Frozen biopsy was performed, and the results were suspicious for a mesenchymal tumor; therefore, the patient underwent chemotherapy with doxorubicin and ifosfamide.

A pathological examination showed diffuse infiltration with a necrotic area of short spindle cells and small round cells with irregular nuclei (Fig. 3A). The mitotic rate was high (60 per 10 HPF).

Immunohistochemical analysis was performed using antibodies directed against epithelial markers (EMA, cytokeratin 7, cytokeratin 20, CAM5.2, and AE1/3), mesenchymal markers (desmin and vimentin) (Fig. 3B,C),

and neural markers (synaptophysin and s-100 protein). Unfortunately, the atypical immunohistochemical features made diagnosis of this tumor difficult. The tumor cells were positive when stained with antibodies to EMA, vimentin, desmin, and synaptophysin, but were negative for s-100 protein and some epithelial markers (cytokeratin 7, cytokeratin 20, CAM5.2, and AE1/3).

Cytogenetic analysis was necessary for a definitive diagnosis in this case. In our case, detection of the chromosomal translocation $t(11;22)(p13;q12)$ allowed for diagnosis by fluorescence in situ hybridization [7] and polymerase chain reaction (PCR).

PCR revealed fusion of the Ewing sarcoma gene (*EWS*) and Wilms tumor gene (*WT1*), and break-apart FISH detected translocations in the *EWS* gene (Fig. 4A,B). PCR and FISH were carried out according to the manufacturer's instructions, as described previously [8, 9].

The patient underwent treatment with five cycles of VDC regimens (vincristine, doxorubicin, and cyclophosphamide) and three cycles of VAC regimens (vincristine, actinomycin D, and cyclophosphamide) after we established the definitive diagnosis. However, the patient's disease progressed throughout all three cycles of VAC regimens, and she died of her disease 9 months after her initial diagnosis.

Discussion

The differential diagnosis of small round cell tumor includes Ewing sarcoma, embryonal rhabdomyosarcoma, Wilms tumor, neuroblastoma, medulloblastoma, synovial sarcoma, and lymphoma; these tumors are often confused with one another [10]. A diagnosis of DSRCT is usually made by a combination of the histologic appearance and immunohistochemical staining results. Therefore, atypical histologic and immunohistochemical features can pose a diagnostic dilemma. Generally, DSRCTs show the typical histologic features of a large necrotic area, sharply demarcated nests of small round cells, or spindle cells embedded in a desmoplastic stroma. The small nests of cells are variably sized and contain hyperchromatic nuclei with faintly eosinophilic, scanty cytoplasm [3, 11–13]. Most DSRCTs coexpress epithelial markers (cytokeratin, EMA), mesenchymal markers (desmin, vimentin), and neural markers (synaptophysin, s-100 protein) [3]. In our case, however, immunohistochemical staining disclosed negativity for some epithelial markers (cytokeratin 7, cytokeratin 20, CAM5.2, and AE1/3) and EMA positivity. In contrast, expression of mesenchymal markers (desmin and vimentin) was positive. In particular, the desmin positivity made it possible to diagnose a malignant myometrial tumor. Embryonal rhabdomyosarcoma, synovial sarcoma, ES/PNET, and an unusual type of leiomyosarcoma should be




Figure 3. (A) The tumor shows diffusely infiltrating short spindle cells and small round cells ($\times 200$). (B) Immunohistochemical staining for desmin shows a dot-like perinuclear reaction in the tumor cells ($\times 400$) and (C) positive staining with vimentin ($\times 200$).




Figure 4. (A) Fluorescence in situ hybridization using a break-apart probe for the Ewing sarcoma breakpoint region 1 gene is positive. (B) *EWS/WT1* rearrangement reverse-transcription PCR demonstrates the 268-bp product.

considered in such cases. The exact histogenesis was unclear in the present case.

Definitive diagnosis was achieved in our case by cytogenetic analysis. DSRCT is generally characterized by a chromosomal translocation $t(11;22)(p13;q12)$ that results in the fusion of the genes *EWS* and *WT1* [7, 11, 13–15].

DSRCTs that are negative for epithelial markers are relatively rare, and only three such cases have been reported. Cytogenetic analysis may resolve diagnostic dilemmas such as that in our case [11, 16, 17].

With respect to treatment of DSRCT, chemotherapy with an intensive alkylator-based regimen is associated with better survival than is standard-dose chemotherapy. This regimen has a risk of toxicity that requires intensive transfusions and antibiotic support [4, 18]. Complete surgical resection is also associated with improved survival. However, debulking surgery is usually impossible at an

advanced stage [4, 19, 20]. Despite such aggressive therapy, the outcome is poor. In one study, 25 of 35 patients died of widespread metastases within a mean of 25.2 months from the time of their diagnosis [21]. Although our patient received chemotherapy using VDC and VAC, she finally died of local tumor progression.

Our patient presented with atypical histologic and immunohistochemical features of DSRCT. The various differential diagnoses of DSRCT may occasionally generate confusion. Cytogenetic analysis may solve diagnostic dilemmas such as that in our case. Further studies are required to establish a standard treatment for DSRCT.

Conflict of Interest

None declared.

References

- Gerald, W. L., and J. Rosai. 1989. Case 2. Desmoplastic small cell tumor with divergent differentiation. *Pediatr. Pathol.* 9:177–183.
- Gerald, W. L., H. K. Miller, H. Battifora, M. Miettinen, E. G. Silva, and J. Rosai. 1991. Intra-abdominal desmoplastic small round-cell tumor. Report of 19 cases of a distinctive type of high-grade polyphenotypic malignancy affecting young individuals. *Am. J. Surg. Pathol.* 15:499–513.
- Ordóñez, N. G. 1998a. Desmoplastic small round cell tumor: II: an ultrastructural and immunohistochemical study with emphasis on new immunohistochemical markers. *Am. J. Surg. Pathol.* 22:1314–1327.
- Lal, D. R., W. T. Su, S. L. Wolden, K. C. Loh, S. Modak, and M. P. La Quaglia. 2005. Results of multimodal treatment for desmoplastic small round cell tumors. *J. Pediatr. Surg.* 40:251–255.
- Gerald, W. L., M. Ladanyi, E. de Alava, M. Cuatrecasas, B. H. Kushner, M. P. LaQuaglia, et al. 1998. Clinical, pathologic, and molecular spectrum of tumors associated with t(11;22)(p13;q12): desmoplastic small round-cell tumor and its variants. *J. Clin. Oncol.* 16:3028–3036.
- Goldblum, J. R., S. W. Weiss, and A. L. Folpe. 2013. Enzinger and Weiss's soft tissue tumors: expert consult: online. Elsevier Health Sciences, Philadelphia; 2013.
- Zhang, P. J., J. R. Goldblum, B. R. Pawel, C. Fisher, T. L. Pasha, and F. G. Barr. 2003. Immunophenotype of desmoplastic small round cell tumors as detected in cases with EWS-WT1 gene fusion product. *Modern Pathol.* 16:229–235.
- Yamaguchi, U., T. Hasegawa, Y. Morimoto, U. Tateishi, M. Endo, F. Nakatani, et al. 2005. A practical approach to the clinical diagnosis of Ewing's sarcoma/primitive neuroectodermal tumour and other small round cell tumours sharing EWS rearrangement using new fluorescence in situ hybridisation probes for EWSR1 on formalin fixed, paraffin wax embedded tissue. *J. Clin. Pathol.* 58:1051–1056.
- Miura, Y., Y. Keira, J. Ogino, K. Nakanishi, H. Noguchi, T. Inoue, et al. 2012. Detection of specific genetic abnormalities by fluorescence in situ hybridization in soft tissue tumors. *Pathol. Int.* 62:16–27.
- Reynolds, C. P., R. G. Smith, and E. P. Frenkel. 1981. The diagnostic dilemma of the "small round cell neoplasm": catecholamine fluorescence and tissue culture morphology as markers for neuroblastoma. *Cancer* 48:2088–2094.
- Zhang, J., J. Dalton, and C. Fuller. 2007. Epithelial marker-negative desmoplastic small round cell tumor with atypical morphology: definitive classification by fluorescence in situ hybridization. *Arch. Pathol. Lab. Med.* 131:646–649.
- Lae, M. E., P. C. Roche, L. Jin, R. V. Lloyd, and A. G. Nascimento. 2002. Desmoplastic small round cell tumor: a clinicopathologic, immunohistochemical, and molecular study of 32 tumors. *Am. J. Surg. Pathol.* 26: 823–835.
- Guinee, D. G., and T. C. Allen. 2008. Primary pleural neoplasia: entities other than diffuse malignant mesothelioma. *Arch. Pathol. Lab. Med.* 132:1149–1170.
- Yang, Y., H. Wang, Y. P. Wang, W. Zhang, Y. Z. Hui, Z. H. Wang, et al. 2005. [Detection of EWS-WT1 fusion transcripts in paraffin-embedded tissues for desmoplastic small round cell tumor]. *Beijing da xue xue bao Yi xue ban = Journal of Peking University Health sciences* 37:325–328.
- Gerald, W. L., J. Rosai, and M. Ladanyi. 1995. Characterization of the genomic breakpoint and chimeric transcripts in the EWS-WT1 gene fusion of desmoplastic small round cell tumor. *Proc. Natl Acad. Sci. USA* 92:1028–1032.
- Trupiano, J. K., S. K. Machen, F. G. Barr, and J. R. Goldblum. 1999. Cytokeratin-negative desmoplastic small round cell tumor: a report of two cases emphasizing the utility of reverse transcriptase-polymerase chain reaction. *Modern Pathol.* 12:849–853.
- Dorsey, B. V., L. E. Benjamin, F. Rauscher, B. Klencke, A. P. Venook, R. S. Warren, et al. 1996. Intra-abdominal desmoplastic small round-cell tumor: expansion of the pathologic profile. *Modern Pathol.* 9:703–709.
- Kushner, B. H., M. P. LaQuaglia, N. Wollner, P. A. Meyers, K. L. Lindsley, F. Ghavimi, et al. 1996. Desmoplastic small round-cell tumor: prolonged progression-free survival with aggressive multimodality therapy. *J. Clin. Oncol.* 14:1526–1531.
- Talarico, F., D. Iusco, L. Negri, and D. Belinelli. 2007. Combined resection and multi-agent adjuvant chemotherapy for intra-abdominal desmoplastic small round cell tumour: case report and review of the literature. *Il Giornale di chirurgia* 28:367–370.
- Liu, Q., N. Liu, and D. Chen. 2014. Primary desmoplastic small round cell tumor of the duodenum. *Eur. J. Med. Res.* 19:38.
- Ordóñez, N. G. 1998b. Desmoplastic small round cell tumor: I: a histopathologic study of 39 cases with emphasis on unusual histological patterns. *Am. J. Surg. Pathol.* 22:1303–1313.