



The central limit theorem for lacunary series

Fukuyama, Katusi
Takahashi, Shigeru

(Citation)

Proceedings of the American Mathematical Society, 127(2):599-608

(Issue Date)

1999-02

(Resource Type)

journal article

(Version)

Accepted Manuscript

(Rights)

First published in Proceedings of the American Mathematical Society 127(2) (Feb 1999), published by the American Mathematical Society. ©1999 American Mathematical Society.

(URL)

<https://hdl.handle.net/20.500.14094/90003853>



THE CENTRAL LIMIT THEOREM FOR LACUNARY SERIES

KATUJI FUKUYAMA AND SHIGERU TAKAHASHI

???, ?, 1997

ABSTRACT. In this paper, the central limit theorem for lacunary trigonometric series is proved. Two gap conditions by Erdős and Takahashi are extended and unified. The criterion for the Fourier character of lacunary series is also given.

1. INTRODUCTION

It is well known that lacunary trigonometric series $\sum a_i \cos(2\pi n_i \omega + \phi_i)$ behaves like random series when $\{n_i\}$ increases very fast. For example, if $\{n_i\}$ has Hadamard gaps, i.e. $n_{i+1}/n_i > q > 1$, the series converges or diverges almost everywhere according as $\sum a_i^2$ converges or diverges. (Kolmogorov [3] and Zygmund [10].) It is also known that the series is not a Fourier series of integrable function when $\sum a_i^2 = \infty$. (Zygmund [10].)

As to the central limit theorem for the series with Hadamard gaps, Salem-Zygmund [4] proved: If $A_n = (\frac{1}{2} \sum_{i=1}^n a_i^2)^{1/2} \rightarrow \infty$ and $a_n = o(A_n)$ are satisfied, then

$$(1.1) \quad \frac{1}{A_n} \sum_{i=1}^n a_i \cos(2\pi n_i \omega + \phi_i) \xrightarrow{\mathcal{D}} N_{0,1}$$

holds on probability space $(\Omega, d\omega/|\Omega|)$, when $\Omega \subset [0, 1]$ has positive measure. Here $|\cdot|$ denotes Lebesgue measure, $N_{0,1}$ the standard normal distribution, and $\xrightarrow{\mathcal{D}}$ convergence in law.

Erdős [1] relaxed the gap condition to

$$(1.2) \quad n_{i+1}/n_i > 1 + c_i/\sqrt{i} \quad \text{where} \quad c_i \rightarrow \infty,$$

and proved (1.1) for $a_n \equiv 1$. Takahashi [6] proved that $a_n \equiv 1$ can be relaxed to

$$(1.3) \quad A_n \rightarrow \infty \quad \text{and} \quad a_n = O(A_n/\sqrt{n}).$$

Takahashi [7] also proved (1.1) assuming

$$(1.4) \quad n_{i+1}/n_i > 1 + c/i^\alpha, \quad A_n \rightarrow \infty \quad \text{and} \quad a_n = o(A_n/n^\alpha),$$

1991 *Mathematics Subject Classification*. Primary 60F05, 42A55.

Key words and phrases. Lacunary series, Central limit theorem, Fourier series, Fourier Stieltjes series.

where $c > 0$ and $0 \leq \alpha \leq 1/2$. It should be noted that there is no implication between (1.2) and (1.3), and (1.4). Indeed, if we put $\alpha = 1/2$, the gap condition of (1.4) is weaker than (1.2), but if we put $a_n \equiv 1$, we must put $\alpha < 1/2$ in (1.4), which is stronger than (1.2).

From (1.1), by the way, we can deduce that the series is not a Fourier series. Therefore, the series is not a Fourier series under (1.4). Takahashi [8] proved that this claim remains valid even if we relax $a_n = o(A_n/n^\alpha)$ to $a_n = O(A_n/n^\alpha)$. Under this condition, (1.1) does not hold generally. A counterexample was constructed by Takahashi [9]. Previously, Erdős [1] had noted the existence of such an example for $\alpha = 1/2$.

Although these results have been considered to be best possible, we still have the following examples excluded from the above scheme: Under the conditions

$$n_{i+1}/n_i > 1 + c/\sqrt{i \log i}, \quad A_n \rightarrow \infty \quad \text{and} \quad a_n = o(A_n/\sqrt{n \log n}),$$

the central limit theorem (1.1) holds. Even if we relax the last condition to $a_n = O(A_n/\sqrt{n \log n})$, the series is not a Fourier series, but there are counterexamples for (1.1).

In this note we introduce a more general gap condition and prove theorems including all the above results and examples.

Theorem 1. *Let us suppose the following conditions:*

$$(1.5) \quad \lambda(i) > \lambda \quad \text{for some } \lambda > 0;$$

$$(1.6) \quad \lambda(i+1) - \lambda(i) = o(1);$$

$$(1.7) \quad n_{i+1}/n_i > 1 + c/\lambda(i) \quad \text{for some } c > 0;$$

$$(1.8) \quad a_n = o(A_n/\lambda(n)) \quad \text{and} \quad A_n \rightarrow \infty.$$

Then (1.1) holds on any $\Omega \subset [0, 1]$ with positive measure.

In Theorem 1, the next condition is implicitly assumed:

$$(1.9) \quad \sum_{i=1}^{\infty} \frac{1}{\lambda^2(i)} = \infty.$$

Or, more precisely, the existence of $\{a_i\}$ satisfying (1.8) is equivalent to (1.9). Indeed, if (1.9) is false, the contradiction $A_n^2 = O(\frac{1}{2} \sum_{i=1}^n A_i^2/\lambda^2(i)) = o(A_n^2)$ follows. In case (1.9) is valid, we can construct $\{a_i\}$ satisfying (1.8), by putting $A_n = \exp(\frac{1}{2} \sum_{i=1}^n e_i)$ and $a_n^2 = (A_n^2 - A_{n-1}^2)/2$, where $\{e_i\}$ satisfies $0 < e_i = o(1/\lambda^2(i))$ and $\sum_{i=1}^{\infty} e_i = \infty$.

Although Theorem 1 contains assumptions that generalize (1.4), we can derive the next corollary which assumes the generalization of (1.2) and (1.3). The condition (1.9) is again implicitly assumed for $\{\Lambda(i)\}$.

Corollary. *Let us suppose the following conditions:*

$$\Lambda(i) > 0 \quad \text{and} \quad \Lambda(i) \rightarrow \infty;$$

$$\Lambda(i+1) - \Lambda(i) = O(1);$$

$$n_{i+1}/n_i > 1 + c_i/\Lambda(i) \quad \text{where } c_i \rightarrow \infty;$$

$$a_n = O(A_n/\Lambda(n)) \quad \text{and} \quad A_n \rightarrow \infty.$$

Then (1.1) holds on any $\Omega \subset [0, 1]$ with positive measure.

Actually, we will prove that the assumption of Theorem 1 is equivalent to that of Corollary. Thus, our result extends and unifies previous results.

Since $a_n = o(A_n)$ is necessary for (1.1) (Salem-Zygmund [4]), if we assume (1.8), the condition (1.5) is indispensable for Theorem 1.

The next theorem asserts that (1.8) cannot be weakened.

Theorem 2. *Suppose that $\{\lambda(i)\}$ satisfies $\lambda(i) > 0$, (1.6) and (1.9). Then there exist sequences $\{n_i\}$ and $\{a_i\}$ satisfying (1.7),*

$$(1.10) \quad a_n = O(A_n/\lambda(n)) \quad \text{and} \quad A_n \rightarrow \infty,$$

such that the central limit theorem (1.1) does not hold on $\Omega = [0, 1]$.

Finally, we state a result on the Fourier character of the series. Since (1.11) is always true under Hadamard's gap condition, it includes that of Zygmund [10].

Theorem 3. *Let us assume (1.5), (1.6), (1.7), (1.9),*

$$(1.11) \quad a_n = O(A_n \ell(A_n)/\lambda(n)) \quad \text{and} \quad A_n \rightarrow \infty,$$

where $\ell(x) = \sqrt{\log x \log \log x \dots \log^{(\gamma)} x}$ and $\gamma \in \mathbf{N}$. Then the series diverges almost surely and is neither a Fourier series nor a Fourier-Stieltjes series.

Before closing the introduction, we note that the same results for lacunary Walsh series can be proved in the same way.

2. THE CENTRAL LIMIT THEOREM

Let us put $n_0 = 1$ and $\lambda(0) = 2\lambda$, and introduce the following notation:

$$\begin{aligned} p(0) &= 0, & p(k) &= \max\{i \mid n_i \leq 2^k\} \quad (k \geq 1), & l(k) &= p(k+1) - p(k), \\ P(k) &= \mathbf{N} \cap (p(k), p(k+1)], & \mu(k) &= \max_{i \in P(k)} \lambda(i), & \nu(k) &= \min_{i \in P(k)} \lambda(i). \end{aligned}$$

Since $\{n_i\}$ diverges to infinity, $\{p(k)\}$ does also. If $p(k) + 1 < p(k+1)$, we have

$$2 > \frac{n_{p(k+1)}}{n_{p(k)+1}} > \prod_{i=p(k)+1}^{p(k+1)-1} \left(1 + \frac{c}{\lambda(i)}\right) > 1 + \sum_{i=p(k)+1}^{p(k+1)-1} \frac{c}{\lambda(i)} > 1 + c \frac{l(k) - 1}{\mu(k)}.$$

From this and $\mu(k) > \lambda$, it follows that

$$(2.1) \quad l(k) = O(\mu(k)).$$

By (1.6), we have $\lambda(i) = o(i)$ and $\mu(k) = o(p(k+1))$. Applying this to (2.1) we get

$$(2.2) \quad p(k+1) \sim p(k).$$

Applying (1.6) and (2.1), we have

$$0 \leq \frac{\mu(k) - \nu(k)}{\mu(k)} \leq \sum_{i=p(k)}^{p(k+1)-1} \frac{|\lambda(i+1) - \lambda(i)|}{\mu(k)} = o\left(\frac{l(k)}{\mu(k)}\right) = o(1).$$

This implies $\mu(k) \sim \nu(k)$, and hence $\mu(k) \sim \lambda(p(k+1))$ and $\mu(k+1) \sim \lambda(p(k+1)+1)$ follow. Since $\lambda(i+1) \sim \lambda(i)$ is clear from (1.5) and (1.6), we have

$$(2.3) \quad \mu(k+1) \sim \mu(k).$$

The next lemma plays an important role in the proof.

Lemma 1. *For any given integers j, k, h , and q satisfying*

$$j < k \quad \text{and} \quad p(j) + 1 < h \leq p(j+1) < p(k) + 1 < q \leq p(k+1),$$

the number of solutions (n_r, n_i) of the equation

$$n_q - n_r = n_h - n_i \quad \text{where} \quad p(j) < i < h \quad \text{and} \quad p(k) < r < q,$$

is at most $2^{j-k+1}\mu(k)/c$.

Proof. If (n_r, n_i) is a solution, we have

$$n_r = n_q - (n_h - n_i) > n_q - 2^j > n_q(1 - 2^{j-k}) \geq n_q(1 + 2^{j-k+1})^{-1}.$$

Let us denote the least (or greatest) index of n_r 's by m_1 (or m_2). Dividing $n_q \geq n_{m_2+1}$ by $n_q(1 + 2^{j-k+1})^{-1} \leq n_{m_1}$, we have

$$1 + 2^{j-k+1} \geq \frac{n_{m_2+1}}{n_{m_1}} \geq \prod_{m=m_1}^{m_2} \left(1 + \frac{c}{\lambda(m)}\right) \geq 1 + \frac{c(m_2 - m_1 + 1)}{\mu(k)}. \quad \square$$

The next lemma can be proved in the same way.

Lemma 2. *For any given integers j, k, h , and q satisfying*

$$j+1 < k \quad \text{and} \quad p(j+1) < h \leq p(j+2) < p(k+1) < q \leq p(k+2),$$

the number of solutions (n_r, n_i) of the equation

$$n_q - n_r = n_h - n_i \quad \text{where} \quad p(j) < i \leq p(j+1) \quad \text{and} \quad p(k) < r \leq p(k+1),$$

is at most $2^{j-k+2}\mu(k)/c$.

In the proof of Theorem 1, we assume $\phi_i = 0$ to simplify notation. The general case can be proved in the same way. We apply the following result ([5]).

Theorem A. *Let $\{d_i\}$ be a sequence of real numbers and put*

$$f_n(\omega) = \sum_{i=1}^n d_i \cos 2\pi i\omega, \quad \Delta_k = f_{2^{k+1}} - f_{2^k}, \quad D_n = \left(\frac{1}{2} \sum_{i=1}^n d_i^2 \right)^{1/2},$$

and $B_k = D_{2^{k+1}}$. Suppose that the following conditions are satisfied:

$$(2.4) \quad \int_0^1 \left| \frac{1}{B_n^2} \sum_{k=1}^n (\Delta_k^2(\omega) + 2\Delta_k(\omega)\Delta_{k+1}(\omega)) - 1 \right| d\omega \rightarrow 0,$$

$$(2.5) \quad B_n \rightarrow \infty \quad \text{and} \quad \sup_{\omega \in [0,1]} |\Delta_n(\omega)| = o(B_n).$$

Then for any $\Omega \subset [0, 1]$ with positive measure, the law of f_n/D_n on $(\Omega, d\omega/|\Omega|)$ converges weakly to the standard normal distribution.

We apply this by putting $\Delta_k(\omega) = \sum_{i \in P(k)} a_i \cos 2\pi n_i \omega$ and $B_k = A_{p(k+1)}$. Let us put $C_k = \|\Delta_k\|$ where $\|\cdot\|$ denotes $L^2[0,1]$ -norm. Obviously we have $C_k^2 = B_k^2 - B_{k-1}^2$. By (1.8) and (2.1), we have

$$(2.6) \quad \sup_{\omega \in [0,1]} |\Delta_k(\omega)| \leq \sum_{i \in P(k)} |a_i| \leq l(k) \max_{i \in P(k)} |a_i| = o\left(\mu(k) \frac{B_k}{\nu(k)}\right) = o(B_k).$$

This implies $C_k = o(B_k)$ and $B_{k+1}^2/B_k^2 = (1 - C_{k+1}^2/B_{k+1}^2)^{-1} = 1 + o(1)$. The next estimate follows from the Schwarz inequality:

$$(2.7) \quad \sum_{q \in P(k)} |a_q| \leq l^{1/2}(k) C_k = O(\mu^{1/2}(k) C_k).$$

To prove (2.4), we divide Δ_k^2 into three parts; putting

$$U_k = \frac{1}{2} \sum_{q \in P(k)} a_q \sum_{r \in P(k)} a_r \cos 2\pi(n_q + n_r)\omega \quad \text{and} \\ V_k = \sum_{p(k) < r < q \leq p(k+1)} a_q a_r \cos 2\pi(n_q - n_r)\omega,$$

we have $\Delta_k^2 - C_k^2 = U_k + V_k$. From (2.6), it follows that

$$\|U_k\| \leq \frac{1}{2} \sum_{q \in P(k)} |a_q| C_k = o(B_k C_k) \quad \text{and} \quad \|V_k\| = o(B_k C_k).$$

Since $\{U_k\}$ is an orthogonal sequence, we have

$$\left\| \sum_{k=1}^n U_k \right\|^2 = \sum_{k=1}^n \|U_k\|^2 = o\left(B_k^2 \sum_{k=1}^n C_k^2\right) = o(B_k^4).$$

Noting this and $\|\sum(U_k + V_k)\|^2 \leq 2\|\sum U_k\|^2 + 2\|\sum V_k\|^2$, we have

$$\left\| \sum_{k=1}^n (\Delta_k^2 - C_k^2) \right\|^2 = o(B_k^4) + 4 \sum_{1 \leq j < k \leq n} \int_0^1 V_k(\omega) V_j(\omega) d\omega.$$

In a similar way, we can prove

$$\left\| \sum_{k=1}^n \Delta_k \Delta_{k+1} \right\|^2 = o(B_k^4) + 4 \sum_{1 \leq j < k \leq n} \int_0^1 W_k(\omega) W_j(\omega) d\omega,$$

where $W_k = \sum_{q \in P(k+1)} a_q \sum_{r \in P(k)} a_r \cos 2\pi(n_q - n_r)\omega$. By Lemma 1, (1.8) and (2.7), we have

$$\left| \int_0^1 V_k(\omega) V_j(\omega) d\omega \right| \leq \sum_{q \in P(k)} |a_q| \sum_{h \in P(j)} |a_h| \max_{r \in P(k)} |a_r| \max_{i \in P(j)} |a_i| \frac{2^{j-k+1} \mu(k)}{c} \\ = o(B_k B_j \mu^{1/2}(k) \mu^{-1/2}(j) 2^{j-k} C_k C_j).$$

Because of (2.3), for large enough M , we have $\mu(k)/\mu(j) \leq M2^{k-j}$. Hence we have

$$\begin{aligned}
\sum_{1 \leq j < k \leq n} \int_0^1 V_k(\omega) V_j(\omega) d\omega &= o(B_n^2) \sum_{k=2}^n C_k \sum_{j=1}^{k-1} \sqrt{2}^{j-k} C_j \\
&= o(B_n^2) \sum_{k=2}^n C_k \left(\sum_{j=1}^{k-1} \sqrt{2}^{j-k} C_j^2 \right)^{1/2} \left(\sum_{j=1}^{k-1} \sqrt{2}^{j-k} \right)^{1/2} \\
&= o(B_n^2) \left(\sum_{k=2}^n C_k^2 \right)^{1/2} \left(\sum_{k=2}^n \sum_{j=1}^{k-1} \sqrt{2}^{j-k} C_j^2 \right)^{1/2} \\
&= o(B_n^3) \left(\sum_{j=1}^{n-1} C_j^2 \sum_{k=j+1}^n \sqrt{2}^{j-k} \right)^{1/2} \\
&= o(B_n^4).
\end{aligned}$$

Similarly we have $\sum_{1 \leq j < k \leq n} \int_0^1 W_k(\omega) W_j(\omega) d\omega = o(B_n^4)$. These estimates yield

$$\left\| \sum_{k=1}^n (\Delta_k^2 - C_k^2) \right\| = o(B_n^2) \quad \text{and} \quad \left\| \sum_{k=1}^n \Delta_k \Delta_{k+1} \right\| = o(B_n^2),$$

which imply (2.4). \square

Next we prove the Corollary. Let us put $\rho(i) = 1/(n_{i+1}/n_i - 1)$. Let $\Delta x(i)$ denote $x(i+1) - x(i)$.

We now assume $0 < \Lambda(i) \rightarrow \infty$, $\Delta \Lambda(i) = O(1)$, $\rho(i) = o(\Lambda(i))$ and $\rho(i) \leq \Lambda(i)$, and hereafter construct $\lambda(i)$ which satisfies $2\lambda \vee \rho(i) \leq \lambda(i)$, $\lambda(i) = o(\Lambda(i))$ and $\Delta \lambda(i) = o(1)$. The conditions of Theorem 1 are clearly derived from these. Put $\lambda = \frac{1}{2} \inf_i \Lambda(i)$. Let us first construct sequences $1 = i_0 < i_1 < i_2 < \dots$ and $\{\Lambda_0(i)\}$, $\{\Lambda_1(i)\}$, $\{\Lambda_2(i)\}$, \dots such that

$$(2.8) \quad \rho(i) = o(\Lambda_n(i)) \quad \text{and} \quad \Lambda_n(i) \rightarrow \infty \quad (i \rightarrow \infty, n \geq 0)$$

$$(2.9) \quad 2\lambda \vee \rho(i) \leq \Lambda_n(i) \leq \Lambda_{n-1}(i) \quad \text{and} \quad \Delta \Lambda_n(i) = \frac{1}{2} \Delta \Lambda_{n-1}(i) \quad (n \geq 1, i \geq i_n)$$

$$(2.10) \quad \Lambda_{n-1}(i) \leq \frac{2}{3} \Lambda_{n-2}(i) \quad (n \geq 2, i \geq i_n).$$

These sequences are constructed inductively in n . First we put $i_0 = 1$ and $\Lambda_0(i) = \Lambda(i)$. It is clear that (2.8) is satisfied for $n = 0$. After i_{n-1} and $\{\Lambda_{n-1}(i)\}$ have been constructed, we define i_n and $\{\Lambda_n(i)\}$ as follows: We can take $j > i_{n-1}$ such that

$$(2.11) \quad \rho(i) \leq \Lambda_{n-1}(i)/2 \quad \text{and} \quad \Lambda_{n-2}(i_{n-1}) \leq \Lambda_{n-2}(i)/3 \quad (i \geq j).$$

(The second condition must be omitted in case $n = 1$.) Let us take $i_n \geq j$ such that $\Lambda_{n-1}(i_n) = \min_{i \geq j} \Lambda_{n-1}(i)$ holds, and define $\Lambda_n(i)$ by

$$\Lambda_n(i) = \begin{cases} \Lambda_{n-1}(i) & i < i_n, \\ (\Lambda_{n-1}(i) + \Lambda_{n-1}(i_n))/2 & i \geq i_n. \end{cases}$$

By definition, (2.11) holds if we put $j = i_n$, and $\Lambda_{n-1}(i) \geq \Lambda_n(i_n)$ holds for all $i \geq i_n$. Therefore $\Lambda_n(i) \leq \Lambda_{n-1}(i)$ holds for $i \geq i_n$, and the rest of (2.9) is clear from definition and the first inequality of (2.11). (2.10) follows from the last inequality of (2.11). By definition, (2.8) is clear, and the sequences are well constructed.

If we put $\lambda(i) = \Lambda_n(i)$ for $i_n \leq i < i_{n+1}$, it satisfies $2\lambda \vee \rho(i) \leq \lambda(i)$, $\Delta\lambda(i) = \Delta\Lambda_n(i) = (\frac{1}{2})^n \Delta\Lambda(i) = o(1)$ and $\lambda(i) = \Lambda_n(i) \leq \Lambda_{n-1}(i) \leq (\frac{2}{3})^{n-1} \Lambda(i) = o(\Lambda(i))$. \square

Finally, we derive Theorem 1 from Corollary. By this we see that Theorem 1 and Corollary are equivalent. We now assume (1.5), (1.6), (1.7) or $\rho(i) = O(\lambda(i))$, and (1.8), and derive the conditions of Corollary. Conditions (1.5) and (1.8) imply $a_i/A_i \rightarrow 0$. We can therefore take an increasing sequence $\tilde{\lambda}(i)$ of positive numbers such that $\tilde{\lambda}(i)a_i/A_i \rightarrow 0$ and $\Delta\tilde{\lambda}(i) = o(1)$. If we put $\lambda_0(i) = \lambda(i) + \tilde{\lambda}(i)$, we have $0 < \lambda_0(i) \rightarrow \infty$, $\Delta\lambda_0(i) = o(1)$, $\rho(i) = O(\lambda_0(i))$ and $\lambda_0(i)a_i/A_i \rightarrow 0$. Next we construct $1 = i_0 < i_1 < i_2 < \dots$ and $\{\lambda_1(i)\}, \{\lambda_2(i)\}, \dots$ such that $\lambda_n(i) \geq \lambda_{n-1}(i)$,

$$0 < \lambda_n(i) \rightarrow \infty, \quad \Delta\lambda_n(i) = o(1), \quad \text{and} \quad \lambda_n(i)a_i/A_i \rightarrow 0 \quad (i \rightarrow \infty, n \geq 0)$$

$$|\Delta\lambda_n(i)| \leq 1, \quad \Delta\lambda_n(i) = 2\Delta\lambda_{n-1}(i), \quad \left| \frac{\lambda_n(i)a_i}{A_i} \right| \leq 1, \quad (i \geq i_n, n \geq 1)$$

$$\lambda_{n-1}(i) \geq \frac{3}{2}\lambda_{n-2}(i) \quad (i \geq i_n, n \geq 2).$$

It can be achieved first by taking i_{n+1} to satisfy $\lambda_n(i_{n+1}) = \min_{i \geq i_{n+1}} \lambda_n(i)$,

$$\left| \frac{\lambda_n(i)a_i}{A_i} \right| \leq \frac{1}{2}, \quad |\Delta\lambda_n(i)| \leq \frac{1}{2} \quad \text{and} \quad \lambda_{n-1}(i) \geq 2\lambda_{n-1}(i_n), \quad (i \geq i_{n+1}),$$

and then putting $\lambda_{n+1}(i) = 2\lambda_n(i) - \lambda_n(i_{n+1})$ if $i \geq i_{n+1}$ and $\lambda_{n+1}(i) = \lambda_n(i)$ otherwise. If we put $\Lambda(i) = \lambda_n(i)$ for $n_i \leq i < n_{i+1}$, we can verify the conditions on $\Lambda(i)$ in a similar way as before. \square

3. CONSTRUCTION OF COUNTEREXAMPLES

We may assume $\lambda(i) \rightarrow \infty$, since the condition $a_n = o(A_n)$ is necessary for (1.1). There is no loss of generality if we assume $c = 1$ and $\lambda(i) \geq 1$. Let us denote by $\|\cdot\|_\infty$ the sup-norm on $[0, 1]$. Redefine $\{p(k)\}$ by

$$p(0) = 0 \quad \text{and} \quad p(k) = \max \left\{ j \left| \sum_{i=1}^j \frac{1}{\lambda(i)} \leq k. \right. \right\} \quad (k \geq 1),$$

and define $l(k)$, $P(k)$, $\mu(k)$ and $\nu(k)$ as before by using new $\{p(k)\}$.

If $p(k) + 1 < p(k+1)$, we have

$$\frac{l(k)}{\nu(k)} + \frac{1}{\nu(k+1)} + \frac{1}{\nu(k-1)} \geq \sum_{i=p(k)}^{p(k+1)+1} \frac{1}{\lambda(i)} \geq 1 \geq \sum_{i=p(k)+1}^{p(k+1)} \frac{1}{\lambda(i)} \geq \frac{l(k)}{\mu(k)},$$

which implies $\liminf l(k)/\nu(k) \geq 1$ and $l(k) \leq \mu(k)$. By using $l(k) \leq \mu(k)$, in the same way as before, we can prove (2.2), $\mu(j) \sim \nu(j)$, and (2.3) in turn. Consequently

we have $\mu(j) \sim \nu(j) \sim l(j) \rightarrow \infty$, and therefore we can take $j_0 \geq 1$ such that $\mu(j)/2 \leq l(j) \leq 2\nu(j)$ for $j \geq j_0$. We note that

$$\sum_{j=j_0}^{\infty} \frac{1}{l(j)} = \sum_{j=j_0}^{\infty} \frac{l(j)}{l^2(j)} \geq \sum_{j=j_0}^{\infty} \sum_{i \in P(j)} \frac{1}{4\lambda^2(i)} = \infty.$$

Let us put

$$a_i = \begin{cases} 1 & \text{if } i \leq p(j_0), \\ A_{p(j)}/l(j) & \text{if } i \in P(j), j \geq j_0, \end{cases} \quad \text{and} \quad b_j = a_{p(j+1)},$$

and define Δ_k as before. We easily have $A_{p(j_0)} > 0$, $a_i = O(A_i/\lambda(i))$ and

$$(3.1) \quad A_{p(k+1)}^2 = A_{p(j_0)}^2 \prod_{j=j_0}^k \left(1 + \frac{1}{2l(j)}\right) \geq \frac{A_{p(j_0)}^2}{2} \sum_{j=j_0}^k \frac{1}{l(j)} \rightarrow \infty.$$

Let us take an increasing sequence $\{q(j)\}$ of integers such that

$$q(j_0) = p(j_0) + 1 \quad \text{and} \quad 2^{q(j+1)-q(j)} \geq \max\{2l(j+1), \pi A_{p(j)} l^2(j) j^2\},$$

and introduce $\{n_i\}$ by

$$n_{p(j)+l} = \begin{cases} 2^{p(j)+l} & \text{if } j < j_0, 1 \leq l \leq l(j), \\ 2^{q(j)} l & \text{if } j \geq j_0, 1 \leq l \leq l(j). \end{cases}$$

If $j \geq j_0$ and $1 \leq l \leq l(j)$, (1.7) is verified by noting $l(j) \leq 2\nu(j) \leq 2\lambda(p(j) + l)$;

$$\frac{n_{p(j)+l+1}}{n_{p(j)+l}} \geq 1 + \frac{1}{l(j)} \geq 1 + \frac{1}{\lambda(p(j) + l)} \quad \text{and} \quad \frac{n_{p(j)+1}}{n_{p(j)}} = \frac{2^{q(j)-q(j-1)}}{l(j)} \geq 2.$$

By using the Dirichlet kernel, we see that there exists an absolute constant $c_0 > 0$ such that for all integers m and l , $P(|\sum_{j=1}^l \cos 2\pi m j \omega| > l/e) \geq c_0/l$. Applying this, we have $P(|\Delta_j| \geq b_j l(j)/e) \geq c_0/l(j)$. Note that $b_j l(j)/e = A_{p(j)}/e$ for $j \geq j_0$. If we put $J_m = \{j = j_0, \dots, m \mid e A_{p(j)} \geq A_{p(m+1)}\}$, we have

$$\sum_{j=1}^m P\left(|\Delta_j| \geq \frac{A_{p(m+1)}}{e^2}\right) \geq \sum_{j \in J_m} P\left(|\Delta_j| \geq \frac{A_{p(j)}}{e}\right) \geq \sum_{j \in J_m} \frac{c_0}{l(j)}.$$

Since we have $A_{p(m+1)}/A_{p(j)} \leq \exp(\frac{1}{2} \sum_{k=j+1}^m 1/l(k))$, by putting $J'_m = \{j = j_0, \dots, m \mid \sum_{k=j+1}^m 1/l(k) \leq 2\}$, we have $J'_m \subset J_m$ and hence

$$(3.2) \quad \sum_{j=1}^m P\left(|\Delta_j| \geq \frac{A_{p(m+1)}}{e^2}\right) \geq \sum_{j \in J'_m} \frac{c_0}{l(j)} \rightarrow 2c_0.$$

Let $\omega = \sum_{r=1}^{\infty} 2^{-r} d_r(\omega)$ be the dyadic expansion of ω , and put

$$\tilde{X}_j(\omega) = \sum_{i \in P(j)} a_i \cos 2\pi n_i \left(\sum_{r=q(j)+1}^{q(j+1)} 2^{-r} d_r(\omega) \right) \quad \text{and} \quad X_j = \tilde{X}_j - E \tilde{X}_j.$$

Clearly, $\{X_j\}$ is an independent sequence. Because of $E\Delta_j = 0$ and the estimate

$$\|\Delta_j - \tilde{X}_j\|_\infty \leq b_j \sum_{i \in P(j)} 2\pi n_i 2^{-q(j+1)} \leq \pi b_j l^2(j) 2^{q(j)-q(j+1)} \leq 1/j^2,$$

we have $\|\Delta_j - X_j\|_\infty \leq 2/j^2$. If we put $\sigma_j^2 = EX_j^2$ and $s_m^2 = \sum_{j=1}^m \sigma_j^2$, we get

$$|s_m - A_{p(m+1)}| \leq \sum_{j=1}^m \|\Delta_j - X_j\|_\infty \leq 4 \quad \text{and} \quad |\sigma_j - E^{1/2} \Delta_j^2| \leq \frac{2}{j^2}.$$

Combining these with (3.1) and $E\Delta_j^2 = A_{p(j)}^2/2l(j)$, we have

$$(3.3) \quad s_m \rightarrow \infty \quad \text{and} \quad \sigma_m = O(1/m^2) + E^{1/2} \Delta_m^2 = o(A_{p(m)}) = o(s_m).$$

If the central limit theorem (1.1) holds, then it holds for $\{X_i\}$. The Lindeberg theorem claims that (Cf. Chapter XV of Feller [2]) under (3.3), the central limit theorem $s_m^{-1} \sum_{i=1}^m X_k \xrightarrow{D} N_{0,1}$ implies the Lindeberg condition $s_m^{-2} \sum_{i=1}^m E(X_i^2 : |X_i| > \varepsilon s_m) \rightarrow 0$ for all $\varepsilon > 0$. From this we have

$$\lim_{m \rightarrow \infty} \sum_{i=1}^m P(|X_i| > \varepsilon s_m) = 0 \quad \text{for all } \varepsilon > 0.$$

On the other hand, by (3.2) we have

$$0 < c_0 \leq \sum_{i=1}^m P(|\Delta_i| > A_{p(m+1)}/e^2) \leq \sum_{i=1}^m P(|X_i| > s_m/e^3)$$

for large m . These contradict each other. \square

4. FOURIER CHARACTER OF LACUNARY SERIES

We prove Theorem 3 for $\gamma = 2$. The general case can be proved by iterating the following argument γ times. Let us put $S_n(\omega) = \sum_{i=1}^n a_i \cos(2\pi n_i \omega + \phi_i)$. We may assume $\delta = \sup_i |a_i| < \infty$, otherwise the conclusion is clear. Let us put

$$r(0) = 0 \quad \text{and} \quad r(m) = \max\{i \mid A_i \leq \delta m, \} \quad (m \geq 1).$$

By $\delta^2 m^2 \geq A_{r(m)}^2 = A_{r(m)+1}^2 - a_{r(m)+1}^2/2 > \delta^2(m^2 - 1/2)$, we have $A_{r(m)} \sim \delta m$, $A_{r(m)}^2 - A_{r(m-1)}^2 \sim 2\delta^2 m$ and $A_{r(m)}^{-1} - A_{r(m-1)}^{-1} = O(m^{-2})$.

First we assume that $\{S_n\}$ converges on some set E with $|E| > 0$ and derive a contradiction. Let us put $b_i = a_i/A_i$, $B_m = \frac{1}{2} \sum_{i=1}^m b_i^2$ and $T_m(\omega) = \sum_{i=1}^m b_i \cos 2\pi n_i \omega$. Since each term of T_m is a product of the term of S_m and non-negative decreasing sequence $1/A_i$, by using the Abel's Theorem (Cf. (2.4) of Chapter I in Zygmund [11]), $\{T_n\}$ converges on E . Since we have

$$B_{r(m)}^2 = \sum_{k=1}^m \sum_{i=r(k-1)+1}^{r(k)} \frac{a_i^2}{2A_i^2} \sim \sum_{k=1}^m \frac{A_{r(k)}^2 - A_{r(k-1)}^2}{A_{r(k)}^2} \sim \sum_{k=1}^m \frac{2}{k} \sim \log m^2 \sim \log A_{r(m)}^2,$$

we get $B_k^2 \sim \log A_k^2$, and hence we can prove

$$(4.1) \quad b_i = o(B_i \sqrt{\log B_i / \lambda(i)}) \quad \text{and} \quad B_i \rightarrow \infty.$$

Let us repeat the above argument, i.e., take a sequence $\{r'(m)\}$ as $B_{r'(m)} \sim m$ and define $c_i = b_i/B_i$, $C_k = \frac{1}{2} \sum_{i=1}^k c_i^2$ and $Z_m(\omega) = \sum_{i=1}^m c_i \cos 2\pi n_i \omega$. In the same way we can prove that $C_k^2 \sim \log B_k^2$ and hence

$$(4.2) \quad c_i = o(C_i / \lambda(i)) \quad \text{and} \quad C_i \rightarrow \infty,$$

and that Z_∞ converges on E and thereby $Z_m/C_m \rightarrow 0$ on E . By (4.2) we can apply Theorem 1 and conclude that Z_m/C_m converges to $N_{0,1}$ on E , which is a contradiction.

Next we assume that the series is a Fourier series or a Fourier-Stieltjes series. Let us take $\rho \in (1/2, 1)$. By the Riesz-Kolmogorov inequality (Cf. (6.8) or (6.27) of Chapter VII in Zygmund [11]), we have $E|S_m|^\rho = O(1)$.

Let us redefine b_i as $b_i = a_i/A_{r(m+1)}$ if $r(m) < i \leq r(m+1)$, and define B_k and T_m as before by using these b_i . In a similar way, we can prove $B_k^2 \sim \log A_k^2$ and (4.1). For any m , let us take n such that $r(n) < m \leq r(n+1)$. By applying the Abel's partial summation (the Abel transformation) to $T_m = (S_m - S_{r(n)})/A_{r(n+1)} + \sum_{k=1}^n (S_{r(k)} - S_{r(k-1)})/A_{r(k)}$, we have

$$\begin{aligned} E|T_m|^\rho &= E \left| \frac{1}{A_{r(n+1)}} S_m + \sum_{k=1}^n \left(\frac{1}{A_{r(k)}} - \frac{1}{A_{r(k+1)}} \right) S_{r(k)} \right|^\rho \\ &\leq \frac{1}{A_{r(n+1)}^\rho} E|S_m|^\rho + \sum_{k=1}^n \left(\frac{1}{A_{r(k)}} - \frac{1}{A_{r(k+1)}} \right)^\rho E|S_{r(k)}|^\rho. \end{aligned}$$

Since $\left(\frac{1}{A_{r(k)}} - \frac{1}{A_{r(k+1)}} \right)^\rho = O(k^{-2\rho})$ is summable in k , we have $E|T_m|^\rho = O(1)$.

Let us redefine c_i as $c_i = b_i/B_{r'(m+1)}$ if $r'(m) < i \leq r'(m+1)$, and define C_k and Z_m as before. Then we have (4.2) and $E|Z_m|^\rho = O(1)$. Thus Z_m/C_m converges to $N_{0,1}$ in law, and to 0 in L^ρ -sense and hence in probability. This is a contradiction. \square

REFERENCES

1. P. Erdős, *On trigonometric series with gaps*, Magyar Tud. Akad. Mat. Kutató Int. Közl. **7** (1962), 37–42.
2. W. Feller, *An introduction to probability theory and its applications* vol. II, second ed., Wiley, New York, 1971.
3. A. N. Kolmogorov, *Une contribution à l'étude de la convergence des séries de Fourier*, Fund. Math. **5** (1924), 26–27.
4. R. Salem & A. Zygmund, *On lacunary trigonometric series*, Proc. Nat. Acad. Sci. **33** (1947), 333–338.
5. S. Takahashi, *A version of the central limit theorem for trigonometric series*, Tôhoku Math. J. **16** (1964), 384–398.
6. ———, *On trigonometric series with gaps*, Tôhoku Math. J. **17** (1965), 227–234.
7. ———, *On lacunary trigonometric series*, Proc. Japan Acad. **41** (1965), 503–506.
8. ———, *On the lacunary Fourier series*, Tôhoku Math. J. **19** (1967), 79–85.
9. ———, *On lacunary trigonometric series II*, Proc. Japan Acad. **44** (1968), 766–770.
10. A. Zygmund, *On the convergence of lacunary trigonometric series*, Fund. Math. **16** (1930), 90–97; *Correction*, Fund. Math. **18** (1932), 312.

11. ———, *Trigonometric series* vol. I, second ed., Cambridge Univ. Press, Cambridge, (1959).

DEPARTMENT OF MATHEMATICS, KOBE UNIVERSITY, ROKKO, KOBE, 657 JAPAN
E-mail address: `fukuyama@math.s.kobe-u.ac.jp`

YAMAMICHI-CHO 21-1-601, HIROSAKI, 036 JAPAN