



Diabetic chorea as a neurological complication

Kenzaka, Tsuneaki
Fujikawa, Moemi
Matsumoto, Masanori

(Citation)

Clinical Case Reports, 6(7):1398-1399

(Issue Date)

2018-07

(Resource Type)

journal article

(Version)

Version of Record

(Rights)

© 2018 The Authors. Clinical Case Reports published by John Wiley & Sons Ltd.
This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

(URL)

<https://hdl.handle.net/20.500.14094/90005513>



Diabetic chorea as a neurological complication

Tsuneaki Kenzaka^{1,2}  | Moemi Fujikawa² | Masanori Matsumoto²

¹Division of Community Medicine and Career Development, Kobe University Graduate School of Medicine, Kobe, Japan

²Department of Internal Medicine, Hyogo Prefectural Kaibara Hospital, Tanba, Japan

Correspondence

Tsuneaki Kenzaka, Division of Community Medicine and Career Development, Kobe University Graduate School of Medicine, Kobe, Hyogo, Japan.

Email: smile.kenzaka@jichi.ac.jp

Key Clinical Message

Diabetic chorea accompanies hyperglycemic states or drastic changes in blood glucose levels and involves the acute onset of unilateral or bilateral choreatic movements. Diabetes is sometimes discovered due to diabetic chorea, and thus, diabetic chorea is considered an important neurological complication.

KEYWORDS

diabetes mellitus, diabetic chorea, neurological complication

The patient was a 57-year-old man being treated for bipolar disorder. He had been diagnosed with type 2 diabetes, but was not receiving drug therapy. A month before

the examination, his depressive symptoms worsened because of stress at work. He was spending more time at home, mainly consuming soft drinks, sweet breads, and other foods that were high in sugar. One week before the admission to our hospital, his right arm began moving involuntarily. The symptoms worsened, so he came for outpatient examination. Blood test results were as follows: blood glucose, 573 mg/dL; HbA1c, 16.0%; calculated serum osmolality, 315 mOsm/kg; pH, 7.359; pCO₂, 41.9 mm Hg; and bicarbonate ions, 23.1 mEq/L. He did not present with liver dysfunction or electrolyte imbalance or thyroid dysfunction, and was negative for urinary ketones. Head CT showed isodense regions in the left putamen (Figure 1). Video S1 shows the choreatic movements observed during inpatient observation. Glucose control improved with insulin therapy, and after about 1 week, the choreatic movements also improved. Based on the acute onset of unilateral choreatic movements that improved along with glucose control, the patient was diagnosed with diabetic chorea, which is also recognized as chorea, hyperglycemia, basal ganglia syndrome (C-H-BG).



FIGURE 1 Isodensity to mild hyperdensity in the left putamen in head CT scan

CONFLICT OF INTEREST

None of the authors have any financial interests to disclose, nor do they have any conflict of interests to declare.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

AUTHORS' CONTRIBUTIONS

TK: managed redaction, correction, and reconstruction of the manuscript. MF: managed the case and redaction and correction of the manuscript. MK: assisted with clinical management of the case and correction of the manuscript. All authors read and approved the final manuscript.

ORCID

Tsuneaki Kenzaka  <http://orcid.org/0000-0002-3120-6605>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Kenzaka T, Fujikawa M, Matsumoto M. Diabetic chorea as a neurological complication. *Clin Case Rep*. 2018;6:1398–1399. <https://doi.org/10.1002/ccr3.1579>