



Ectopic expression of band 3 anion transport protein in colorectal cancer revealed in an autoimmune hemolytic anemia patient

Kawamoto, Shinichiro ; Kamesaki, Toyomi ; Masutani, Ryota ; Kitao, Akihito ; Hatanaka, Kazuo ; Imakita, Masami ; Tamaki, Toshiharu ;...

(Citation)

Human Pathology, 83:193-198

(Issue Date)

2019-01

(Resource Type)

journal article

(Version)

Accepted Manuscript

(Rights)

© 2018 Elsevier.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

(URL)

<https://hdl.handle.net/20.500.14094/90005661>



Title: Ectopic Expression of Band 3 Anion Transport Protein in Colorectal Cancer Revealed in an Autoimmune Hemolytic Anemia Patient.

Authors: Shinichiro Kawamoto, MD, PhD¹, Toyomi Kamesaki, MD, PhD², Ryota Masutani, MT³, Akihito Kitao, MD⁴, Kazuo Hatanaka, MD⁵, Masami Imakita, MD, PhD⁶, Toshiharu Tamaki, MD, PhD⁷, Takayuki Takubo, MD, PhD⁸

Addresses of the institutions

¹ Department of Transfusion Medicine and Cell Therapy, Kobe University Hospital, Hyogo, Japan

² Center for Community Medicine, Jichi Medical University, Tochigi, Japan

³ Department of Central Clinical Laboratory, Osaka Medical College Hospital, Osaka, Japan

⁴ Department of Medical Oncology/Hematology, Kobe University Hospital, Hyogo, Japan

⁵ Department of Hematology, Osaka Red Cross Hospital, Osaka, Japan

⁶ Pathology division of Rinku General Medical Center, Osaka, Japan

⁷ Department of Internal Medicine, Nogami Hospital, Osaka, Japan

⁸ Japanese Red Cross Society, Osaka, Japan

Corresponding author: Shinichiro Kawamoto M.D., PhD.

Address: Chuoh-ku Kusunokicho 7-5-2, Kobe City, 650-0017, Hhyogo, Japan

- 1 **E-mail address:** skawamo@med.kobe-u.ac.jp
- 2 **TEL:** +81-078-382-5111
- 3 **FAX:** +81-078-382-5821
- 4 **Declarations of interest:** none
- 5 **Number of tables/figures:** 1/2

Summary

Cancer patients occasionally have anemia with high mean corpuscular volume in addition to iron deficiency anemia. Secondary autoimmune hemolytic anemia (AIHA) following cancer is also observed with low frequency. To date, no causal mechanisms for these disease states have been reported.

Here, we present the case of an 80-year-old woman with AIHA that was resistant to prednisolone.

Further examinations revealed primary adenocarcinoma of the sigmoid colon and primary squamous cell carcinoma in the right lung. After resections of these tumors, her anemia partially improved until a colon cancer-derived metastatic tumor was detected in the left lung.

Immunoprecipitation of erythrocyte membrane proteins with an autoantibody followed by mass spectrometry/western blotting identified band 3 as the target of the autoantibody.

Immunohistochemical analysis revealed ectopic expression of band 3 in the colon adenocarcinoma. To our knowledge, this is the first report that identifies the cause in a case of anemia without bleeding in a cancer patient and that defines a mechanism underlying secondary AIHA following cancer progression.

Key words

AIHA, Autoantibody, Band 3, Colon cancer, Mass spectrometry

1. Introduction

Secondary autoimmune hemolytic anemia (AIHA) is attributed to primary diseases, including lymphoid neoplasms, collagen diseases, and ovarian cysts. Typically, treatment of primary diseases leads to AIHA remission. Cases of AIHA secondary to cancer are rare [1].

Band 3 protein, also known as anion exchanger 1, is an anion transporter that exchanges HCO_3^- and Cl^- in equal proportions [2]. It is abundantly expressed in red blood cells (RBCs) and renal tubule cells. Band 3 is one target of autoantibodies associated with idiopathic AIHA. Other targets include Rhesus group D antigen (RhD) and glycophorin A [3]. Aberrant expression of band 3 has been reported in gastrointestinal (GI) cancers, including esophageal, gastric, and colon cancers, and band 3 expression is a factor in the pathogenesis and poor prognosis of patients with GI cancers [4, 5]. The relationship between AIHA onset and aberrant expression of band 3 in cancer cells remains unclear.

In this report, we identify the possible cause of an anemia with previously unknown etiology in a cancer patient, by analyzing secondary AIHA following colon cancer using mass spectrometry and immunological techniques. This report provides critical information in the fields of oncology, hematology, and immunology.

2. Materials and methods

2.1. Isolation of Ghosts and IgG autoantibodies

Whole blood was collected into EDTA-containing tubes, and RBCs were isolated using Ficoll-Paque PLUS in accordance with the manufacturer's instructions (GE, Cat. No. 17144002, NJ, USA). RBCs were processed for isolation of ghost membranes in accordance with Baker's method [6]. The IgG autoantibodies were isolated from serum using an IgG purification kit (Dojindo Molecular Technologies, Inc. Cat. No. AP02, Rockville, USA). Immunoprecipitation was performed using the isolated autoantibodies with Protein G Sepharose (Calbiochem, Cat. No. IP05, USA).

2.2. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis and immunoblotting

Gradient polyacrylamide gels (10%–20%) and polyvinylidene difluoride membranes were purchased from ATTO Inc. (Figure 1; Cat. ET-1020L, Cat. AE-6668, Tokyo, Japan). Any kD™ Mini-PROTEAN TGX Precast Gels and a Trans-Blot Turbo Transfer Pack were purchased from BIO-RAD Laboratories Inc. (Figure 2; Cat. #456-9033, Cat #170-4156, USA). Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and immunoblotting were performed by conventional methods using the following buffers: lysis buffer (50 mM Tris-HCl pH 8.0, 0.1% Tween 20, 0.1% SDS, 150 mM NaCl, and 0.5 mM EDTA), 2× sample buffer (0.25 M Tris-HCl, pH 6.8, 20% 2-mercaptoethanol, 4% SDS, 20% sucrose, and 0.002% bromophenol blue), running buffer (25 mM Tris, 0.192 M glycine, and 0.1% SDS), and blocking buffer; 3% skim milk in

Tween-PBS (137 mM NaCl, 2.7 mM KCl, 8.1 mM Na₂HPO₄·12H₂O, 1.5 mM KH₂PO₄, and 0.1% Tween 20). After incubation with the indicated antibodies, membranes were incubated with secondary antibodies. Detection was performed by enhanced chemiluminescence (ECL). Antibodies and reagents for western blotting were commercially purchased as follows: molecular marker, Amersham (Cat. No. RNP800, NJ, USA) and BIO-RAD (Cat. No. #1610374, USA); peroxidase-labeled secondary anti-human IgG (H+L) antibody, Cytodiagnostics (Cat. ABH-01-11, ON, Canada); rabbit polyclonal anti-slc4a1 antibody, MBL (Cat. BMP012, Aichi, Japan); horse radish peroxidase (HRP)-conjugated anti-rabbit-IgG secondary antibody, Santa Cruz Biotech (Cat. SC-2004, USA); anti-human RhD antibody, Sigma-Aldrich (Cat. No. sab2107753, MO, USA); anti-band3 antibody, MBL (Cat. BMP012, Nagoya, Aichi, Japan); and ECL substrate, Merck Millipore (Cat. No. WBLUR0100, MA, USA).

2.3. Mass spectrometry

After staining SDS-PAGE gels with Coomassie G-250 (Invitrogen, Cat. No. LC6060, Carlsbad, CA, USA), visualized protein bands were extracted from the gel. After purification of the extracted proteins, samples were analyzed by mass spectrometry (liquid chromatography ion-trap time-of-flight mass spectrometry, Shimadzu Co.).

3. A case report

An 80-year-old woman with a history of mycoplasma pneumonia, but with no history of transfusions presented with macrocytic anemia. Examination of the upper digestive tract was not performed, and no occult blood was found in her stool sample. An abdominal ultrasonogram revealed no abnormalities, but a computed tomography (CT) scan revealed a small nodule in her lower right lung.

Because her anemia worsened, she was referred to our hospital. Based on the results of her laboratory tests, including Coombs test and bone marrow aspiration, she was diagnosed with AIHA (Table). As anemia worsened (hemoglobin; Hb 79 g/L, hematocrit; Ht 24.6 L/L, mean corpuscular volume; MCV 116 fL, reticulocyte; Ret $0.17 \times 10^{12}/L$), prednisolone was initiated at a dose of 0.5 mg/kg. This treatment achieved only a partial response. We therefore suspected that a malignancy might be causing her secondary AIHA. The results of repeated tests for occult blood in stool samples were negative. A chest CT was performed to follow-up on the lesion that was previously detected in her lower right lung. Tumor growth was observed in this follow-up CT scan (Fig. 1A). Positron emission tomography (PET) revealed no metastasis of the tumor to the lymph nodes (Fig. 1B); however, fluorodeoxyglucose uptake was also observed in the sigmoid colon (Fig. 1C). Through colonoscopy, advanced cancer was confirmed (Fig. 1D), and a diagnosis of adenocarcinoma was made based on histopathological examination of the colon lesion. The

lung tumor was resected, and histopathologically determined to be squamous cell carcinoma. After subsequent colon cancer surgery, the patient's anemia improved slightly (Hb 136 g/L, Ht 42.2 L/L, MCV 99 fL, Ret $0.10 \times 10^{12}/L$) with low-dose prednisolone (5 mg/day). Over time, the number of reticulocytes gradually increased again, and anemia worsened 2 months post surgery (Hb 103 g/L, Ht 32.2 L/L, MCV 105 fL, Ret $0.14 \times 10^{12}/L$). This indicated AIHA recurrence and lung or colon cancer relapse. The patient's chest CT scan revealed a small nodule in her left lung (Fig. 1E). This tumor was resected, and histopathologically identified as adenocarcinoma, indicating colon cancer metastasis (Fig. 1 F). We therefore suspected that the adenocarcinoma might have caused the AIHA. After the left lung surgery, the patient refused treatment with sufficient prednisolone to treat her AIHA and adjuvant chemotherapy for colon cancer due to fear of their side effects. Her AIHA was thus managed with 5 mg/day prednisolone, leaving her with moderate anemia (Hb 105 g/L, Ht 33.6 L/L, MCV 107 fL, Ret $0.13 \times 10^{12}/L$). The last follow up with this patient was three years post surgery, although detailed information was not available.

4. Results

After obtaining informed consent from the patient, experiments were performed to characterize the autoantibody and its target. Results of the Coombs test and flow cytometry revealed that the class of immunoglobulin that bound the patient's erythrocytes was IgG (data not shown). IgG was collected from the patient's plasma sample using protein G sepharose and used for

immunoprecipitation and western blotting of RBC membrane proteins. Proteins of three molecular weights (*MW*), in the range of 24,000–30,000 daltons, were detected in immunoprecipitates using the autoantibody as the primary antibody and an HRP-conjugated anti-human IgG secondary antibody (Fig. 1G). These immunoprecipitates were not detected with the secondary antibody alone, indicating that these bands were not immunoglobulins. Because the molecular weight of RhD is 31,000 daltons, we suspected that one of the largest proteins detected was RhD. As expected, probing immunoprecipitates with an anti-RhD antibody as primary antibody detected similar protein bands (Fig. 1H). We hypothesized that the colon adenocarcinoma expressed a molecule that had homology with RhD antigen.

Histopathological examinations were performed using the same anti-RhD antibody. The lung cancer and normal colon tissue were not stained with the antibody (data not shown). Colon adenocarcinoma was only modestly stained. We concluded that this staining was non-specific because this tissue stained weakly with anti-rabbit secondary antibody alone (data not shown). This suggests that the target of the patient's autoantibody was not the protein with cross-reacting epitope to RhD antigen.

To further characterize the autoantibody, three bands that were immunoprecipitated from RBC membranes with the autoantibody were extracted from the gel, fragmented, and subjected to mass spectrometry. Band 3 was detected in the immunoprecipitates, although the intact molecular

weight of band 3 protein is approximately 100,000 daltons. The largest band, approximately 31,000 daltons, contained peptides that partially matched band 3 amino acid residues 139–263 and 541–878, and covered almost the entire sequence (Fig. 2A). Immunoblotting of the immunoprecipitates with a rabbit anti-band 3 polyclonal antibody, focusing on low molecular weight proteins, detected three similar bands, although one of these was a new band at 17,000 daltons rather than the band previously detected at 24,000 daltons (Fig. 2B).

Proteins expressed in the patient's surgically resected colon and lung samples were analyzed immunohistochemically using the same anti-band 3 antibody used to probe immunoprecipitations. The colon adenocarcinoma stained strongly with the anti-band 3 antibody (Fig. 2C), whereas normal colon tissue and lung squamous carcinoma did not (Fig. 2D, E). The metastatic lesion of colon adenocarcinoma found in the left lung also stained with the antibody (Fig. 2F).

5. Discussion

In the present report, we confirmed that band 3 protein was ectopically expressed in adenocarcinoma cells of the sigmoid colon and it was targeted by autoantibodies in a patient with secondary AIHA, whose fecal immunological test (FIT) was negative.

The Coombs test is not commonly performed for patients with GI cancers, because anemia is most frequently due to bleeding and improves after tumor resection. However, cancers that are associated with anemia often show high mean corpuscular volumes (MCVs), instead of iron

1 deficiency anemia patterns. Few previous investigators have suspected that production of
2 autoantibodies to erythrocytes may be another cause of anemia in cancer patients [1, 7, 8]. To date,
3 no precise analysis of the cause of antibody production, or identification of autoantibody targets
4 has been performed.

5 It is possible that natural secretion of anti-band 3 autoantibody into the serum and binding to
6 erythrocyte membranes are enhanced in many patients with colorectal cancer due to phagocytosis
7 of cancer cells, and antigen presentation by immune cells such as $\gamma\delta$ T cells in intraepithelial
8 lymphocytes [9]. Disturbing the balance of the $\gamma\delta$ T cell repertoire has been shown to induce
9 autoantibody binding to hematopoietic cells [10]. In healthy individuals, naturally occurring anti-
10 band 3 antibodies exist that remove senescent RBCs from the reticuloendothelial system [11]. In
11 this way, immunological responses can cause high MCV anemia that reflects the increase in
12 reticulocytes without bleeding in cancer patients.

13 Two issues arose out of our protein analysis results. The first relates to the reaction between the
14 anti-RhD antibody and the protein that is precipitated by the autoantibody (Fig. 1H). Although
15 the reason for this reaction is unclear, a previous report supports this phenomenon [12]. Gordon
16 et al. reported that an anti-C IgG autoantibody to the Rhesus blood group system appeared in a
17 patient with adenocarcinoma [13]. Band 3 may have been expressed in this patient's
18 adenocarcinoma, and the anti-C antibody also reacted with band 3. In support of this explanation,

1 we found in our current study that the anti-RhD antibody moderately reacted with colon
2 adenocarcinoma (data not shown). The lack of RhD peptide detection in our mass spectrometry
3 analysis of immunoprecipitated bands suggests the absence of anti-RhD autoantibodies in the
4 serum of the present patient.

5 The second issue is that the band 3 fragments detected had molecular weights of 17,000–30,000
6 daltons, while the putative molecular weight of full-length band 3 is 100,000 daltons (Fig. 1G,
7 H and Fig. 2B). Baker's method for preparing RBC protein membranes, erythrocyte 'ghosts,'
8 does not cleave Band 3 [6]. We also confirmed that our reducing conditions, using 2-

9 mercaptoethanol, did not cleave band 3 (data not shown). This result agrees with that of a
10 previous report, which analyzed senescent cell antigens related to band 3 [14]. Proteolytic
11 fragments of band 3 are recognized by physiologic IgG autoantibodies [15]. As residual
12 leukocyte debris after preparation of erythrocyte 'ghosts' has proteolytic activity, it is possible
13 that this debris caused the fragmentation of band 3 during storage [16]. In addition, caspase-3, a
14 member of the cysteine-aspartic acid protease family, might be responsible for our observation,
15 because parts of the N-terminal cytoplasmic and transmembrane domains of band 3 are known
16 to be cleaved by caspase-3 [17]. Caspase-3 fragments of band 3 possess autoantibody
17 recognition sites. The amino acid residues 160–878 of band 3 that were detected by mass
18 spectrometry in the current study also possess antigenicity [18,19]. As caspase-3 is expressed in

colon cancer and is considered to be a poor prognostic factor for patients, autoantibody binding to band 3 could increase owing to cancer-related stress [20].

Autoantibody binding to erythrocyte membranes of patients with AIHA is measurable, even in those with Coombs-negative AIHA [21]. The amount of Red-blood-cell-bound IgG in the present case was higher than normal reference levels (Table) and was detectable by flow cytometry (data not shown). Therefore, the cancer-related anti-band 3 autoantibody can potentially be quantified by flow cytometry and might be useful for screening of colorectal cancers that substitute FIT [22]. Considering the follow up duration of the current case after surgery, screening by flow cytometry might improve survival of colorectal cancer patients even in advanced stages.

In conclusion, the autoimmunity induced by ectopic expression of band 3 in patients with colon adenocarcinoma correlates with development of secondary AIHA, and causes anemia without bleeding, possibly due to phagocytosis of autoantibody-bound RBCs by macrophages.

Acknowledgments

We thank Dr. Naoya Hatano (Kobe University) and Emi Yanagida (Hokkaido University) for performing the mass spectrometry and immunohistochemical analyses, respectively, and Toshiyuki Ikemoto and Hiroko Tanada (Osaka Medical College) for performing flow cytometry

1 analyses. We also appreciate the assistance of Dr Akinori Nishikawa (Wakayama Medical
2 University) and Dr. Hiroyoshi Ichihara (Fuchu Hospital) for obtaining further information.

3

4 **Disclosure Statement**

5 The authors declare that they have no conflicts of interest.


References

1. Puthenparambil, J., Lechner, K., and Kornek, G. Autoimmune hemolytic anemia as a paraneoplastic phenomenon in solid tumors: A critical analysis of 52 cases reported in the literature. *Wien Klin Wochenschr.* 2010; 122: 229-236.
2. Reithmeier, R.A., Casey, J.R., Kalli, A.C., Sansom, M.S., Alguel, Y., and Iwata S. Band 3, the human red cell chloride/bicarbonate anion exchanger (AE1, SLC4A1), in a structural context. *Biochim Biophys Acta.* 2016; 1858: 1507-1532.
3. Baker, R.N., Casswell, K.M., Reid, M.E., Sokol, R.J., and Elson, C.J. Identification of autoantigens in autoimmune haemolytic anaemia by a non-radioisotope immunoprecipitation method. *Br J Haematol.* 1992; 82: 126-132.
4. Shen, W.W., Wu, J., Cai, L., Liu, B.Y. *et al.* Expression of Anion Exchanger 1 Sequesters p16 in the Cytoplasm in Gastric and Colonic Adenocarcinoma. *Neoplasia.* 2007; 9: 812-9.
5. Shiozaki, A., Kudou, M., Ichikawa, D. *et al.* Expression and role of anion exchanger 1 in esophageal squamous cell carcinoma. *Oncotarget.* ; 8: 17921-17935.
6. Baker, R.N. Electrophoretic Analysis of Erythrocyte Membrane Proteins and Glycoproteins from Different Species. *Comp Hematol Intern.* 1991; 1: 155-160
7. Green, H.N., Wakefield, J., and Littlewood, G. The nature of cancer anemia and its bearing on the immunological theory of cancer. *British Med J.* 1957; 2: 779-784.

8. Sokol, R.J., Brooker, D.J., and Stamp, R. Erythrocyte autoantibodies, autoimmune haemolysis, and carcinoma. *J Clin. Pathol.* 1994; 47: 340-343.
9. Lafont, V., Sanchez, F., Laprevotte, E. *et al.* Plasticity of $\gamma\delta$ T Cells: Impact on the Anti-Tumor Response. *Front Immunol.* 2014; 5: 622.
10. Huang, Y., Heiser, R.A., Detanico, T.O. *et al.* $\gamma\delta$ T cells affect IL-4 production and B-cell tolerance. *Proc Natl Acad Sci U S A.* 2015; 112: E39-48.
11. Pantaleo, A., Giribaldi, G., Mannu, F., Arese, P., Turrini, F. Naturally occurring anti-band 3 antibodies and red blood cell removal under physiological and pathological conditions. *Autoimmun Rev.* 2008; 7: 457-462.
12. Victoria, E.J., Mahan, L.C., and Masouredis, S.P. Anti-Rh0(D) IgG binds to band 3 glycoprotein of the human erythrocytemembrane. *Proc Natl Acad Sci USA.* 1981; 78: 2898-2902.
13. Gordon, P.A., Baylis, P.H., and Bird, G.W.G. Tumour-induced autoimmune haemolytic anemia. *Br Med J.* 1976; 6025: 1569-1570.
14. Kay, M.M., Goodman, S.R., Sorensen, K. *et al.* Senescent cell antigen is immunologically related to band 3. *Proc Natl Acad Sci USA.* 1983; 80: 1631-1635.
15. Kay, M.M.B. Localization of senescent cell antigen on band 3. *Proc Natl Acad Sci USA.* 1984; 81: 5753-5757.

- 1 16. Fairbanks, G., Steck, TL., Wallach, DF. Electrophoretic analysis of the major polypeptides of
2 the human erythrocyte membrane. *Biochemistry*. 1971; 10: 2606-2617.
- 3 17. Mandal, D., Baudin-Creuz, V., Bhattacharyya, A. *et al*. Caspase-3-mediated Proteolysis of
4 the N-terminal Cytoplasmic Domain of the Human Erythroid Anion Exchanger 1 (Band 3). *J*
5 *Biol Chem*. 2003; 278: 52551-52558.
- 6 18. Kay, M. Immunoregulation of cellular life span. *Ann N Y Acad Sci*. 2005; 1057: 85-111.
- 7 19. Moriyama, R., Kawamatsu, S., Kondo, Y., Tomida, M., and Makino, S. Antigenic
8 determinant of the Cytoplasmic domain of Band 3 from Bovine Erythrocyte Membrane.
9 *Arch Biochem Biophys*. 1989; 274: 130-137.
- 10 20. Jonges, L.E., Nagelkerke, J.F., Ensink, N.G. *et al*. Caspase-3 activity as a prognostic factor in
11 colorectal carcinoma. *Lab Invest*. 2001; 81: 681-688.
- 12 21. Kamesaki, T., Oyamada, T., Omine, M., Ozawa, K., and Kajii, E. Cut-off value of red-
13 blood-cell-bound IgG for the diagnosis of Coombs-negative autoimmune hemolytic anemia.
14 *Am J Hematol*. 2009; 84: 98-101.
- 15 22. Inadomi JM. Screening for Colorectal Neoplasia. *N Engl J Med*. 2017; 376: 149-156.

1



2

3 Figure 1. Autoantibody induced in a patient with colon and lung double cancer.

4 A, B) A squamous cell carcinoma detected in the right lung by Chest CT and PET CT examination.

5 C) A colon adenocarcinoma detected by PET CT examination. D) An advanced sigmoid colon

6 adenocarcinoma without bleeding revealed by colonoscopy. E) A left lung metastatic lesion of

7 recurrent colon adenocarcinoma. F) Hematoxylin-Eosin staining of left lung metastatic lesion of

8 colon adenocarcinoma - original magnification x40.



9 G) Western blotting of RBC ghost immunoprecipitates by autoantibody. Lane 1: molecular

10 marker. Lane 2: Membrane was incubated with autoantibodies followed by HRP-conjugated anti-

11 human IgG (H+L) secondary antibody. Lane 3: Membrane was incubated with HRP-conjugated

12 anti-human IgG secondary antibody without autoantibody. Arrow indicates IgG heavy chain. H)

1 Detection of similar size bands by anti-RHD antibody. Immunoprecipitate of RBC ghost by
2 autoantibody was subjected to western blotting using anti-human RhD polyclonal antibody and
3 HRP-conjugated anti-rabbit secondary antibody.
4 Abbreviations: M molecular marker; IP immunoprecipitate; Ab antibody; H immunoglobulin
5 heavy chain.
6 .



7
8 Figure 2. Band 3 protein detected in immunoprecipitates and colon adenocarcinoma.
9 A) Amino acid sequences detected by mass spectrometry from a 31,000-dalton-
10 immunoprecipitated band. Red letters indicate detected peptides found in full length band 3 amino

- 1 acid sequence. B) Immunoprecipitates detected by recombinant anti-band 3 antibody. C-F)
- 2 Histopathological staining with anti-band 3 antibody. C) Colon adenocarcinoma - original
- 3 magnification x400. D) Normal colon tissue - original magnification x100. E) Right lung
- 4 squamous cell carcinoma - original magnification x100. F) Left lung metastatic lesion of colon
- 5 adenocarcinoma - original magnification x40.