



Unconjugated interferon-stimulated gene 15 specifically interacts with the hepatitis C virus NS5A protein via domain I

Minami, Nanae ; Abe, Takayuki ; Deng, Lin ; Matsui, Chieko ; Fukuhara, Takasuke ; Matsuura, Yoshiharu ; Shoji, Ikuo

(Citation)

Microbiology and Immunology, 61(7):287-292

(Issue Date)

2017-07

(Resource Type)

journal article

(Version)

Accepted Manuscript

(Rights)

© 2017 The Societies and John Wiley & Sons Australia, Ltd. This is the peer reviewed version of the following article: [Microbiology and Immunology, 61(7):287-292, 2017], which has been published in final form at <https://doi.org/10.1111/1348-0421.12493>. This article may be used for non-commercial purposes in accordance with Wiley Terms...

(URL)

<https://hdl.handle.net/20.500.14094/90005839>



Note

Unconjugated interferon-stimulated gene 15 specifically interacts with the hepatitis C virus NS5A protein via domain I

Short running title: Unconjugated ISG15-HCV NS5A interaction

Nanae Minami,¹ Takayuki Abe,¹ Lin Deng,¹ Chieko Matsui,¹ Takasuke Fukuhara,²
Yoshiharu Matsuura,² and Ikuo Shoji^{1,*}

¹Division of Infectious Disease Control, Center for Infectious Diseases, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe, Hyogo, 650-0017, Japan

²Department of Molecular Virology, Research Institute for Microbial Diseases, Osaka University, 3-1 Yamada-oka, Suita, Osaka 565-0871, Japan

***To whom correspondence should be addressed:** Ikuo Shoji, M.D., Ph.D.

Division of Infectious Disease Control, Center for Infectious Diseases, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe, Hyogo 650-0017, Japan.

Tel: +81-78-382-5500. **FAX:** +81-78-382-5519.

E-mail: ishoji@med.kobe-u.ac.jp

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, Typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/1348-0421.12493.

© 2017 The Societies and John Wiley & Sons Australia, Ltd

ABSTRACT

Interferon-stimulated gene 15 (ISG15) is a ubiquitin-like protein induced by type I interferon. Although several groups have reported ISGylation of the HCV NS5A protein, it is still unclear whether ISGylation of NS5A has anti-viral or pro-viral effects in HCV infection. In the present study, we examined the role of ISGylation-independent, unconjugated ISG15 in HCV infection. Immunoprecipitation analyses revealed that ISG15 specifically interacted with NS5A domain I. The ISG15 mutants lacking a C-terminal glycine residue essential for ISGylation still interacted with NS5A protein. Taken together, these results suggest that unconjugated ISG15 may affect the functions of HCV NS5A through protein-protein interaction.

Keywords: HCV, ISG15, NS5A

List of Abbreviations: Apolipoprotein A1 (ApoA1), Chikungunya virus (CHIKV), FK506-binding protein (FKBP8), Heat shock protein 27 (HSP27), Hepatitis C virus (HCV), Hepatocyte nuclear factor 1 α (HNF-1 α), Interferon (IFN), Interferon-stimulated gene 15 (ISG15), 2'-5'oligoadenylate synthase (2'-5'OAS), ovarian tumor protein 7B (OTUD7B), Proximity ligation assay (PLA), Virus-like particle (VLP).

Revised manuscript, changes are written using red text.

Subject section: Virology

Specific Fields: Animal RNA virus

Word count

Abstract: 100 words

Main Text: 1488 words

Figures: 3

MAIN TEXT

Hepatitis C virus (HCV) is a major cause of chronic liver disease affecting 130-170 million people worldwide (1). Chronic HCV infection causes steatosis, fibrosis, cirrhosis and hepatocellular carcinoma (HCC) (2). HCV belongs to the Hepacivirus genus of the Flaviviridae family. The positive sense single-strand RNA genome encodes a polyprotein that is cleaved by cellular signalases and viral proteases into 10 viral proteins: three structural proteins (core, E1 and E2), the p7 polypeptide, and six non-structural proteins (NS2, NS3, NS4A, NS4B, NS5A, and NS5B) (3). Unlike other HCV non-structural proteins such as viral protease or polymerase, HCV NS5A lacks any enzymatic motif. Instead, it has been well established that NS5A is an essential viral component for efficient viral replication through the recruitment and interaction with various cellular host factors (4).

The interferon-stimulated gene 15 (ISG15) is a type I interferon (IFN)-inducible gene and a ubiquitin-like protein containing two ubiquitin-like domains (5). Because the modification of protein by ISG15 is similar to that by ubiquitylation, protein conjugation by ISG15 is termed ISGylation. ISGylation requires three distinct enzymes: an ATP-dependent activating enzyme for ISG15 (E1, Ube1L), an ISG15-specific carrier protein/conjugating enzyme (E2, UbcH8) and several ISG15-specific conjugating enzymes (E3, Herc5). Expression of ISG15 and the conjugation of ISG15 to target proteins are strongly promoted by IFN- α/β treatment, dsRNA, and viral or bacterial infection (6). In addition to existing in its conjugated form, ISG15 is present in an unconjugated form intracellularly and is also released into the extracellular space (7).

HCV infection triggers intrahepatic synthesis of several ISGs in patients (8, 9). ISG15 is one of the ISGs which are highly induced by HCV infection (10). ISG15 has

already been shown to function as an antiviral factor against numerous virus infections, but to act as a proviral factor in HCV infection (11-14). Conversely, other groups have reported that ISG15 mediates antiviral activity against HCV infection, and that this activity may occur through the proteasomal degradation pathway and in a Herc5-independent manner (15, 16). Therefore, the potential roles of ISG15 in HCV infection are still under debate. Evidence is accumulating that unconjugated ISG15 may modulate viral infection independent of ISGylation (17-19). In the present study, we attempted to gain further insight into the role of ISGylation-independent, unconjugated ISG15 in HCV infection.

First, we sought to determine whether ISG15 physically interacts with any HCV proteins in Huh-7 cells. For this purpose, Huh-7 cells were co-expressed with a series of myc-His₆-tagged HCV proteins together with HA-ISG15 (Fig. 1). Immunoprecipitation analysis revealed that NS5A-myc-His₆ was co-immunoprecipitated with HA-ISG15 (Fig. 1a, upper panel, lane 3). On the other hand, FLAG-ISG15 was co-immunoprecipitated with HA-NS5A (Fig. 1b, upper panel, lane 3). Although only NS3-myc-His₆ was co-immunoprecipitated with HA-ISG15 (Fig. 1a, upper panel, lane 5), HA-NS3 was not co-immunoprecipitated with FLAG-ISG15 (Fig. 1b, upper panel, lane 5) when NS3 was expressed as NS3/4A in the cells (Fig. 1b). NS4A is known to be responsible for the appropriate subcellular localization of the NS3-4A complex at the organelle's membranes (20). Thus, we concluded that NS3 may not interact with ISG15 in the proper subcellular localization. In addition, neither myc-His₆-tagged NS4B, NS5B or p7 protein was co-immunoprecipitated with HA-ISG15 (Fig. 1a, upper panel, lane 6; Fig. 1c, upper panel, lane 5; Fig. 1e, upper panel, lane 3). As shown in Fig. 1d, the HCV core protein was not co-immunoprecipitated with HA-ISG15 (Fig. 1d, upper

panel, lane 3). These results suggest that ISG15 specifically interacts with the HCV NS5A protein.

To determine whether endogenous ISG15 interacts with HCV NS5A in HCV-replicating cells, the HCV subgenomic replicon cells, Huh 9-13 (21), were treated with 1.5×10^3 IU/mL IFN- α (Sigma-Aldrich, St. Louis, MO) for 18 hr. Endogenous ISG15 was co-immunoprecipitated with NS5A protein using anti-NS5A monoclonal antibody (mAb) but not with control mouse IgG (Fig. 1f, top panel, lanes 4 and 3). These results suggest that endogenous ISG15 indeed interacts with the HCV NS5A protein expressed from HCV-replicating cells.

To determine whether the interaction between unconjugated ISG15 and HCV NS5A protein is ISGylation-independent, we constructed ISG15 mutants with amino acid substitutions in the C-terminal LRLRGG motif (mutated from LRLRGG to LRLRAA, referred to as ISG15 G156/157A) or deleting LRLRGG motif (referred to as ISG15 Δ 152-157) (Fig. 2a). These two ISG15 mutants are incapable of forming ISG15-conjugates (22).

To determine whether these two ISG15 mutants can interact with the NS5A protein, HA-NS5A was co-expressed with either FLAG-ISG15, FLAG-ISG15 Δ 152-157, or FLAG-ISG15 G156/157A in Huh-7 cells, respectively. The cell extracts were subjected to immunoprecipitation with anti-FLAG mAb or anti-HA polyclonal antibody (pAb). As shown in Fig. 2b, HA-tagged NS5A was co-immunoprecipitated with FLAG-ISG15 Δ 152-157 and FLAG-ISG15 G156/157A (Fig. 2b, top panel, lanes 3 and 4). These results suggest that unconjugated ISG15 interacts with HCV NS5A in an ISGylation-independent manner.

To examine the subcellular localization of ISG15 and NS5A using a confocal

microscope, the HCV 1b full-length replicon (RCYM1) (23) was expressed with a vector control, FLAG-ISG15, FLAG-ISG15 Δ 152-157, or FLAG-ISG15 G156/157A, respectively. The immunofluorescence staining revealed that unconjugated ISG15 was colocalized with NS5A in the perinuclear region in HCV replicon cells (Fig. 2c, merged panels). Moreover, to verify the colocalization of ISG15 and NS5A, we performed an *in situ* proximity ligation assay (PLA) using Duolink In Situ PLA (Sigma-Aldrich) (24). Red fluorescent spots were observed in HCV replicon cells expressing FLAG-ISG15, FLAG-ISG15 G156/157A, or FLAG-ISG15 Δ 152-157, whereas no signal was detected in the vector control (Fig. 2d). These results suggest that unconjugated ISG15 colocalizes with the HCV NS5A protein in an ISGylation-independent manner.

Finally, to determine the ISG15-binding site on NS5A protein, we used a series of NS5A-deletion mutants constructed previously (25, 26). The series of NS5A-deletion mutants were co-expressed with FLAG-ISG15 in Huh-7 cells (Fig. 3a). Immunoprecipitation analysis was performed with anti-FLAG antibody, followed by immunoblotting with anti-HA PAb, anti-Myc MAb, or anti-ISG15 MAb, respectively (Fig. 3b and 3c). Among the HA-tagged NS5A mutants shown in Fig. 3a, the N-terminal-deletion mutants lacking the region from aa 357 to 447, aa 250 to 447, or aa 214 to 447 were not co-immunoprecipitated with FLAG-tagged ISG15 (Fig. 3, upper panel, lanes 7-9), whereas other C-terminal-deletion mutants were co-immunoprecipitated with FLAG-tagged ISG15 (Fig 3b, upper panel, lanes 2-6). Co-immunoprecipitation analysis using the Myc-His₆-tagged NS5A 1-126, NS5A 1-146, or NS5A 1-447 revealed that these NS5A mutants were co-precipitated with FLAG-ISG15 (Fig. 3c, upper panel, lanes 6-8). These results suggest that the region from aa 1 to 126 within the domain-I of NS5A is responsible for the interaction with

unconjugated ISG15.

Taken together, these findings suggest that unconjugated ISG15 interacts with NS5A in transfected cells and in HCV-replicating cells. Moreover, the immunofluorescence staining and the *in situ* PLA assay also demonstrated the colocalization of unconjugated ISG15 with the HCV NS5A protein in an ISGylation-independent manner. We mapped the region ranging from aa 1 to 126 within the domain I of NS5A, which was the region found to be responsible for the interaction between unconjugated ISG15 and NS5A. As far as we know, this is the first report to show the potential role of unconjugated ISG15 in HCV infection.

After the discovery of ISG15, much effort was invested in studying ISGylation. Recently, however, it was reported that ISG15^{-/-} mice are more susceptible to Chikungunya virus (CHIKV) infection, whereas Ube1L^{-/-} mice display similar susceptibility to CHIKV as wild-type mice, suggesting a conjugation-independent role for ISG15 on the CHIKV infection (17). Other studies have also showed that an unconjugated ISG15 interacts with E3 ubiquitin ligase Nedd4 and blocks the Nedd4-mediated ubiquitylation of Ebola virus VP40 for the efficient viral particle release (18, 19). These results may imply that viral infection is modulated through unconjugated ISG15. Our results raise the possibility that unconjugated ISG15 may have some roles in the HCV life cycle and in viral pathogenesis.

HCV NS5A is a proline-rich, hydrophilic phosphoprotein that functions as a key regulator of HCV RNA replication and assembly (27). NS5A can interact with a wide variety of host cell proteins, such as HNF-1 α , ApoA1, 2'-5' OAS, La, HSP27, PKR, FKBP8, and OTUD7B (25, 26, 28-34). Here we demonstrated a specific interaction between unconjugated ISG15 and NS5A via the domain I of NS5A (Fig. 3). To elucidate

the physiological significance of the unconjugated ISG15-NS5A interaction, it will be necessary to examine whether unconjugated ISG15 is involved in the viral replication, assembly, or immune evasion from the host innate immune response. Our results suggest that unconjugated ISG15 as well as ISGylation forms of NS5A are present in HCV-replicating cells. Co-existence of unconjugated- and conjugated-ISG15 interactions with NS5A may affect versatile functions of NS5A protein in HCV replication and in HCV pathogenesis. Further analysis will be required to elucidate the biological significance of the interplay between unconjugated- and conjugated-ISG15.

Collectively, we provided evidence indicating that unconjugated ISG15 interacts with HCV NS5A in an ISGylation-independent manner. Understanding the role of unconjugated ISG15 in HCV infection may shed new light on the molecular mechanisms of the HCV life cycle and HCV pathogenesis.

ACKNOWLEDGMENTS

We thank Ms. Yasuko Kozaki for secretarial work. This study was supported in part by grants-in-aid for research on hepatitis from the Ministry of Health, Labour, and Welfare, Japan, and the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan, JSPS KAKENHI, and Japan Agency for Medical Research and Development (AMED).

DISCLOSURE

The authors declare that they have no conflicting interests.

REFERENCES

1. Lavanchy D. (2009) The global burden of hepatitis C. *Liver Int* **29 Suppl 1**: 74-81.
2. Jacobson I.M., Davis G.L., El-Serag H., Negro F., Trepo C. (2010) Prevalence and challenges of liver diseases in patients with chronic hepatitis C virus infection. *Clin Gastroenterol Hepatol* **8**: 924-33; quiz e117.
3. Moradpour D., Penin F., Rice C.M. (2007) Replication of hepatitis C virus. *Nat Rev Microbiol* **5**: 453-63.
4. He Y., Staschke K.A., Tan S.-L. (2006) HCV NS5A: A Multifunctional Regulator of Cellular Pathways and Virus Replication. In: SI T., ed. eds. *Hepatitis C Viruses: Genomes and Molecular Biology*. Norfolk (UK): Horizon Bioscience, pp. 267-92.
5. Haas A.L., Ahrens P., Bright P.M., Ankel H. (1987) Interferon induces a 15-kilodalton protein exhibiting marked homology to ubiquitin. *J Biol Chem* **262**: 11315-23.
6. Loeb K.R., Haas A.L. (1992) The interferon-inducible 15-kDa ubiquitin homolog conjugates to intracellular proteins. *J Biol Chem* **267**: 7806-13.
7. D'cunha J., Ramanujam S., Wagner R.J., Witt P.L., Knight E., Jr., Borden E.C. (1996) In vitro and in vivo secretion of human ISG15, an IFN-induced immunomodulatory cytokine. *J Immunol* **157**: 4100-8.
8. Mihm S., Frese M., Meier V., Wietzke-Braun P., Scharf J.G., Bartenschlager R., Ramadori G. (2004) Interferon type I gene expression in chronic hepatitis C. *Lab Invest* **84**: 1148-59.
9. Sarasin-Filipowicz M., Oakeley E.J., Duong F.H., Christen V., Terracciano L., Filipowicz W., Heim M.H. (2008) Interferon signaling and treatment outcome in chronic hepatitis C. *Proc Natl Acad Sci U S A* **105**: 7034-9.
10. Chen L., Borozan I., Feld J., Sun J., Tannis L.L., Coltescu C., Heathcote J., Edwards A.M., Mcgilvray I.D. (2005) Hepatic gene expression discriminates responders and nonresponders in treatment of chronic hepatitis C viral infection. *Gastroenterology* **128**: 1437-44.
11. Broering R., Zhang X., Kottlil S., Trippler M., Jiang M., Lu M., Gerken G., Schlaak J.F. (2010) The interferon stimulated gene 15 functions as a proviral

- factor for the hepatitis C virus and as a regulator of the IFN response. *Gut* **59**: 1111-9.
12. Chen L., Sun J., Meng L., Heathcote J., Edwards A.M., Mcgilvray I.D. (2010) ISG15, a ubiquitin-like interferon-stimulated gene, promotes hepatitis C virus production in vitro: implications for chronic infection and response to treatment. *J Gen Virol* **91**: 382-8.
 13. Arnaud N., Dabo S., Akazawa D., Fukasawa M., Shinkai-Ouchi F., Hugon J., Wakita T., Meurs E.F. (2011) Hepatitis C virus reveals a novel early control in acute immune response. *PLoS Pathog* **7**: e1002289.
 14. Real C.I., Megger D.A., Sitek B., Jahn-Hofmann K., Ickenstein L.M., John M.J., Walker A., Timm J., Kuhlmann K., Eisenacher M., Meyer H.E., Gerken G., Broering R., Schlaak J.F. (2013) Identification of proteins that mediate the pro-viral functions of the interferon stimulated gene 15 in hepatitis C virus replication. *Antiviral Res* **100**: 654-61.
 15. Kim M.J., Yoo J.Y. (2010) Inhibition of hepatitis C virus replication by IFN-mediated ISGylation of HCV-NS5A. *J Immunol* **185**: 4311-8.
 16. Domingues P., Bamford C.G., Boutell C., Mclauchlan J. (2015) Inhibition of hepatitis C virus RNA replication by ISG15 does not require its conjugation to protein substrates by the HERC5 E3 ligase. *J Gen Virol* **96**: 3236-42.
 17. Werneke S.W., Schilte C., Rohatgi A., Monte K.J., Michault A., Arenzana-Seisdedos F., Vanlandingham D.L., Higgs S., Fontanet A., Albert M.L., Lenschow D.J. (2011) ISG15 is critical in the control of Chikungunya virus infection independent of Ube1L mediated conjugation. *PLoS Pathog* **7**: e1002322.
 18. Okumura A., Pitha P.M., Harty R.N. (2008) ISG15 inhibits Ebola VP40 VLP budding in an L-domain-dependent manner by blocking Nedd4 ligase activity. *Proc Natl Acad Sci U S A* **105**: 3974-9.
 19. Malakhova O.A., Zhang D.E. (2008) ISG15 inhibits Nedd4 ubiquitin E3 activity and enhances the innate antiviral response. *J Biol Chem* **283**: 8783-7.
 20. Wolk B., Sansonno D., Krausslich H.G., Dammacco F., Rice C.M., Blum H.E., Moradpour D. (2000) Subcellular localization, stability, and trans-cleavage competence of the hepatitis C virus NS3-NS4A complex expressed in

- tetracycline-regulated cell lines. *J Virol* **74**: 2293-304.
21. Lohmann V., Korner F., Koch J., Herian U., Theilmann L., Bartenschlager R. (1999) Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science (New York, NY)* **285**: 110-3.
 22. Lenschow D.J., Giannakopoulos N.V., Gunn L.J., Johnston C., O'guin A.K., Schmidt R.E., Levine B., Virgin H.W.T. (2005) Identification of interferon-stimulated gene 15 as an antiviral molecule during Sindbis virus infection in vivo. *J Virol* **79**: 13974-83.
 23. Murakami K., Ishii K., Ishihara Y., Yoshizaki S., Tanaka K., Gotoh Y., Aizaki H., Kohara M., Yoshioka H., Mori Y., Manabe N., Shoji I., Sata T., Bartenschlager R., Matsuura Y., Miyamura T., Suzuki T. (2006) Production of infectious hepatitis C virus particles in three-dimensional cultures of the cell line carrying the genome-length dicistronic viral RNA of genotype 1b. *Virology* **351**: 381-92.
 24. Hayashi M., Deng L., Chen M., Gan X., Shinozaki K., Shoji I., Hotta H. (2016) Interaction of the hepatitis B virus X protein with the lysine methyltransferase SET and MYND domain-containing 3 induces activator protein 1 activation. *Microbiology and immunology* **60**: 17-25.
 25. Matsui C., Shoji I., Kaneda S., Sianipar I.R., Deng L., Hotta H. (2012) Hepatitis C virus infection suppresses GLUT2 gene expression via downregulation of hepatocyte nuclear factor 1alpha. *J Virol* **86**: 12903-11.
 26. Matsui C., Rosalyn Sianipar I., Minami N., Deng L., Hotta H., Shoji I. (2015) A single-amino-acid mutation in hepatitis C virus NS5A disrupts physical and functional interaction with the transcription factor HNF-1alpha. *J Gen Virol* **96**: 2200-5.
 27. Macdonald A., Harris M. (2004) Hepatitis C virus NS5A: tales of a promiscuous protein. *J Gen Virol* **85**: 2485-502.
 28. Shi S.T., Polyak S.J., Tu H., Taylor D.R., Gretch D.R., Lai M.M. (2002) Hepatitis C virus NS5A colocalizes with the core protein on lipid droplets and interacts with apolipoproteins. *Virology* **292**: 198-210.
 29. Taguchi T., Nagano-Fujii M., Akutsu M., Kadoya H., Ohgimoto S., Ishido S., Hotta H. (2004) Hepatitis C virus NS5A protein interacts with 2',5'-oligoadenylate synthetase and inhibits antiviral activity of IFN in an IFN

- sensitivity-determining region-independent manner. *J Gen Virol* **85**: 959-69.
30. Houshmand H., Bergqvist A. (2003) Interaction of hepatitis C virus NS5A with La protein revealed by T7 phage display. *Biochem Biophys Res Commun* **309**: 695-701.
 31. Choi Y.W., Tan Y.J., Lim S.G., Hong W., Goh P.Y. (2004) Proteomic approach identifies HSP27 as an interacting partner of the hepatitis C virus NS5A protein. *Biochem Biophys Res Commun* **318**: 514-9.
 32. Gale M.J., Jr., Korth M.J., Tang N.M., Tan S.L., Hopkins D.A., Dever T.E., Polyak S.J., Gretch D.R., Katze M.G. (1997) Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the nonstructural 5A protein. *Virology* **230**: 217-27.
 33. Okamoto T., Nishimura Y., Ichimura T., Suzuki K., Miyamura T., Suzuki T., Moriishi K., Matsuura Y. (2006) Hepatitis C virus RNA replication is regulated by FKBP8 and Hsp90. *Embo j* **25**: 5015-25.
 34. Sianipar I.R., Matsui C., Minami N., Gan X., Deng L., Hotta H., Shoji I. (2015) Physical and functional interaction between hepatitis C virus NS5A protein and ovarian tumor protein deubiquitinase 7B. *Microbiol Immunol* **59**: 466-76.

FIGURE LEGENDS

Fig. 1. ISG15 specifically interacts with HCV NS5A protein.

HA-ISG15 was co-expressed with NS3-myc-His₆, NS4B-myc-His₆, or NS5A-myc-His₆ (a), NS5B-myc-His₆ (c), P7-myc-His₆ (e) or FLAG-Core (d) in Huh-7 cells, and subjected to immunoprecipitation (IP) and immunoblotting (IB) using the indicated antibodies. FLAG-ISG15 was co-expressed with HA-NS5A or HA-NS3/4A (b), and subjected to IP and IB using the indicated antibodies. (f) HCV subgenomic replicon cells (Con1, genotype 1b) were treated with 1.5×10^3 IU/ml recombinant IFN- α for 18 hr, and then the cell lysates were subjected to IP with anti-NS5A mAb or control mouse IgG, followed by IB with anti-NS5A mAb or anti-ISG15 mAb, respectively.

Fig. 2. ISG15 interacts with HCV NS5A protein in an ISGylation-independent manner.

(a) Schematic representation of ISG15 and its mutants (indicated as ISG15 Δ 152-157 and ISG15 G156/157A). UBL, Ubiquitin-like domain; UBL, LRLRGG, Leu-Arg-Leu-Arg-Gly-Gly. (b) HA-NS5A was co-expressed with FLAG-ISG15, FLAG-ISG15 Δ 152-157, and FLAG-ISG15 G156/157A in Huh-7 cells, followed by immunoprecipitation (IP) with anti-FLAG mAb, coupled with immunoblotting (IB) with anti-HA rabbit pAb or anti-FLAG mAb, respectively. The arrow indicates co-immunoprecipitated HA-NS5A. The asterisk indicates a non-specific band. (c) HCV full-genome replicon cells (Con1, genotype 1b, termed as RCYM1) were transfected with pCAG-FLAG-ISG15, pCAG-FLAG-ISG15 Δ 152-157, or pCAG-FLAG-ISG15 G156/157A. The cells were fixed, permeabilized, and stained with anti-FLAG rabbit pAb and anti-NS5A mouse mAb, followed by incubation with anti-rabbit Alexa Fluor®

488 and anti-mouse Alexa Fluor® 594, respectively. Nuclei were stained with Hoechst 33342. (d) *In situ* proximity ligation assay (PLA). HCV replicon RCYM1 cells were transfected with either FLAG-ISG15, FLAG-ISG15 Δ 152-157, or FLAG-ISG15 G156/157A. Cells were fixed, permeabilized, and incubated with anti-NS5A mouse mAb and anti-FLAG rabbit pAb, followed by staining with PLA. The red fluorescent spots indicate the interactions with NS5A and ISG15 or mutant ISG15. Nuclei were stained with Hoechst 33342.

Fig. 3. Mapping of ISG15-binding domain on the NS5A protein.

(a) Schematic representation of the NS5A and deletion mutants. The N-terminal- and C-terminal-deletion mutants of NS5A are fused to the HA-tag (gray region, a to h) or myc-His₆-tag (lattice region, i and j), respectively. Closed boxes represent proteins that are bound specifically to the unconjugated ISG15 protein and open boxes represent those that are not bound. (b and c) FLAG-ISG15 was co-expressed with one of the above-described HA-NS5A-deletion mutants in Huh7 cells. Cell lysates were immunoprecipitated (IP) with anti-FLAG antibody coupled with immunoblotting (IB) with anti-HA rabbit pAb, anti-c-myc mouse mAb, or anti-ISG15 mouse mAb, respectively.




Figure 1. Minami et al.

Figure 1

220x151mm (300 x 300 DPI)




Figure 2. Minami et al.

Figure 2

196x154mm (300 x 300 DPI)




Figure 3. Minami et al.

Figure 3

192x145mm (300 x 300 DPI)