



Glucagon secretions are impaired in patients with fulminant type 1 diabetes

Komada, Hisako
Hirota, Yushi
Ogawa, Wataru

(Citation)

Journal of Diabetes Investigation, 10(3):866-867

(Issue Date)

2019-05

(Resource Type)

journal article

(Version)

Version of Record

(Rights)

© 2019 The Authors. Journal of Diabetes Investigation published by Asian Association for the Study of Diabetes (AASD) and John Wiley & Sons Australia, Ltd.
This is an open access article under the terms of the Creative Commons Attribution - NonCommercial License, which permits use, distribution and reproduction in any mediu...

(URL)

<https://hdl.handle.net/20.500.14094/90006078>



Glucagon secretions are impaired in patients with fulminant type 1 diabetes

We have read with great interest the articles by Murase-Mishibe *et al.*¹ and Takahashi *et al.*² studying glucagon secretion in fulminant type 1 diabetes patients. Murase-Mishibe *et al.* investigated plasma glucagon levels of fulminant type 1 diabetes patients ($n = 3$) and acute-onset type 1 diabetes patients ($n = 6$) during a meal tolerance test, and stated in their study that “no defect in glucagon secretion was observed” in fulminant type 1 diabetes without providing the information for statistical analysis. Takahashi *et al.* investigated plasma glucagon levels of fulminant type 1 diabetes patients ($n = 5$) and age- and diabetes duration-matched type 1 diabetes patients ($n = 5$) during an arginine challenge test (ACT), and stated that “the curve of glucagon levels were similar between the two groups.” It is again unclear how the authors analyzed the similarity between the groups.

We have recently investigated glucagon secretion during ACT in various types of diabetes patients, including four fulminant type 1 diabetes patients, five slowly progressive insulin-dependent diabetes patients, 18 type 1 diabetes patients, 40 type 2 diabetes patients and 16 pancreatic diabetes (PD) patients, and showed that the area under the curve of glucagon levels in fulminant type 1 diabetes patients was significantly smaller than those of type 1 diabetes patients, but was similar to PD patients³. We now show the time course of our previous data of glucagon secretion during ACT in these patients (Figure 1). The analysis of the




Figure 1 | Plasma glucose, serum glucagon and serum C-peptide during an arginine challenge test in various type of diabetes patients. Characteristics of the study participants and study methods were previously described³. Data are shown as the mean \pm standard deviation. CPR, C-peptide reactivity; FT1D, fulminant type 1 diabetes; IRG, immunoreactive glucagon; PD, pancreatic diabetes; SP1DD, slowly progressive insulin-dependent diabetes; T1D, acute-onset type 1 diabetes; T2D, type 2 diabetes.

*Corresponding author. Yushi Hirota

Tel.: +81-78-382-5861

Fax: +81-78-382-2080

E-mail address: hirota@med.kobe-u.ac.jp

Received 12 February 2019; revised 5 March 2019;

accepted 8 March 2019

time course data with repeated-measures ANOVA showed that the glucagon levels of fulminant type 1 patients diabetes were significantly lower than that of type 1 diabetes patients ($P = 0.003$), but similar to PD patients ($P = 0.397$), consistent with our previous analysis³. Given that Takahashi's group previously showed that glucagon secretion in patients with fulminant type 1 diabetes was substantially decreased within 6 months after the onset of the disease⁴, the two groups of investigators discussed whether the duration of the disease could affect the ability of glucagon secretion^{1,2}. The duration of the disease of our fulminant type 1 diabetes patients was 1, 6, 12 and 100 months (29.8 ± 47.0 months).

The numbers of study participants with fulminant type 1 diabetes in our (4) and the two investigator groups (3 and 5) were small. It is obvious that a study with a larger number of participants is required to obtain a definite conclusion. It is possible that the difference in assays is attributable to the apparent inconsistency of our and the other groups' results. We used a radioimmunoassay that has been used conventionally to determine the levels of glucagon, and evidence suggests that a newly launched sandwich enzyme-linked immunoassay more accurately reflects the hormone levels⁵. Takahashi *et al.* showed that whereas the absolute values determined by the radioimmunoassay and by the enzyme-linked immunoassay were

different, the relative changes in these two assays were similar⁴. Metabolic testing of our study (ACT) and that of Murase-Mishibe *et al.* (a meal tolerance test) differed, which also might be attributable to the inconsistency. Furthermore, whereas Murase-Mishibe *et al.* carried out the test with healthy volunteers, we only analyzed patients with various types of diabetes.



An interesting finding in our study is that, in contrast to glucagon, secretion of C-peptide in fulminant type 1 diabetes patients is similar to that of type 1 diabetes patients ($P = 0.284$), but smaller than that of PD patients ($P = 0.020$; Figure 1). The difference in the extent of the impairment of the two pancreatic hormone secretions might be related to clinical characteristics of the three types of diabetes primarily caused by insufficient insulin secretion. Again, further analysis with a larger number of participants warrants clarifying this point.

ACKNOWLEDGMENTS

We thank Yoko Okuno and Kazuhiko Sakaguchi for data collection.

DISCLOSURE

The authors declare no conflict of interest.

Hisako Komada, Yushi Hirota* ,
Wataru Ogawa 

Department of Internal Medicine,
Division of Diabetes and Endocrinology,

Kobe University Graduate School of
Medicine, Kobe, Japan

REFERENCES

1. Murase-Mishiba Y, Bessho-Tachibana M, Imagawa A. "Preserved" glucagon secretion in fulminant type 1 diabetes. *J Diabetes Investig* 2019; 10: 186–187.
2. Takahashi N, Chujo D. Response to "Preserved" glucagon secretion in fulminant type 1 diabetes. *J Diabetes Investig* 2019; 10: 188–189.
3. Komada H, Hirota Y, Sakaguchi K, *et al.* Impaired glucagon secretion in patients with fulminant type 1 diabetes mellitus. *Endocrine* 2019; 63: 476–479.
4. Takahashi N, Chujo D, Kajio H, *et al.* Contribution of pancreatic α -cell function to insulin sensitivity and glycemic variability in patients with type 1 diabetes. *J Diabetes Investig* 2018. <https://doi.org/10.1111/jdi.12949>
5. Miyachi A, Kobayashi M, Mieno E, *et al.* Accurate analytical method for human plasma glucagon levels using liquid chromatography-high resolution mass spectrometry: comparison with commercially available immunoassays. *Anal Bioanal Chem* 2017; 409: 5911–5918.

Doi: 10.1111/jdi.13041