



Quasinormal modes of p-forms in spherical black holes

Yoshida, Daisuke
Soda, Jiro

(Citation)

Physical Review D, 99(4):044054-044054

(Issue Date)

2019-02-26

(Resource Type)

journal article

(Version)

Version of Record

(Rights)

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

(URL)

<https://hdl.handle.net/20.500.14094/90006277>



Quasinormal modes of p -forms in spherical black holes

Daisuke Yoshida^{*} and Jiro Soda[†]

Department of Physics, Kobe University, Kobe 657-8501, Japan



(Received 29 January 2019; published 26 February 2019)

We study the quasinormal modes of p -form fields in spherical black holes in D dimensions. Using the spherical symmetry of the black holes and gauge symmetry, we show the p -form field can be expressed in terms of the coexact p -form and the coexact $(p - 1)$ -form on the sphere S^{D-2} . These variables allow us to find the master equations. By utilizing the S -deformation method, we explicitly show the stability of p -form fields in the spherical black hole spacetime. Moreover, using the WKB approximation, we calculate the quasinormal modes of the p -form fields in $D(\leq 10)$ dimensions.

DOI: [10.1103/PhysRevD.99.044054](https://doi.org/10.1103/PhysRevD.99.044054)

I. INTRODUCTION

Black holes in general relativity are important from various perspectives. In fact, they are sources of gravitational waves, they provide a way to test general relativity in the strong gravity regime, and they can be a key to quantum gravity. Before discussing this physics, we have to show the stability of black holes. In fact, the stability of black holes is often nontrivial. Historically, the stability of black holes has been studied since the seminal papers by Regge and Wheeler [1] and Zerilli [2]; for examples, see [3–5]. From the point of view of a unified theory such as string theory, it is natural to consider black holes in higher dimensions. Indeed, higher-dimensional black holes may be created at the accelerator such as the LHC [6]. Thus, the stability analysis is also generalized to higher dimensions [7–10]. In higher dimensions, however, Einstein's general relativity is not a unique possibility. Rather, Lovelock gravity is natural in higher dimensions [11,12]. Therefore, the stability of black holes in Lovelock gravity has been studied [13–18]. Moreover, in contrast to four-dimensional general relativity where only scalar, electromagnetic and gravitational fields can reside in the black hole spacetime, there exist p -form fields in higher dimensions. To our best knowledge, no work of p -form fields in higher dimensional black hole spacetime has been done. The purpose of this paper is to study the stability of p -form fields in black hole spacetime and obtain quasinormal modes of p -form fields.

To study the behavior of various physical fields in spherical black holes, we must derive the master equations. For this purpose, we express a p -form field in terms of a coexact p -form and coexact $(p - 1)$ -form on the sphere and derive the master equation for each component. If the effective potential in the master equation is positive outside the event horizon of black holes, the p -form field is stable [19]. However, it turns out that the effective potential for a p -form field has a negative region for some parameters. This region may cause the instability of p -form fields in spherical black holes. Nevertheless, we succeed in proving the stability of p -form fields using the S -deformation method [8].

Given the stability, we can calculate the quasinormal modes of p -form fields in the black hole spacetime. We use the WKB method [20–22] to calculate the quasinormal modes of p -form fields in $D(\leq 10)$ dimensions. Since a p -form field has two components, there are two quasinormal modes, namely, one for each component. It is shown that the quasinormal modes of the p -form field in D dimensions reflect duality relations.

The organization of the paper is as follows. In Sec. II, we review the properties of p -form fields. In particular, we count the physical degrees of freedom (d.o.f.) of a p -form field. In Sec. III, we consider p -form fields in spherical black holes. We represent a p -form field by coexact form fields on the sphere. We also discuss spherical harmonics. In Sec. IV, we obtain the master equations for the p -form field in spherical black holes in arbitrary dimensions. The master variables are a coexact p -form and a coexact $(p - 1)$ -form on the sphere. We check their d.o.f. match to the physical d.o.f. of a p -form field. We also find useful duality relations for the effective potentials. It turns out that the effective potential has a negative region for some cases. In Sec. V, therefore, we have explicitly verified the stability of the p -form field using the S -deformation technique. In Sec. VI, we also investigated the quasinormal modes of

^{*}dice-k.yoshida@stu.kobe-u.ac.jp

[†]jiro@phys.sci.kobe-u.ac.jp

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

p -form fields using the WKB approximation. Section VII is devoted to conclusion.

II. BASICS OF p -FORM FIELDS

In this section, we introduce the action for p -form fields and explain the gauge invariance in the system. We also calculate physical d.o.f. of a p -form field. We refer the reader to the textbook [23] for a more detailed explanation of the p -forms on manifolds.

The action of the p -form field is given by

$$S = -\frac{1}{2} \int \mathbf{F} \wedge * \mathbf{F} \\ = -\frac{1}{2(p+1)!} \int d^D x \sqrt{-g} F_{a_1 \dots a_{p+1}} F^{a_1 \dots a_{p+1}}, \quad (1)$$

where we used the Hodge operator $*$. Here, D is the dimension of the spacetime. We defined the p -form field A_p as follows:

$$A_p \equiv \frac{1}{p!} A_{a_1 \dots a_p} dx^{a_1} \wedge dx^{a_2} \wedge \dots \wedge dx^{a_p} \quad (2)$$

and the field strength \mathbf{F} is defined by

$$\mathbf{F} \equiv dA. \quad (3)$$

The operator d is the exterior derivative satisfying the identity

$$d^2 = 0. \quad (4)$$

Using the Hodge operator, we can define the coderivative δ as

$$\delta \equiv (-1)^{D(p+1)+1} * d *. \quad (5)$$

Note that the coderivative also satisfies the identity

$$\delta^2 = 0. \quad (6)$$

The equations of motion for the form field can be deduced as

$$d\mathbf{F} = 0 \quad \text{and} \quad \delta\mathbf{F} = 0. \quad (7)$$

This system has the symmetry under the dual transformation

$$\tilde{\mathbf{F}} = *\mathbf{F}. \quad (8)$$

Because of this symmetry, we do not need to consider a p -form field with the rank higher than

$$p_{\max} = \left\lfloor \frac{D}{2} \right\rfloor - 1 = \left\lfloor \frac{n}{2} \right\rfloor. \quad (9)$$

Here, $\lfloor \cdot \rfloor$ denotes the Gauss symbol. So, we concentrate on the form fields $A_p (0 \leq p \leq p_{\max})$ in arbitrary dimensions.

The p -form field has the invariance under the gauge transformation

$$A_p \rightarrow \tilde{A}_p = A_p + d\Xi_{p-1} \quad (10)$$

with an arbitrary $(p-1)$ -form field Ξ_{p-1} . This is because the field strength is defined by (3). The gauge parameter Ξ_{p-1} itself has the degeneracy

$$\Xi_{p-1} \rightarrow \tilde{\Xi}_{p-1} = \Xi_{p-1} + d\Xi_{p-2} \quad (11)$$

with an arbitrary $(p-2)$ -form field Ξ_{p-2} . Hence, in order to count the physical d.o.f. of the p -form field A_p , we need to take into account these degrees generated by the transformations $\Xi_{p-1}, \Xi_{p-2}, \dots, \Xi_0$. Taking into account that components $A_{0a_2 \dots a_p}$ are not dynamical, the formula for physical d.o.f. is given by

$$_{D-1}C_p - _{D-1}C_{p-1} + _{D-1}C_{p-2} - \dots = _{D-2}C_p. \quad (12)$$

For example, in $D = 4$, a 2-form field has one physical d.o.f.

III. p -FORM FIELDS IN SPHERICAL BLACK HOLES

In this subsection, we study the decomposition of the p -form field in black hole spacetime in terms of form fields on the sphere. Then, we eliminate some components using gauge transformations. We also discuss eigenvalues of spherical harmonics for p -forms.

In general relativity, the spherical black hole is known as the Schwarzschild black hole. It is not difficult to generalize the Schwarzschild black hole to higher dimensions $D > 4$, and the solutions are called a Schwarzschild-Tangherlini black hole [24] expressed by the metric

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2 q_{AB} dx^A dx^B, \quad (13)$$

where $f(r)$ is given by

$$f(r) = 1 - \frac{\mu}{r^{n-1}}, \quad (14)$$

and n is defined as $n \equiv D - 2$. Here, q_{AB} is the metric of the sphere S^n and the spherical coordinate is expressed by

$$x^A = (\theta^1, \dots, \theta^n). \quad (15)$$

A. Decomposition of a p -form field in terms of coexact form fields on the sphere

We use the notation \hat{A}_p to denote a p -form field on the sphere, that is, \hat{A}_p is written by

$$\hat{A}_p = A_{A_1 \dots A_p} \mathcal{D}^{A_1 \dots A_p}, \quad (16)$$

where

$$\mathcal{D}^{A_1 \dots A_p} \equiv dx^{A_1} \wedge \dots \wedge dx^{A_p}. \quad (17)$$

We can write down the components of A_p as follows:

$$\begin{aligned} A_p &= \frac{1}{(p-2)!} A_{t r A_1 \dots A_{p-2}} dt \wedge dr \wedge \mathcal{D}^{A_1 \dots A_{p-2}} \\ &+ \frac{1}{(p-1)!} A_{t A_1 \dots A_{p-1}} dt \wedge \mathcal{D}^{A_1 \dots A_{p-1}} \\ &+ \frac{1}{(p-1)!} A_{r A_1 \dots A_{p-1}} dr \wedge \mathcal{D}^{A_1 \dots A_{p-1}} + \frac{1}{p!} A_{A_1 \dots A_p} \mathcal{D}^{A_1 \dots A_p}. \end{aligned} \quad (18)$$

If we define the components of A_p as

$$U_{A_1 \dots A_{p-2}} \equiv A_{t r A_1 \dots A_{p-2}}, \quad (19)$$

$$V_{A_1 \dots A_{p-1}} \equiv A_{t A_1 \dots A_{p-1}}, \quad (20)$$

$$W_{A_1 \dots A_{p-1}} \equiv A_{r A_1 \dots A_{p-1}}, \quad (21)$$

$$X_{A_1 \dots A_p} \equiv A_{A_1 \dots A_p}, \quad (22)$$

then A_p is written by

$$A_p = dt \wedge dr \wedge \hat{U}_{p-2} + dt \wedge \hat{V}_{p-1} + dr \wedge \hat{W}_{p-1} + \hat{X}_p. \quad (23)$$

Thus, the p -form can be expressed by the one $(p-2)$ -form \hat{U}_{p-2} , the two $(p-1)$ -forms \hat{V}_{p-1} and \hat{W}_{p-1} and the one p -form \hat{X}_p . The identity

$${}_D C_p = {}_{D-2} C_{p-2} + 2 {}_{D-2} C_{p-1} + {}_{D-2} C_p \quad (24)$$

guarantees the matching of d.o.f.

Using the Hodge decomposition on the sphere S^n , a p -form field \hat{A}_p can be decomposed as

$$\hat{A}_p = \hat{d}\hat{A}_{p-1} + \hat{A}_p, \quad \text{for } 1 \leq p < n-1, \quad (25)$$

where we have introduced the coexact form $\hat{\delta}\hat{A}_p = 0$. From this decomposition theorem on the sphere, we can write

down a more useful expansion. In fact, for $p \geq 2$, we can express \hat{A}_p by the coexact form

$$\begin{aligned} \hat{A}_p &= \hat{d}\hat{A}_{p-1} + \hat{A}_p \\ &= \hat{d}(\hat{d}\hat{A}_{p-2} + \hat{A}_{p-1}) + \hat{A}_p \\ &= \hat{d}\hat{A}_{p-1} + \hat{A}_p. \end{aligned} \quad (26)$$

This result shows that the general form field on S^n is expressed by only the coexact form fields. Thus, the arbitrary p -form field A_p is given by

$$\begin{aligned} A_p &= dt \wedge dr \wedge \hat{U}_{p-2} + dt \wedge \hat{V}_{p-1} + dr \wedge \hat{W}_{p-1} + \hat{X}_p \\ &= dt \wedge dr \wedge (\hat{d}\hat{U}_{p-3} + \hat{U}_{p-2}) + dt \wedge (\hat{d}\hat{V}_{p-2} + \hat{V}_{p-1}) \\ &\quad + dr \wedge (\hat{d}\hat{W}_{p-2} + \hat{W}_{p-1}) + (\hat{d}\hat{X}_{p-1} + \hat{X}_p). \end{aligned} \quad (27)$$

B. Gauge fixing of p -form field A_p

The p -form field A_p has the gauge invariance under the transformation by B_{p-1} , that is,

$$A_p \rightarrow \tilde{A}_p = A_p + dB_{p-1}. \quad (28)$$

Now starting from the general expression for A_p in D dimensions, we can deduce the following result,

$$A_p = dt \wedge dr \wedge \hat{U}_{p-2} + dt \wedge \hat{V}_{p-1} + dr \wedge \hat{W}_{p-1} + \hat{X}_p, \quad (29)$$

by using the gauge transformation for A_p .

C. Spherical harmonics for coexact p -form field

We review the spherical harmonics of the p -form field following [25]. The Laplace-Beltrami operator is defined by

$$\hat{\Delta} \equiv \hat{\delta}\hat{d} + \hat{d}\hat{\delta}. \quad (30)$$

The spherical harmonics of the coexact p -form field $\hat{\mathcal{Y}}_p$ is defined by

$$\hat{\mathcal{Y}}_p = \frac{1}{p!} \mathcal{Y}_{A_1 \dots A_p} \mathcal{D}^{A_1 \dots A_p} \quad (31)$$

which satisfies

$$\hat{\delta}\hat{\mathcal{Y}}_p = 0 \quad (32)$$

and

$$\hat{\Delta}\hat{\mathcal{Y}}_p = \lambda_p \hat{\mathcal{Y}}_p. \quad (33)$$

The last equation becomes

$$\hat{\delta} \hat{d} \hat{\mathcal{Y}}_p = \lambda_p \hat{\mathcal{Y}}_p \quad (34)$$

by using the identity (6). Here, λ_p is given by

$$\lambda_p \equiv (\ell + p)(\ell + n - p - 1), \quad (35)$$

and ℓ is a positive integer, $\ell = 1, 2, \dots, \infty$. On the sphere S^n , the left-hand side of the equation becomes

$$\hat{\delta} \hat{d} \hat{\mathcal{Y}}_p = \frac{1}{p!} (-\tilde{\Delta} - p(n - p)) \mathcal{Y}_{A_1 \dots A_p} \mathcal{D}^{A_1 \dots A_p}. \quad (36)$$

Here, we defined $\tilde{\Delta}$ as

$$\tilde{\Delta} \mathcal{Y}_{A_1 \dots A_p} = \mathcal{Y}_{A_1 \dots A_p} :^A :_A. \quad (37)$$

Then, the spectrum of the coefficient of the p -form harmonics $\hat{\mathcal{Y}}_p$ is

$$-(\tilde{\Delta} - p(n - p)) \mathcal{Y}_{A_1 \dots A_p} = \lambda_p \mathcal{Y}_{A_1 \dots A_p}. \quad (38)$$

We rewrite this equation as follows:

$$\tilde{\Delta} \mathcal{Y}_{A_1 \dots A_p} = -\gamma_p^{(n)} \mathcal{Y}_{A_1 \dots A_p}, \quad (39)$$

where

$$\gamma_p^{(n)} \equiv \lambda_p - p(n - p). \quad (40)$$

With the harmonics $\hat{\mathcal{Y}}_p$, the coefficient of the general coexact p -form field \hat{A}_p can be expanded as

$$\mathcal{A}_{A_1 \dots A_p}(x) = \sum_{l, \sigma} \mathcal{A}_{\ell, \sigma}(t, r) \mathcal{Y}_{A_1 \dots A_p}^{\ell, \sigma}(x^A), \quad (41)$$

where σ denotes other indices to characterize the degeneracy. For simplicity, we denote $\mathcal{A}_{\ell, \sigma}(t, r)$ just as \mathcal{A} .

IV. MASTER EQUATIONS FOR p -FORM FIELD

In this section, we derive the master equations for the master variable Ψ in the Schrödinger form

$$-\ddot{\Psi} + \partial_x^2 \Psi - V\Psi = 0, \quad (42)$$

where V is the effective potential and x is the tortoise coordinate defined by

$$\frac{d}{dx} = f \frac{d}{dr}. \quad (43)$$

In Eq. (7), the first equation is trivially satisfied from the identity (4), and the second equation $\delta \mathbf{F} = 0$ in the coordinate basis is expressed as follows:

$$\frac{1}{\sqrt{-g}} \partial_a (\sqrt{-g} F^{aa_1 \dots a_p}) = 0. \quad (44)$$

This equation can be decomposed into four patterns.

The first pattern we consider is

$$\frac{1}{\sqrt{-g}} \partial_a (\sqrt{-g} F^{atr A_1 \dots A_{p-2}}) = 0. \quad (45)$$

Substituting the components into the above equation, we obtain

$$-\frac{1}{r^{2(p-1)}} (\tilde{\Delta} - (p-2)(n - (p-2))) \mathcal{U}^{A_1 \dots A_{p-2}} = 0. \quad (46)$$

This yields

$$(\ell + p - 2)(\ell + 1 + n - p) \mathcal{U}_{\ell, \sigma} = 0. \quad (47)$$

Since we are considering $p \geq 2$, and generally $n > p$, the quantity $(\ell + p - 2)(\ell + 1 + n - p)$ is always positive. Hence, the coefficient $\mathcal{U}_{\ell, \sigma}$ must vanish for all ℓ , namely,

$$\hat{\mathcal{U}}_{p-2} = 0. \quad (48)$$

Thus, the $(p-2)$ -form $\hat{\mathcal{U}}_{p-2}$ in Eq. (29) is not dynamical.

The second pattern is the following:

$$\frac{1}{\sqrt{-g}} \partial_a (\sqrt{-g} F^{at A_1 \dots A_{p-1}}) = 0. \quad (49)$$

It is easy to get

$$\begin{aligned} & \frac{1}{r^{n-2(p-1)}} f \partial_r (r^{n-2(p-1)} (\dot{\mathcal{W}} - \mathcal{V}')) \\ & + \frac{1}{r^2} (\gamma_{p-1}^{(n)} + (p-1)(n-p+1)) \mathcal{V} = 0. \end{aligned} \quad (50)$$

The third pattern is given by

$$\frac{1}{\sqrt{-g}} \partial_a (\sqrt{-g} F^{ar A_1 \dots A_{p-1}}) = 0. \quad (51)$$

This leads to

$$\ddot{\mathcal{W}} - \dot{\mathcal{V}}' + \frac{f}{r^2} (\gamma_{p-1}^{(n)} + (p-1)(n-p+1)) \mathcal{W} = 0. \quad (52)$$

The final pattern

$$\frac{1}{\sqrt{-g}} \partial_a (\sqrt{-g} F^{a A_1 \dots A_p}) = 0 \quad (53)$$

generates two equations for the $(p-1)$ -form component $\mathcal{V}^{A_1 \dots A_{p-1}}$, $\mathcal{W}^{A_1 \dots A_{p-1}}$ and p -form component $\mathcal{X}^{A_1 \dots A_p}$ as

$$\dot{\mathcal{V}} - \frac{1}{r^{n-2p}} f \partial_r (r^{n-2p} f \mathcal{W}) = 0 \quad (54)$$

and

$$-\ddot{\mathcal{X}} + \frac{1}{r^{n-2p}} f \partial_r (r^{n-2p} f \mathcal{X}') - \frac{f}{r^2} (\gamma_p^{(n)} + p(n-p)) \mathcal{X} = 0. \quad (55)$$

A. Master equations

Now, we can derive the master equations. These general results reproduce the effective potential in Eqs. (15) and (16) in [26] and in Eq. (99) in [27] as special cases and agree with the results in [28,29].

1. Coexact p -form

From Eq. (55), we can read off the master variable for the p -form component as

$$\Psi_p = \frac{1}{r^a} \mathcal{X} \quad \text{and} \quad a \equiv \frac{2p-n}{2}. \quad (56)$$

Assuming the time dependence of Ψ_p as

$$e^{-i\omega t},$$

we obtain the master equation

$$-\partial_x^2 \Psi_p + V_{p,p}^{(n)} \Psi_p = \omega^2 \Psi_p \quad (57)$$

with the effective potential $V_{p,p}^{(n)}$:

$$V_{p,p}^{(n)} = \frac{f}{r^2} \left((\ell + p)(\ell + n - p - 1) + \frac{n-2p}{2} \left(r f' + \frac{n-2p-2}{2} f \right) \right). \quad (58)$$

2. Coexact $(p-1)$ -form

The $(p-1)$ -form components are contained in Eqs. (50), (52) and (54), but the master equation is derived in Eqs. (52) and (54). Using the master variable Ψ_{p-1} for the $(p-1)$ -form

$$\Psi_{p-1} \equiv \frac{f}{r^a} \mathcal{W} \quad \text{and} \quad \mathcal{V} = r^{2a} f \partial_r \left(\frac{f}{r^{2a}} \int \mathcal{W} dt \right), \quad (59)$$

we can deduce the master equation as

$$-\partial_x^2 \Psi_{p-1} + V_{p,p-1}^{(n)} \Psi_{p-1} = \omega^2 \Psi_{p-1}. \quad (60)$$

Here, we assumed the same time dependence as before.

The effective potential $V_{p,p-1}^{(n)}$ is given by

$$V_{p,p-1}^{(n)} = \frac{f}{r^2} \left((\ell + p - 1)(\ell + n - p) + \frac{n-2p}{2} \left(\frac{n-2p+2}{2} f - r f' \right) \right). \quad (61)$$

Note that Eq. (50) is trivially satisfied.

B. Degrees of freedom and dual relations

We have shown that the D -dimensional p -form field can be represented by the coexact p -form and coexact $(p-1)$ -form fields. The condition for the coexact form $\delta \hat{\mathcal{A}}_p = 0$ can be solved as

$$\hat{\mathcal{A}}_p = \delta \hat{\mathcal{B}}_{p+1}. \quad (62)$$

However, $\hat{\mathcal{B}}_{p+1}$ has a freedom $\hat{\mathcal{B}}_{p+1} + \delta \hat{\mathcal{B}}_{p+2}$. This argument continues up to the maximum value n . Hence, the d.o.f. of the coexact p -form are given by

$${}_n C_{p+1} - {}_n C_{p+2} + {}_n C_{p+3} + \cdots = {}_n C_p - {}_n C_{p-1} + {}_n C_{p-2} + \cdots = {}_{n-1} C_p. \quad (63)$$

Similarly, we obtain ${}_{n-1} C_{p-1}$ for the d.o.f. of the coexact $(p-1)$ -form. Note that the identity

$${}_{n-1} C_p + {}_{n-1} C_{p-1} = {}_n C_p \quad (64)$$

exactly coincides with the physical d.o.f. of a p -form field.

The duality plays an important role in form fields. Indeed, we found the following duality relations:

$$V_{n-p,n-p}^{(n)} = V_{p,p-1}^{(n)}, \quad (65)$$

$$V_{n-p,n-p-1}^{(n)} = V_{p+1,p+1}^{(n)}. \quad (66)$$

In particular, in even dimensions, we have the degeneracy

$$V_{\frac{n}{2},\frac{n}{2}}^{(n)} = V_{\frac{n}{2},\frac{n}{2}-1}^{(n)}. \quad (67)$$

Later, we will see this degeneracy in the quasinormal mode spectrum.

C. Examples of effective potentials

In four dimensions where $n = 2$, the master equation for the $p = 0$ -form \mathbf{A}_0 becomes

$$V_{0,0}^{(2)} = \frac{f}{r^2} (\ell(\ell+1) + r f'). \quad (68)$$

From the master equations of the $p = 1$ -form \mathbf{A}_1 , we see that the effective potential of the coexact 1-form is

$$V_{1,1}^{(2)} = \ell(\ell+1) \frac{f}{r^2} \quad (69)$$

and that of the coexact 0-form reads

$$V_{1,0}^{(2)} = \ell(\ell + 1) \frac{f}{r^2}. \quad (70)$$

The effective potentials for the coexact 1-form and the coexact 0-form components in A_1 are the same. Our results agree with the expression (99) in [27]. Moreover, since the coexact 2-forms on the sphere S^2 do not exist, the master equation for the $p = 2$ -form A_2 becomes single. The effective potential of the coexact 1-form component is given by

$$V_{2,1}^{(2)} = \frac{f}{r^2} (\ell(\ell + 1) + rf'). \quad (71)$$

As is expected, this effective potential is the same expression as that for the 0-form field in Eq. (68). This confirms that we can consider only the form fields with a rank larger than p_M .

In five dimensions where $n = 3$, the master equation for the $p = 0$ -form A_0 becomes

$$V_{0,0}^{(3)} = \frac{f}{r^2} \left(\ell(\ell + 2) + \frac{3}{2} \left(rf' + \frac{f}{2} \right) \right). \quad (72)$$

We can also read off the effective potentials in the master equations for the 1-form A_1 . The effective potential of the coexact 1-form component reads

$$V_{1,1}^{(3)} = \frac{f}{r^2} \left((\ell + 1)^2 + \frac{1}{2} \left(rf' - \frac{1}{2}f \right) \right) \quad (73)$$

and that of the coexact 0-form component is given by

$$V_{1,0}^{(3)} = \frac{f}{r^2} \left(\ell(\ell + 2) + \frac{1}{2} \left(\frac{3}{2}f - rf' \right) \right). \quad (74)$$

In five dimensions, we need not consider a 2-form field because of the duality. Here, for the 2-form A_2 . The effective potential of the coexact 2-form component is given by

$$V_{2,2}^{(3)} = \frac{f}{r^2} \left(\ell(\ell + 2) + \frac{1}{2} \left(\frac{3}{2}f - rf' \right) \right) \quad (75)$$

and that of the coexact 1-form component becomes

$$V_{2,1}^{(3)} = \frac{f}{r^2} \left((\ell + 1)^2 + \frac{1}{2} \left(rf' - \frac{1}{2}f \right) \right). \quad (76)$$

The effective potential (76) is the same as Eq. (73), and the effective potential (75) is the same as Eq. (74). These results just reflect the duality relations (65) and (66).

It is also easy to explicitly write down the master equations for the p -form fields in higher dimensions.

V. STABILITY ANALYSIS

In this section, we show the stability of the p -form field in arbitrary dimensions. The stability of p -form fields in black hole spacetime is nontrivial because the effective potential has a negative region as is shown in Fig. 1. To show the stability of the fields around the black holes, the S -deformation method [7–10,30] is useful. Hence, first, we shortly review the S -deformation method. Secondly, we show the stability of the effective potential for each component of the p -form field.

Let us start with the master equation

$$\omega^2 \Psi = A \Psi, \quad (77)$$

where we defined the operator A as

$$A \equiv -\frac{d^2}{dx^2} \Psi + V \Psi, \quad (78)$$

and we assume the time dependence of Ψ as

$$\Psi \propto e^{-i\omega t}. \quad (79)$$

If $\omega^2 < 0$ for the boundary conditions $\Psi \rightarrow 0$ and $d\Psi/dx \rightarrow 0$ at $x \rightarrow \pm\infty$, this solution is unstable and exponentially grows. So, if we want to show the stability, we have to show $\omega^2 > 0$. The S -deformation is defined by using the new operator D_x as follows:




FIG. 1. The effective potentials for the 1-form in 10 dimensions are plotted. The potential for the coexact 0-form component has a negative region. Here, we took $\ell = 1$ and $\mu = 1$.

$$D_x \equiv \frac{d}{dx} + S(x). \quad (80)$$

Then, Eq. (77) is modified as

$$\begin{aligned} \omega^2 |\Psi|^2 &= \Psi^* A \Psi \\ &= \Psi^* \left[-\frac{d^2}{dx^2} + V \right] \Psi \\ &= -\frac{d}{dx} (\Psi^* D_x \Psi) + |D_x \Psi|^2 + \bar{V} |\Psi|^2, \\ \bar{V} &\equiv V + \frac{dS}{dx} - S^2. \end{aligned} \quad (81)$$

Suppose we find the continuous function, S , which makes $\bar{V} > 0$ for $-\infty < x < \infty$. Then, after integration, we have

$$\begin{aligned} \omega^2 \int_{-\infty}^{\infty} dx |\Psi|^2 &= -[\Psi^* D_x \Psi]_{-\infty}^{\infty} \\ &+ \int_{-\infty}^{\infty} dx [|D_x \Psi|^2 + \bar{V} |\Psi|^2] > 0. \end{aligned} \quad (82)$$

Here, we assumed the boundary conditions $\Psi \rightarrow 0$ and $d\Psi/dx \rightarrow 0$ at $x \rightarrow \pm\infty$; that is, the Ψ and $d\Psi/dx$ have a compact support, so the first term in Eq. (82) vanishes. The above inequality implies the positivity of ω^2 ; that is, there is no growing mode for Ψ .

In the present case, an appropriate S -deformation exists as follows:

$$S = -\frac{d}{dx} \left(\ln \left(\frac{1}{r^\alpha} \right) \right) \quad (83)$$

which is inspired by [17].

A. Effective potential of coexact p -form

If we choose

$$\alpha = -\frac{n-2p}{2}, \quad (84)$$

the effective potential of the coexact p -form on the sphere becomes

$$\begin{aligned} V_{p,p}^{(n)} &\rightarrow \bar{V}_{p,p}^{(n)} = V_{p,p}^{(n)} + f \frac{dS}{dr} - S^2 \\ &= \frac{f}{r^2} (\ell + p)(\ell + n - p - 1). \end{aligned} \quad (85)$$

This modified potential is positive definite outside the event horizon, because p satisfies $p \leq p_{\max} < n$.

B. Effective potential of coexact $(p-1)$ -form

If we choose

$$\alpha = \frac{n-2p}{2}, \quad (86)$$

the effective potential of the coexact $(p-1)$ -form on the sphere becomes

$$\begin{aligned} V_{p,p-1}^{(n)} &\rightarrow \bar{V}_{p,p-1}^{(n)} = V_{p,p-1}^{(n)} + f \frac{dS}{dr} - S^2 \\ &= \frac{f}{r^2} ((\ell + p)(\ell + n - p - 1) - (n - 2p)) \\ &\geq \frac{f}{r^2} ((1 + p)(1 + n - p - 1) - (n - 2p)) \\ &= \frac{f}{r^2} p(n - p + 1) > 0. \end{aligned} \quad (87)$$

This modified potential is also positive definite outside the event horizon.

From the above analysis, we see that the p -form fields in arbitrary dimensions are stable in the spherical black hole even for other gravity theories as long as f satisfies the positivity $f > 0$ outside the horizon $r > r_h$.

VI. QUASINORMAL MODES

We got the master equations and the effective potentials for each component of the p -form field. In this section we present the quasinormal modes for the general form fields. The quasinormal modes are fundamental vibration modes around a black hole, and these modes are obtained by solving the master equation under the appropriate boundary conditions. The general formalism for calculating quasinormal modes by using WKB approximation has been proposed by Schutz and Will in [20] and subsequently developed by many people in [21,22,31–35]. Here, we summarize the main points of the WKB method for calculating quasinormal modes.

Firstly, we can divide the region into two regions. The region I ($-\infty < x < x_0$) is the one ranging from the top of the effective potential x_0 to the horizon of the black hole. The region II ($x_0 < x < \infty$) is the one ranging from the top of the potential x_0 to the far outside of the black hole, i.e., infinity. The wave traveling to the potential is called the ingoing wave and the wave traveling from the potential is called the outgoing wave. In each region, the solutions of the master equation can be expressed as

$$\begin{cases} \Psi_I = Z_I^{\text{in}} \Psi_I^{\text{in}} + Z_I^{\text{out}} \Psi_I^{\text{out}}, \\ \Psi_{II} = Z_{II}^{\text{in}} \Psi_{II}^{\text{in}} + Z_{II}^{\text{out}} \Psi_{II}^{\text{out}}, \end{cases} \quad (88)$$

where Ψ^{in} and Ψ^{out} represent the ingoing and outgoing waves, respectively. The boundary condition for obtaining quasinormal modes is that there are no ingoing waves:

$$Z_I^{\text{in}} = Z_{\text{II}}^{\text{in}} = 0. \quad (89)$$

Since there are two conditions, only discrete complex eigenvalues are allowed.

In the N th order WKB method, we approximate the function $Q(x)$ defined by

$$Q(x) \equiv \omega^2 - V(x) \quad (90)$$

in terms of a $2N$ th order Taylor expansion around the maximum of the potential as follows:

$$\left. \frac{dQ(x)}{dx} \right|_{x=x_0} = Q_0^{(1)} = 0 \quad \text{and} \quad Q(x) \simeq \sum_{k=0}^{2N} \frac{1}{k!} Q_0^{(k)} (x - x_0)^k. \quad (91)$$

Expressing the wave function using WKB approximation, we can calculate the scattering matrix. Thus, we can get the formula for quasinormal modes as

$$\omega \simeq \sqrt{V_0 + \sqrt{2V_0^{(2)}} \left(n_{\text{tone}} + \frac{1}{2} + \sum_{k=1}^{N-1} \Omega_k \right)}, \quad (92)$$

where $V_0 \equiv V(x_0)$, and $V_0^{(k)}$ is the k th order derivative of the potential

$$V_0^{(k)} \equiv \left. \frac{d^k V(x)}{dx^k} \right|_{x=x_0}, \quad (93)$$

and Ω_1 and Ω_2 are given by

$$\Omega_1 = (-30\beta_1^2 + 6\beta_2)\beta^2 - \frac{7}{2}\beta_1^2 + \frac{3}{2}\beta_2 \quad (94)$$

$$\begin{aligned} \Omega_2 = & (-2820\beta_1^4 + 1800\beta_1^2\beta_2 - 280\beta_1\beta_3 - 68\beta_2^2 + 20\beta_4)\beta^3 \\ & + (-1155\beta_1^4 + 918\beta_1^2\beta_2 - 190\beta_1\beta_3 - 67\beta_2^2 + 25\beta_4)\beta. \end{aligned} \quad (95)$$

Here, β is defined by

$$\beta \equiv n_{\text{tone}} + \frac{1}{2} \quad (96)$$

and $\beta_k (k \geq 1)$ is defined by

$$\beta_k \equiv \frac{V_0^{(k+2)}}{(k+2)!} \left(\frac{1}{2V_0^{(2)}} \right)^{\frac{k+1}{2}}. \quad (97)$$

The higher Ω_k can be derived explicitly, but the expressions of $\Omega_k (k \geq 3)$ are too long to write down them here. So, we give them in the Appendix. The parameter, n_{tone} , is called the tone number of quasinormal modes. This method is

TABLE I. The QNMs of 2-form field A_2 .

D	2-form component
6	$1.2618 - 0.4616i$
7	$1.7509 - 0.5920i$
8	$2.2231 - 0.7192i$
9	$2.6681 - 0.8555i$
10	$3.0791 - 1.0080i$

D	1-form component
6	$1.2618 - 0.4616i$
7	$1.5387 - 0.5652i$
8	$1.8352 - 0.7345i$
9	$2.2630 - 0.8521i$
10	$2.6910 - 0.9444i$

often called the N th order WKB approximation. It is known that in the case of $n_{\text{tone}} < \ell$ this approximation is good. So we focus on the case $n_{\text{tone}} = 0$ in this paper.

Now, we show the quasinormal modes of the p -form fields up to $D = 10$ dimensions. In this case, we need to consider form fields up to $p = 4$. In this study we used the sixth order WKB method, so we need the $\Omega_1, \Omega_2, \dots, \Omega_5$. We choose a set of parameters,

$$(\ell, \mu, n_{\text{tone}}) = (1, 1, 0). \quad (98)$$

In Table I, we showed the quasinormal modes (QNMs) of a 2-form field. As you can see the QNM of the coexact 2-form component and the QNM of the coexact 1-form component in six dimensions coincide. This comes from the duality relations (65) and (66). Except for $D = 8$, the coexact 2-form component decays faster than the coexact 1-form component. In Table II, we listed the QNMs of a 3-form field. In $D = 8$ dimensions, we can see that duality relations hold. In other dimensions, the coexact 2-form component decays faster than the coexact 3-form component. In Table III, we displayed the QNMs of a 4-form field. In this case, only $D = 10$ is relevant. Here, we see the duality relations again. In all cases, we see, as D increases,

TABLE II. The QNMs of 3-form field A_3 .

D	3-form component
8	$2.0779 - 0.6754i$
9	$2.6018 - 0.7640i$
10	$3.1539 - 0.8307i$

D	2-form component
8	$2.0779 - 0.6754i$
9	$2.3795 - 0.7729i$
10	$2.6947 - 0.9197i$

TABLE III. The QNMs of 4-form field A_4 .

D	4-form component
10	$2.9227 - 0.8595i$
D	3-form component
10	$2.9227 - 0.8595i$

the real and imaginary parts of the quasinormal frequency increase.

VII. CONCLUSION

We studied the quasinormal modes of p -form fields in spherical black holes in arbitrary dimensions. Using the spherical symmetry of the black holes and gauge symmetry, we showed that the p -form field can be expressed in terms of the coexact p -form and coexact $(p-1)$ -form on the sphere. These variables allow us to find the master equations. We revealed some relations between the effective potentials in the master equations. We found that the effective potential can have a negative region for some parameters. Therefore, by utilizing the S -deformation method, we explicitly showed the stability of p -form fields in the spherical black hole spacetime. Finally, using the WKB approximation, we calculated the quasinormal modes of p -form fields in $D(\leq 10)$ dimensions. There,

we can see the degeneracy of the spectrum expected from the duality relations we found.

It is interesting to include rotations of black holes in our analysis. Recently, Lunin found the ansatz for the 1-form field in arbitrary dimensions to get the separable equations of motion [36]. It is interesting to investigate p -form fields in higher dimensional rotational black holes. We can also consider higher spin fields in arbitrary dimensions. Resolving the above problems must have implications for string theory. We leave these problems for future work.

ACKNOWLEDGMENTS

We would like to thank A. Ito for useful discussion. D. Y. was supported by Grant-in-Aid for JSPS Research Fellow and JSPS KAKENHI Grant No. 17J00490. J. S. was in part supported by JSPS KAKENHI Grants No. JP17H02894, No. JP17K18778, No. JP15H05895, No. JP17H06359, and No. JP18H04589. J. S. is also supported by JSPS Bilateral Joint Research Projects (JSPS-NRF collaboration) ‘‘String Axion Cosmology.’’

APPENDIX

To perform the sixth order WKB method, we need Ω_3 , Ω_4 and Ω_5 in addition to Ω_1 , Ω_2 . Here, we display them for completeness:

$$\begin{aligned}
\Omega_3 = & (-463020\beta_1^6 + 465300\beta_1^4\beta_2 - 78120\beta_1^3\beta_3 - 99780\beta_1^2\beta_2^2 + 10860\beta_1^2\beta_4 + 19320\beta_1\beta_2\beta_3 \\
& - 1260\beta_1\beta_5 + 1500\beta_2^3 - 660\beta_2\beta_4 - 630\beta_3^2 + 70\beta_6)\beta^4 \\
& + (-418110\beta_1^6 + 479970\beta_1^4\beta_2 - 95460\beta_1^3\beta_3 - 124026\beta_1^2\beta_2^2 + 17070\beta_1^2\beta_4 + 29340\beta_1\beta_2\beta_3 \\
& + 3414\beta_2^3 - 2730\beta_1\beta_5 - 1770\beta_2\beta_4 - 1085\beta_3^2 + 245\beta_6)\beta^2 \\
& - \frac{101479}{4}\beta_1^6 + \frac{131817}{4}\beta_1^4\beta_2 - \frac{14777}{2}\beta_1^3\beta_3 - \frac{40261}{4}\beta_1^2\beta_2^2 + \frac{6055}{4}\beta_1^2\beta_4 + \frac{5667}{2}\beta_1\beta_2\beta_3 \\
& - \frac{1155}{4}\beta_1\beta_5 + \frac{1539}{4}\beta_2^3 - \frac{945}{4}\beta_2\beta_4 - \frac{1107}{8}\beta_3^2 + \frac{315}{8}\beta_6,
\end{aligned} \tag{A1}$$

$$\begin{aligned}
\Omega_4 = & (-95872644\beta_1^8 + 130619664\beta_1^6\beta_2 - 22467312\beta_1^5\beta_3 - 51067800\beta_1^4\beta_2^2 + 3454920\beta_1^4\beta_4 \\
& + 13073760\beta_1^3\beta_2\beta_3 + 5418000\beta_1^2\beta_2^3 - 493920\beta_1^3\beta_5 - 1285200\beta_1^2\beta_2\beta_4 - 732480\beta_1^2\beta_3^2 \\
& - 1140720\beta_1\beta_2^2\beta_3 - 42756\beta_2^4 + 59472\beta_1^2\beta_6 + 98784\beta_1\beta_2\beta_5 + 110544\beta_1\beta_3\beta_4 + 25032\beta_2^2\beta_4 \\
& + 49392\beta_2\beta_3^2 - 5544\beta_1\beta_7 - 3024\beta_2\beta_6 - 5544\beta_3\beta_5 - 1572\beta_4^2 + 252\beta_8)\beta^5 \\
& + (-154601370\beta_1^8 + 231728040\beta_1^6\beta_2 - 45019560\beta_1^5\beta_3 - 101714460\beta_1^4\beta_2^2 + 8269260\beta_1^4\beta_4 \\
& + 29638800\beta_1^3\beta_2\beta_3 + 12782760\beta_1^2\beta_2^3 - 1456560\beta_1^3\beta_5 - 3618360\beta_1^2\beta_2\beta_4 - 1870400\beta_1^2\beta_3^2 \\
& - 3138600\beta_1\beta_2^2\beta_3 - 178330\beta_2^4 + 223720\beta_1^2\beta_6 + 361200\beta_1\beta_2\beta_5 + 354120\beta_1\beta_3\beta_4 \\
& + 118220\beta_2^2\beta_4 + 150360\beta_2\beta_3^2 - 28140\beta_1\beta_7 - 17640\beta_2\beta_6 - 21420\beta_3\beta_5 - 8290\beta_4^2 + 1890\beta_8)\beta^3 \\
& + \left(-\frac{129443349}{4}\beta_1^8 + 53574549\beta_1^6\beta_2 - 11535783\beta_1^5\beta_3 - \frac{53000175}{2}\beta_1^4\beta_2^2 + \frac{4785249}{2}\beta_1^4\beta_4 \right.
\end{aligned}$$

$$\begin{aligned}
& + 8708550\beta_1^3\beta_2\beta_3 + 3909285\beta_1^2\beta_2^3 - 482622\beta_1^3\beta_5 - 1246797\beta_1^2\beta_2\beta_4 - 632340\beta_1^2\beta_3^2 \\
& - 1119415\beta_1\beta_2^2\beta_3 - \frac{305141}{4}\beta_2^4 + 88753\beta_1^2\beta_6 + 149478\beta_1\beta_2\beta_5 + 145417\beta_1\beta_3\beta_4 + \frac{117281}{2}\beta_2^2\beta_4 \\
& + 64731\beta_2\beta_3^2 - \frac{28077}{2}\beta_1\beta_7 - 10521\beta_2\beta_6 - \frac{22029}{2}\beta_3\beta_5 - \frac{19277}{4}\beta_4^2 + \frac{5607}{4}\beta_8 \Big) \beta \quad \text{and} \quad (A2)
\end{aligned}$$

$$\begin{aligned}
\Omega_5 = & (-22598568720\beta_1^{10} + 38797354512\beta_1^8\beta_2 - 6749494080\beta_1^7\beta_3 - 22002812832\beta_1^6\beta_2^2 \\
& + 1080123744\beta_1^6\beta_4 + 6257684160\beta_1^5\beta_2\beta_3 + 4660027680\beta_1^4\beta_2^3 - 165561984\beta_1^5\beta_5 \\
& - 766795680\beta_1^4\beta_2\beta_4 - 413669760\beta_1^4\beta_3^2 - 1443234240\beta_1^3\beta_2^2\beta_3 - 291804240\beta_1^2\beta_2^4 \\
& + 23163840\beta_1^4\beta_6 + 85128960\beta_1^3\beta_2\beta_5 + 90350400\beta_1^3\beta_3\beta_4 + 108679200\beta_1^2\beta_2^2\beta_4 \\
& + 126329280\beta_1^2\beta_2\beta_3^2 + 64196160\beta_1\beta_2^3\beta_3 + 1400784\beta_2^5 - 2919840\beta_1^3\beta_7 - 7531776\beta_1^2\beta_2\beta_6 \\
& - 8618400\beta_1^2\beta_3\beta_5 - 3939600\beta_1^2\beta_4^2 - 6233472\beta_1\beta_2^2\beta_5 - 13981632\beta_1\beta_2\beta_3\beta_4 - 2849280\beta_1\beta_3^3 \\
& - 1023456\beta_2^3\beta_4 - 3116736\beta_2^2\beta_3^2 + 307440\beta_1^2\beta_8 + 487872\beta_1\beta_2\beta_7 + 583520\beta_1\beta_3\beta_6 \\
& + 544320\beta_1\beta_4\beta_5 + 129472\beta_2^2\beta_6 + 487872\beta_2\beta_3\beta_5 + 134736\beta_2\beta_4^2 + 272160\beta_3^2\beta_4 \\
& - 24024\beta_1\beta_9 - 13440\beta_2\beta_8 - 24024\beta_3\beta_7 - 14224\beta_4\beta_6 - 12012\beta_5^2 + 924\beta_{10})\beta^6 \\
& + (-57626387280\beta_1^{10} + 106553134800\beta_1^8\beta_2 - 20386144800\beta_1^7\beta_3 - 65772661920\beta_1^6\beta_2^2 \\
& + 3750215280\beta_1^6\beta_4 + 20631693600\beta_1^5\beta_2\beta_3 + 15479738400\beta_1^4\beta_2^3 - 675647280\beta_1^5\beta_5 \\
& - 2947719600\beta_1^4\beta_2\beta_4 - 1496632200\beta_1^4\beta_3^2 - 5324508000\beta_1^3\beta_2^2\beta_3 - 1135963920\beta_1^2\beta_2^4 \\
& + 113720040\beta_1^4\beta_6 + 390240480\beta_1^3\beta_2\beta_5 + 381413280\beta_1^3\beta_3\beta_4 + 487029840\beta_1^2\beta_2^2\beta_4 \\
& + 513631440\beta_1^2\beta_2\beta_3^2 + 283029600\beta_1\beta_2^3\beta_3 + 9396240\beta_2^5 - 17603880\beta_1^3\beta_7 - 42912240\beta_1^2\beta_2\beta_6 \\
& - 43022280\beta_1^2\beta_3\beta_5 - 19968240\beta_1^2\beta_4^2 - 34807920\beta_1\beta_2^2\beta_5 - 69789120\beta_1\beta_2\beta_3\beta_4 \\
& - 12597200\beta_1\beta_3^3 - 7583760\beta_2^3\beta_4 - 14953960\beta_2^2\beta_3^2 + 2324700\beta_1^2\beta_8 + 3618720\beta_1\beta_2\beta_7 \\
& + 3614520\beta_1\beta_3\beta_6 + 3353280\beta_1\beta_4\beta_5 + 1142120\beta_2^2\beta_6 + 2859360\beta_2\beta_3\beta_5 + 1092720\beta_2\beta_4^2 \\
& + 1464400\beta_3^2\beta_4 - 237930\beta_1\beta_9 - 147000\beta_2\beta_8 - 182490\beta_3\beta_7 - 135380\beta_4\beta_6 \\
& + -82005\beta_5^2 + 12705\beta_{10})\beta^4 \\
& + \left(-26541790065\beta_1^{10} + 53237904993\beta_1^8\beta_2 - 11123381220\beta_1^7\beta_3 - 36045764154\beta_1^6\beta_2^2 \right. \\
& + 2279955006\beta_1^6\beta_4 + 12440307420\beta_1^5\beta_2\beta_3 + 9481289682\beta_1^4\beta_2^3 - 461383776\beta_1^5\beta_5 \\
& - 2012614434\beta_1^4\beta_2\beta_4 - 999867660\beta_1^4\beta_3^2 - 3630132780\beta_1^3\beta_2^2\beta_3 - 809619141\beta_1^2\beta_2^4 \\
& + 89013120\beta_1^4\beta_6 + 304548384\beta_1^3\beta_2\beta_5 + 292426020\beta_1^3\beta_3\beta_4 + 388974714\beta_1^2\beta_2^2\beta_4 \\
& + 392853060\beta_1^2\beta_2\beta_3^2 + 230221620\beta_1\beta_2^3\beta_3 + 9317949\beta_2^5 - 16119726\beta_1^3\beta_7 \\
& - 39660264\beta_1^2\beta_2\beta_6 - 38201310\beta_1^2\beta_3\beta_5 - 18168321\beta_1^2\beta_4^2 - 33071472\beta_1\beta_2^2\beta_5 \\
& - 63944892\beta_1\beta_2\beta_3\beta_4 - 10841880\beta_1\beta_3^3 - 8518614\beta_2^3\beta_4 - 14034096\beta_2^2\beta_3^2 + 2582685\beta_1^2\beta_8 \\
& + 4096764\beta_1\beta_2\beta_7 + 3870930\beta_1\beta_3\beta_6 + 3671892\beta_1\beta_4\beta_5 + 1518048\beta_2^2\beta_6 + 3168732\beta_2\beta_3\beta_5 \\
& + 1390869\beta_2\beta_4^2 + 1530210\beta_3^2\beta_4 - \frac{671517}{2}\beta_1\beta_9 - 236460\beta_2\beta_8 - \frac{476973}{2}\beta_3\beta_7 \\
& \left. + -204771\beta_4\beta_6 - \frac{444381}{4}\beta_5^2 + \frac{101409}{4}\beta_{10} \right) \beta^2 \\
& - \frac{2375536317}{2}\beta_1^{10} + \frac{5112354429}{2}\beta_1^8\beta_2 - 570170440\beta_1^7\beta_3 - 1875235809\beta_1^6\beta_2^2
\end{aligned}$$

$$\begin{aligned}
& + 125451228\beta_1^6\beta_4 + 697320300\beta_1^5\beta_2\beta_3 + 542138237\beta_1^4\beta_2^3 - 27429003\beta_1^5\beta_5 \\
& - 122723430\beta_1^4\beta_2\beta_4 - \frac{121918445}{2}\beta_1^4\beta_3^2 - 226646440\beta_1^3\beta_2^2\beta_3 - \frac{104283313}{2}\beta_1^2\beta_2^4 \\
& + \frac{11623829}{2}\beta_1^4\beta_6 + 20366894\beta_1^3\beta_2\beta_5 + 19607424\beta_1^3\beta_3\beta_4 + 27070372\beta_1^2\beta_2^2\beta_4 \\
& + 27194427\beta_1^2\beta_2\beta_3^2 + 16533060\beta_1\beta_2^3\beta_3 + \frac{1456569}{2}\beta_2^5 - \frac{2336663}{2}\beta_1^3\beta_7 - 2995587\beta_1^2\beta_2\beta_6 \\
& - \frac{5703723}{2}\beta_1^2\beta_3\beta_5 - \frac{2729425\beta_1^2\beta_4^2}{2} - 2594391\beta_1\beta_2^2\beta_5 - 5045766\beta_1\beta_2\beta_3\beta_4 - 854685\beta_1\beta_3^3 \\
& - 735210\beta_2^3\beta_4 - \frac{2301381}{2}\beta_2^2\beta_3^2 + \frac{848925}{4}\beta_1^2\beta_8 + 358344\beta_1\beta_2\beta_7 + \frac{674037}{2}\beta_1\beta_3\beta_6 \\
& + 315150\beta_1\beta_4\beta_5 + \frac{292005}{2}\beta_2^2\beta_6 + 289908\beta_2\beta_3\beta_5 + \frac{269325}{2}\beta_2\beta_4^2 + 143370\beta_3^2\beta_4 \\
& - \frac{259875}{8}\beta_1\beta_9 - \frac{51975}{2}\beta_2\beta_8 - \frac{203931}{8}\beta_3\beta_7 - \frac{89775}{4}\beta_4\beta_6 - \frac{180675}{16}\beta_5^2 + \frac{51975}{16}\beta_{10}. \tag{A3}
\end{aligned}$$

-
- [1] T. Regge and J. A. Wheeler, *Phys. Rev.* **108**, 1063 (1957).
[2] F. J. Zerilli, *Phys. Rev. Lett.* **24**, 737 (1970).
[3] V. Moncrief, *Ann. Phys. (N.Y.)* **88**, 323 (1974).
[4] B. S. Kay and R. M. Wald, *Classical Quantum Gravity* **4**, 893 (1987).
[5] M. Dafermos, G. Holzegel, and I. Rodnianski, *arXiv:1601.06467*.
[6] S. B. Giddings and S. D. Thomas, *Phys. Rev. D* **65**, 056010 (2002).
[7] H. Kodama and A. Ishibashi, *Prog. Theor. Phys.* **110**, 701 (2003).
[8] A. Ishibashi and H. Kodama, *Prog. Theor. Phys.* **110**, 901 (2003).
[9] H. Kodama and A. Ishibashi, *Prog. Theor. Phys.* **111**, 29 (2004).
[10] H. Kodama, *Lect. Notes Phys.* **769**, 427 (2009).
[11] D. Lovelock, *J. Math. Phys.* **12**, 498 (1971).
[12] C. Charmousis, *Lect. Notes Phys.* **769**, 299 (2009).
[13] T. Takahashi and J. Soda, *Phys. Rev. D* **79**, 104025 (2009).
[14] T. Takahashi and J. Soda, *Phys. Rev. D* **80**, 104021 (2009).
[15] T. Takahashi and J. Soda, *Prog. Theor. Phys.* **124**, 711 (2010).
[16] T. Takahashi and J. Soda, *Prog. Theor. Phys.* **124**, 911 (2010).
[17] T. Takahashi, *Stability Analysis of Black Hole Solutions in Lovelock Theory*, Ph.D. thesis, Kyoto University, 2013.
[18] R. A. Konoplya and A. Zhidenko, *J. Cosmol. Astropart. Phys.* **05** (2017) 050.
[19] R. M. Wald, *J. Math. Phys.* **20**, 1056 (1979).
[20] B. F. Schutz and C. M. Will, *Astrophys. J.* **291**, L33 (1985).
[21] S. Iyer and C. M. Will, *Phys. Rev. D* **35**, 3621 (1987).
[22] S. Iyer, *Phys. Rev. D* **35**, 3632 (1987).
[23] Y. Choquet-Bruhat, C. DeWitt-Morette, C. de Witt, M. Dillard-Bleick, and M. Dillard-Bleick, *Analysis, Manifolds and Physics Revised Edition* (Gulf Professional Publishing, North-Holland, 1982).
[24] F. R. Tangherlini, *Nuovo Cimento* **27**, 636 (1963).
[25] R. Camporesi and A. Higuchi, *J. Geom. Phys.* **15**, 57 (1994).
[26] V. Cardoso, J. P. S. Lemos, and S. Yoshida, *Phys. Rev. D* **69**, 044004 (2004).
[27] E. Chaverra, N. Ortiz, and O. Sarbach, *Phys. Rev. D* **87**, 044015 (2013).
[28] D.-P. Du, B. Wang, and R.-K. Su, *Phys. Rev. D* **70**, 064024 (2004).
[29] A. Lopez-Ortega, *Gen. Relativ. Gravit.* **38**, 1565 (2006).
[30] M. Kimura, *Classical Quantum Gravity* **34**, 235007 (2017).
[31] K. D. Kokkotas and B. F. Schutz, *Phys. Rev. D* **37**, 3378 (1988).
[32] E. Seidel and S. Iyer, *Phys. Rev. D* **41**, 374 (1990).
[33] R. A. Konoplya, *Phys. Rev. D* **68**, 024018 (2003).
[34] R. A. Konoplya and A. Zhidenko, *Rev. Mod. Phys.* **83**, 793 (2011).
[35] D. Yoshida and J. Soda, *Phys. Rev. D* **93**, 044024 (2016).
[36] O. Lunin, *J. High Energy Phys.* **12** (2017) 138.