



Central administration of insulin-like growth factor-2 suppresses food intake in chicks

Honda, Kazuhisa
Kewan, Ahmed
Osada, Haruki
Saneyasu, Takaoki
Kamisoyama, Hiroshi

(Citation)

Neuroscience Letters, 751:135797

(Issue Date)

2021-04-23

(Resource Type)

journal article

(Version)

Accepted Manuscript

(Rights)

© 2021 Elsevier B.V.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

(URL)

<https://hdl.handle.net/20.500.14094/90008183>



1 Central administration of insulin-like growth factor-2 suppresses food intake in chicks

2

3 Kazuhisa Honda, Ahmed Kewan, Haruki Osada, Takaoki Saneyasu, and Hiroshi Kamisoyama

4

5 Department of Bioresource Science, Graduate School of Agricultural Science, Kobe University,

6 Kobe 657-8501, Japan

7

8 Running title: IGF-2 suppresses food intake in chicks

9

10 **Corresponding author:**

11 Kazuhisa Honda, Ph.D.

12 Department of Bioresource Science, Graduate School of Agricultural Science, Kobe University,

13 Kobe 657-8501, Japan

14 Tel./fax: +81 78 803 5809

15 E-mail address: honda@tiger.kobe-u.ac.jp

16

17

18 **Abstract**

19 Insulin-like growth factor (IGF)-2 is a multifunctional hormone with structural and functional
20 similarity to IGF-1 in mammals and chickens. We previously showed that intracerebroventricular
21 administration of IGF-1 suppresses food intake in chicks. Also, central administration of IGF-2
22 suppresses food intake in rats. In the present study, we evaluated whether IGF-2 is involved in the
23 regulation of food intake in chicks. We also examined the effects of fasting on the mRNA levels
24 of IGF binding proteins (IGFBPs) in the liver and hypothalamus, because IGFBPs bind IGF-1 and
25 -2 in plasma and block their binding to the receptors, and locally expressed IGFBPs also influence
26 IGFs binding to the receptors in mammals. Intracerebroventricular administration of IGF-2
27 significantly suppressed food intake in chicks. The mRNA levels of IGFBPs in the hypothalamus
28 were not affected by six hours of fasting. On the other hand, six hours of fasting markedly
29 increased the mRNA levels of hepatic IGFBP-1 and -2 (5.47- and 6.95-fold, respectively). The
30 mRNA levels of IGFBP-3 were also significantly increased (1.36-fold) by six hours of fasting,
31 whereas the mRNA levels of IGF-2, IGFBP-4, and -5 were unchanged. These findings suggest
32 that circulating IGF-2 may be involved in satiety signals, but its physiological role may be
33 regulated by IGFBPs production in the liver in chicks.

34

35 **Keywords:** appetite, brain, chicken, feed intake, insulin-like growth factor-2, liver

36

1. Introduction

Insulin-like growth factor (IGF)-1 plays important roles in the muscle development and growth of chickens [6, 7]. We recently found that central administration of IGF-1 suppresses food intake in chicks [12, 13]. The mRNA levels of proopiomelanocortin, an anorexigenic in the hypothalamus, are increased by central administration of IGF-1 [13]. There is evidence that the plasma concentration of insulin and IGF-1 were postprandially elevated in chickens [20]. It is therefore likely that IGF-1 functions as a postprandial satiety hormones in chickens.

IGF-2 is a multifunctional hormone with structural and functional similarity to IGF-1 in mammals and chickens [29]. Central administration of 100 ng (~13.3 pmol) of IGF-2 suppressed food intake in rats [21]. Chicken IGF-2 can bind chicken IGF-1 receptor (IGF-1R) and express the function [8]. In addition, the chicken mannose 6-phosphate receptor (MPR300), which is known to be an IGF-2 receptor (IGF-2R) in mammals, can bind chicken IGF-2 *in vitro* [18] and was expressed in brains of chickens [27]. Feed restriction reduced the plasma IGF-2 concentration in chickens [22]. The mRNA levels of IGF-2 in the liver were increased by a high energy diet in growing chicks [35]. These findings raise the hypothesis that IGF-2 may function as a satiety hormone in chickens.

IGF binding proteins (IGFBPs) bind IGF-1 and -2 in plasma and block their binding to the receptors, indicating that IGFBPs influence the physiological roles of IGFs. For example, addition of either IGFBP-1 or IGFBP-2 to sera reduced free IGF-1 *in vitro* [11]. An increase in IGFBP-1 and reduction in free IGF-1 are accompanied by an increase in IGFBP-1 complexed IGF-1 after fasting in humans [10]. In chickens, only 5% of serum IGF-2 exists in free form, suggesting that the function of almost all plasma IGF-2 is suppressed by IGFBPs [29]. Interestingly, recent evidence demonstrated that locally expressed IGFBPs increased IGFs availability for binding to the receptors, and that IGF exhibited an independent action in mammals and fishes [1, 36].

In the present study, we investigated the possible involvement of IGF-2 in the

63 mechanism of food intake regulation in chicks. We also examined the effects of fasting on the
64 mRNA levels of IGFBPs in the liver and hypothalamus, the production area of IGFBPs and the
65 target site of IGF-2, respectively. Our findings suggest that hepatic IGFBP-1, -2, and -3
66 production may suppress the anorexigenic function of IGF-2 in chicks under *ad libitum* feeding
67 conditions.

68

69 **2. Materials and methods**

70 *2.1. Animals and peptides*

71 This study was approved by the Institutional Animal Care and Use Committee and
72 carried out according to the Kobe University Animal Experimentation Regulations (25-08-01 and
73 27-07-01). Day-old male chicks (White leghorn) were purchased from a local hatchery (Japan
74 Layer K.K., Gifu, Japan). They were given free access to water and a commercial chick starter
75 diet (Nippon Formula Feed Mfg. Co., Ltd., Kanagawa, Japan) in an electrically heated cages
76 (1725 mm x 425 mm x 320 mm) maintained at $28 \pm 2^{\circ}\text{C}$ in a room with an automatically
77 controlled 23 h light/1h dark cycle (23:00-24:00 dark). Room temperature was $22 \pm 2^{\circ}\text{C}$.

78 Amino acid sequencing of IGF-2 showed the presence of 12 amino acid substitutions
79 compared with humans [29]. However, Upton et al. [39] reported that recombinant chicken IGF-2
80 was equipotent with human IGF-2 in both biological and receptor binding studies in chick embryo
81 fibroblasts. Purified chicken MPR 300 binds both chicken and human IGF-2 [18]. The metabolic
82 clearance of chicken IGF-2 and human IGF-2 was similar when administered intravascularly in
83 7-week-old chickens [30]. These findings suggest that human IGF-2 and chicken IGF-2 show
84 similar effects in chicks. Therefore, in the present study, we used human IGF-2 (Novus
85 Biologicals, LLC, Co, USA) instead of chicken IGF-2.

86

87 *2.2. Experiment 1: Effects of central administration of IGF-2 on food intake in chicks*

88 Forty-eight 8-day-old chicks were weighed and allocated to four groups based on body

weight (12 birds in each group). IGF-2 was dissolved in 0.85% (w/v) saline solution containing 0.1% (w/v) Evans Blue. The peptide was intracerebroventricularly administered according to the method of Davis et al. [4] at a volume of 10 μ l after three hours of fasting. Chicks were administered with IGF-2 (0, 30, 100, or 300 pmol). Food intake was measured at 60 and 120 min after administration of IGF-2 in each individual cage (260 mm x 185 mm x 148 mm). Feed and water were supplied in plastic boxes (78 mm x 58 mm x 48 mm and 62 mm x 50 mm x 40 mm, respectively). Each feeder filled with food was pre-weighed. A paper was put under the feeder. At in each time point, any spilled food on the paper was collected and food consumption was weighted using an electric digital balance (Readability: 10 mg). Food intake was calculated as follows:

Food intake = (The amount of food decrease in the feeder) - (The amount of spilled food).

At the end of the experiment, the chicks were euthanized by decapitation. Verification of injection was made by observation of the presence of Evans Blue dye in the lateral ventricle. Data from chicks without Evans Blue dye in the lateral ventricle were omitted. Effects of central administration of IGF-2 on food intake were also measured under an *ad libitum* feeding condition.

2.3. *Experiment 2: Effects of six hours of fasting on IGF-related genes' mRNA levels in the liver and hypothalamus of chicks*

Twelve 7-day-old chicks were weighed, allocated based on body weight, and euthanized by decapitation after 0 or 6 hours of fasting. The liver and diencephalons were collected, and preserved in RNAlater tissue storage reagent (Sigma-Aldrich, St. Louis, Mo, USA). The hypothalamus was excised based on reference to a stereotaxic atlas drawn by Kuenzel and Masson as described previously [16]. Total RNA extraction and cDNA synthesis were performed as described previously [13]. Complementary DNA of IGFBPs were amplified using primers as described previously [14]. Complementary DNA of IGF-1 and IGF-1R were amplified using primers as described previously [12]. Complementary DNA of IGF-2, IGF-2R and ribosomal

115 protein S17 (RPS17: an internal standard) were amplified using primers as follows, IGF-2 sense,
116 5'-CCT GGC TCT GCT GGA AAC C-3' ; antisense, 5'-GAG AGG TCA CGC TCT GAC TTG
117 A-3'; IGF-2R sense, 5'-AAC ATC GGG TGT TTC CTA CAA ATA C-3'; antisense, 5'-TGA TTT
118 GGT GCT GCA ATT TCC-3'; RPS17 sense, 5'- GCG GGT GAT CAT CGA GAA GT-3';
119 antisense, 5'-GCG CTT GTT GGT GTG GAA GT-3'. Messenger RNA levels were quantified in
120 duplicate using the Thermo Scientific PikoReal Real time PCR System (Thermo Fisher Scientific
121 Oy, Vantaa, Finland) and SYBR® Premix Ex Taq™ II (Tli RNaseH Plus) (Takara Bio Inc., Shiga,
122 Japan) according to the manufacturer's recommendations.

123

124 2.4. *Experiment 3: Effects of a peroxisome proliferator-activated receptor alpha (PPAR α)*
125 *agonist on IGF-related genes' mRNA levels in chicken hepatoma cells*

126 There is evidence that a PPAR α agonist significantly upregulates hepatic IGFBP-1 [5]
127 and IGFBP-2 [19] expression in mammalian hepatoma cells. Therefore, we examined the effects
128 of the PPAR α agonist WY14643 on IGFBPs expression in chicken hepatoma (LMH) cells.
129 Degenhardt et al [5] demonstrated the direct effect of WY14643 on gene expression in a human
130 hepatoma cell line (HepG2) at two hours after addition of WY14643. We previously showed that
131 50 μ M of WY14643 significantly upregulated the target gene *carnitine palmitoyltransferase 1A*
132 (*CPT1A*) in LMH cells [16]. According to these findings, we confirmed the effect of a two-hour
133 incubation with 50 μ M of WY14643 on LMH cells. Cell culture, RNA extraction, and cDNA
134 synthesis were performed as described previously [16]. Complementary DNA of CPT1A was
135 amplified using primers as described previously [16].

136

137 2.5. *Data analysis*

138 Data in Experiment 1 were analyzed by the Tukey-Kramer test. Data in Experiment 2
139 were analyzed by Student's t-test. All statistical analyses were performed using the commercial
140 package (StatView version 5, SAS Institute, Cary, NC, USA, 1998).

141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166

3. Results

In the present study, we firstly examined the effect of central administration of IGF-2 on food intake in chicks. Intracerebroventricular administration of 300 pmol of IGF-2 significantly suppressed food intake under both three hours of fasting and *ad libitum* feeding conditions (Fig. 1A and B, respectively).

The hypothalamus plays important roles in the central regulation of food intake in mammals [42] and birds [33]. Hence, in order to evaluate the possible role of central IGF-2 and related proteins in the central regulation of food intake in chicks, we next examined the effects of fasting on the mRNA levels of IGF-2, IGF-1R, IGF-2R, and IGFBPs in the hypothalamus. None of these mRNA levels in the hypothalamus of chicks were affected by six hours of fasting (Table 1).

In order to evaluate the possible role of hepatic IGF-2 and IGFBPs production in response to feeding status in chicks, we next examined the effects of fasting on their mRNA levels in the liver. Six hours of fasting did not influence the mRNA levels of IGF-2 but markedly increased the mRNA levels of IGFBP-1 and -2 (Fig. 2). The mRNA levels of IGFBP-3 were also significantly increased.

We finally examined the effects of WY14643 on IGFBPs expression in LMH cells and found that WY14643 significantly increased the mRNA levels of IGFBP-1, as well as those of a *PPARα* target gene CPT1A (Fig. 3).

Thus, our findings suggest that increased hepatic production of IGFBP-1, -2, and -3 may decrease free IGF-2 in the circulation under *ad libitum* feeding conditions, which in turn suppresses the anorexigenic action of IGF-2 in chicks. Our findings also suggest that the upregulation of hepatic IGFBP-1 expression may be involved in the activation of *PPARα* in chicken liver.

4. Discussion

167 We showed that central administration of IGF-1 [12] and IGF-2 (present study)
168 significantly suppressed food intake in chicks. In rats, both IGF-1 and -2 probably cross the
169 blood-brain barrier via IGFs receptors [32]. As described in the Introduction section, IGFBPs bind
170 IGF-1 and -2 in plasma and block their binding to the receptors. In the present study hepatic
171 IGFBP-1, -2, and -3 mRNA levels showed significantly lower levels in the *ad libitum* feeding
172 condition as compared to the fasting condition. All these findings raise the hypothesis that down
173 regulation of IGFBP-1, -2, and -3 production in the liver elevates plasma free IGFs, which in turn
174 facilitates IGFs crossing to the brain and suppressing food intake in chicks.

175 There are three major IGFBPs, IGFBP-28 (28kDa), -34 (34kDa), and -40 (40kDa) in the
176 plasma of chickens. Plasma IGFBP-28 and -34 were increased after 48 hours of fasting, whereas
177 IGFBP-40 did not show any significant change [3]. IGFBP-28, -34, and -40 are suggested to be
178 IGFBP-1, -2, and -3, respectively [7]. In the present study, mRNA levels of IGFBP-1 and -2 in the
179 liver under six hours of fasting condition were 5.47- and 6.95-fold higher than that under the *ad*
180 *libitum* feeding condition. On the other hand, mRNA levels of IGFBP-3 under the fasting
181 condition were only 1.36-fold higher than that under the feeding condition. The effects of feeding
182 on the plasma concentration of free IGFs in chickens have not been investigated. However, these
183 findings suggest that hepatic IGFBP-1 and -2 production may play important roles in the
184 diet-induced changes in plasma free IGFs levels in chickens.

185 Forty eight hours of fasting significantly decreased the plasma total IGF-2 concentration
186 in 9-week-old fat chickens, but not in lean chickens [3]. Sixteen hours of fasting significantly
187 decreased plasma total IGF-2 concentration in 16-week-old lean chickens [2]. In the present study,
188 six hours of fasting did not influence IGF-2 mRNA levels in the liver in 7-day-old layer chicks. It
189 is therefore possible that six hours of fasting is not enough to elevate the plasma total IGF-2
190 concentration in layer chicks, although the plasma free IGF-2 concentration may be changed.
191 Further study is needed to clarify the effects of fasting on the plasma free IGF-2 concentration in
192 chicks.

193 We previously showed that four hours of fasting significantly elevated plasma
194 non-esterified fatty acid (NEFA) and increased the mRNA levels of hepatic PPAR α in chickens
195 [34]. PPAR α is activated by fatty acids [31]. In mammalian hepatoma cells, hepatic IGFBP-1 [5]
196 and IGFBP-2 [19] expression are upregulated by a peroxisome proliferator-activated receptor α
197 (PPAR α). Therefore, we finally examined the effects of the PPAR α agonist WY14643 on the
198 mRNA levels of IGFBPs and a PPAR α target gene, *CPT1A*, in chicken hepatoma cells and found
199 that the PPAR α agonist significantly increased the mRNA levels of IGFBP-1. It is therefore
200 possible that fasting-elevated plasma NEFA upregulates hepatic IGFBP-1 via PPAR α in chickens,
201 which in turn inhibits the anabolic effects of IGFs. Further study is required to clarify the
202 mechanism underlying the fasting-induced upregulation of IGFBP-2 and -3 expression in chicken
203 liver.

204 In the present study, six hours of fasting increased hepatic IGFBP-3 expression in
205 8-day-old layer chicks. On the other hand, we previously showed that six hours of fasting did not
206 influence hepatic IGFBP-3 expression in 8-day-old broiler chicks. These findings raise the
207 hypothesis that fasting induced IGFBP-3 production may suppress the anorexigenic action of
208 circulating IGFs in layer chicks but not in broiler chicks. However, it is well known that broiler
209 chicks eat more food than layer chicks [26]. Therefore, IGFBP-3 may play a minor role in the
210 regulation of food intake in chicks as compared to IGFBP-1 and -2.

211 Tachibana et al. [38] reported that central administration of 477 pmol of cocaine- and
212 amphetamine-regulated transcript (CART) significantly suppressed food intake under an *ad*
213 *libitum* feeding condition, but the same dose of CART did not suppress food intake in chicks
214 fasted for three hours. Shiraishi et al. [37] reported that central administration of 2 ng of porcine
215 insulin significantly suppressed food intake under an *ad libitum* feeding condition, whereas
216 central administration of 10 ng of porcine insulin did not suppress food intake in chicks fasted for
217 three hours. These findings suggest that the feeding condition can influence the sensitivity to
218 appetite regulatory peptide in chicks. However, in the present study, central administration of 300

219 pmol of IGF-2 significantly suppressed food intake under both three hours of fasting and *ad*
220 *libitum* feeding conditions. Six hours of fasting did not influence the mRNA levels of IGF-1R,
221 IGF-2R, and IGF-BPs in the hypothalamus of chicks. It is therefore likely that short term fasting
222 does not influence the anorexigenic effect of IGF-2 in chicks.

223 IGF-2 plays important roles in chick embryonic development [29]. Liu et al. [23]
224 reported that IGF-2 mRNA levels in the liver and skeletal muscles increased during embryonic
225 growth and showed higher levels in the later stages (embryonic days 17-19). A high concentration
226 (60-80 ng/mL) of plasma IGF-2 during embryonic days 13-21 fell to 40-50 ng/mL after hatching
227 [25]. McMurtry [28] also showed that plasma IGF-2 concentration increased before hatching and
228 decreased after hatching. Holzenberger and Lapointe [15] reported that IGF-2 expression in the
229 chicken brain is downregulated shortly after hatching. These findings raise the hypothesis that
230 IGF-2 suppresses appetite before hatching. Thus, it will be interesting to clarify whether
231 downregulation of IGF-2 production after hatching functions as the trigger for appetite induction
232 in neonatal chicks.

233 Interestingly, the anorexigenic action of IGF-1 has been observed in diabetic rats [24], but
234 not in non-diabetic mammals [9, 21]. These findings raise the hypothesis that the appetite regulatory
235 role of IGF-1 is changed by the physiological condition or type of animal. In the present study, we
236 showed that IGF-2 suppresses food intake in chicks. The anorexigenic action of IGF-2 has been
237 observed in non-diabetic rats [24]. It seems likely that the anorexigenic function of IGF-2 may have
238 been well conserved between mammals and birds. However, we also found that central
239 administration of 300 pmol of IGF-2 did not influence food intake in meat type chicks (unpublished
240 data). Higher doses of IGF-2 may suppress food intake in broiler chicks. Further studies is needed
241 to compare the anorexigenic effect of IGF between different types of chicks in a wider range of
242 doses (i.e. 0-3,000pmol).

243 IGF-2 actions are possibly mediated by both IGF-1R and IGF-2R. Our previous
244 observations suggest that hypothalamic POMC and AKT may be involved in the IGF-1-induced

245 anorexigenic pathway in chicks. However, Versteyhe et al. [40] investigated the gene expression
246 regulated via IGF-1 receptor and found significant difference in responses between equipotent
247 concentrations of IGF-1 and -2 in mice fibroblasts. It is therefore likely that IGF-1 and -2 shows
248 have different actions even through the same receptor. Further study is needed to examine the
249 effects of central administration of IGF-2 on the phosphorylation of signaling molecules and
250 expression of appetite regulating neuropeptides in the hypothalamus of chickens.

251

252 5. Conclusion

253 In the present study, we found that central administration of IGF-2 suppressed food
254 intake in chicks. We also showed that hepatic IGFBP-1, -2, and -3 mRNA levels were markedly
255 increased in response to fasting. These findings suggest that IGF-2, IGFBP-1, -2, and -3 may be
256 involved in the regulation of food intake in chickens.

257

258 Fundings

259 This research did not receive any specific grant from funding agencies in the public,
260 commercial, or not-for-profit sectors.

261

262 References

- 263 [1] J.B. Allard, C. Duan, IGF-binding proteins: why do they exist and why are there so many?
264 Front Endocrinol. 9 (2018) 117. [https://doi: 10.3389/fendo.2018.00117](https://doi.org/10.3389/fendo.2018.00117).
- 265 [2] C. Beccavin, B. Chevalier, L.A. Cogburn, J. Simon, M.J. Duclos. Insulin-like growth factors
266 and body growth in chickens divergently selected for high or low growth rate. J.
267 Endocrinol. 168 (2001) 297-306. [http://doi: 10.1677/joe.0.1680297](http://doi.org/10.1677/joe.0.1680297).
- 268 [3] C. Beccavin, B. Chevalier, J. Simon, M.J. Duclos, Circulating insulin-like growth factors
269 (IGF-I and -II) and IGF binding proteins in divergently selected fat or lean chickens: effect
270 of prolonged fasting, Growth Horm. IGF Res. 9 (1999) 187-194. [https://doi:](https://doi.org/10.1054/gthi.1999.0187)

271 10.1054/ghir.1999.0109.

272 [4] J.L. Davis, D.T. Masuoka, L.K. Gerbrandt, A. Cherkin, Autoradiographic distribution of

273 L-proline in chicks after intracerebral injection, *Physiol. Behav.* 22 (1979) 693-695.

274 [https://doi: 10.1016/0031-9384\(79\)90233-6](https://doi.org/10.1016/0031-9384(79)90233-6).

275 [5] T. Degenhardt, M. Matilainen, K.H. Herzig, T.W. Dunlop, C. Carlberg, The insulin-like

276 growth factor-binding protein 1 gene is a primary target of peroxisome

277 proliferator-activated receptors, *J. Biol. Chem.* 281(2006) 39607-39619. [http://doi:](http://doi.org/10.1074/jbc.M605623200)

278 [10.1074/jbc.M605623200](http://doi.org/10.1074/jbc.M605623200).

279 [6] M.J. Duclos, Insulin-like growth factor-I (IGF-1) mRNA levels and chicken muscle growth, *J.*

280 *Physiol. Pharmacol.* 56 (2005) 25-35.

281 [7] M.J. Duclos, C. Beccavin, J. Simon, Genetic models for the study of insulin-like growth

282 factors (IGF) and muscle development in birds compared to mammals, *Domest. Anim.*

283 *Endocrinol.* 17 (1999) 231-243. [https://doi: 10.1016/s0739-7240\(99\)00040-5](https://doi.org/10.1016/s0739-7240(99)00040-5).

284 [8] M.J. Duclos, C. Goddard, Insulin-like growth factor receptors in chicken liver membranes:

285 binding properties, specificity, developmental pattern and evidence for a single receptor

286 type, *J. Endocrinol.* 125 (1990) 199-206. [https://doi: 10.1677/joe.0.1250199](https://doi.org/10.1677/joe.0.1250199).

287 [9] L.A. Foster, N.K. Ames, R.S. Emery, Food intake and serum insulin responses to

288 intraventricular infusions of insulin and IGF-I, *Physiol. Behav.* 50 (1991) 745-749.

289 [https://doi: 10.1016/0031-9384\(91\)90012-d](https://doi.org/10.1016/0031-9384(91)90012-d).

290 [10] J. Frystyk, K. Højlund, K.N. Rasmussen, S.P. Jørgensen, M. Wildner-Christensen, H. Ørskov,

291 Development and clinical evaluation of a novel immunoassay for the binary complex of

292 IGF-I and IGF-binding protein-1 in human serum, *J. Clin. Endocrinol. Metab.* 87 (2002)

293 260-266. [https://doi: 10.1210/jcem.87.1.8147](https://doi.org/10.1210/jcem.87.1.8147).

294 [11] J. Frystyk, P. Ivarsen, R.K. Støving, R. Dall, T. Bek, C. Hagen, H. Ørskov, Determination of

295 free insulin-like growth factor-I in human serum: comparison of ultrafiltration and

296 direct immunoradiometric assay, *Growth Horm. IGF Res.* 11 (2001) 117-127.

- 297 [https://doi: 10.1054/ghir.2001.0197](https://doi.org/10.1054/ghir.2001.0197).
- 298 [12] S. Fujita, K. Honda, D. Hiramoto, M. Gyu, M. Okuda, S. Nakayama, M. Yamaguchi, T.
 299 Saneyasu, H. Kamisoyama, Central and peripheral administrations of insulin-like
 300 growth factor-1 suppress food intake in chicks, *Physiol. Behav.* 179 (2017) 308-312.
 301 [https://doi: 10.1016/j.physbeh.2017.07.001](https://doi.org/10.1016/j.physbeh.2017.07.001).
- 302 [13] S. Fujita, K. Honda, M. Yamaguchi, S. Fukuzo, T. Saneyasu, H. Kamisoyama, Role of
 303 insulin-like growth factor-1 in the central regulation of feeding behavior in chicks, *J.*
 304 *Poult. Sci.* 56 (2019) 270-276. [https://doi: 10.2141/jpsa.0180127](https://doi.org/10.2141/jpsa.0180127).
- 305 [14] S. Fujita, M. Yamaguchi, D. Hiramoto, T. Saneyasu, K. Honda, H. Kamisoyama, Effects of
 306 fasting and refeeding on the mRNA levels of insulin-like growth factor-binding proteins
 307 in chick liver and brain, *J. Poult. Sci.* 55 (2018) 269-273. [https://doi:](https://doi.org/10.2141/jpsa.0180005)
 308 [10.2141/jpsa.0180005](https://doi.org/10.2141/jpsa.0180005).
- 309 [15] M. Holzenberger, F. Lapointe, Expression of insulin-like growth factor-I (IGF-I) and IGF-II
 310 in the avian brain: relationship of in situ hybridization patterns with IGF type 1 receptor
 311 expression, *Int. J. Dev. Neurosci.* 18 (2000) 69-82. [https://doi:](https://doi.org/10.1016/s0736-5748(99)00076-3)
 312 [10.1016/s0736-5748\(99\)00076-3](https://doi.org/10.1016/s0736-5748(99)00076-3).
- 313 [16] K. Honda, T. Saneyasu, K. Aoki, T. Shimatani, T. Yamaguchi, H. Kamisoyama, Correlation
 314 analysis of hypothalamic mRNA levels of appetite regulatory neuropeptides and several
 315 metabolic parameters in 28-day-old layer chickens, *Anim. Sci. J.* 86 (2015) 517-522.
 316 [https://doi: 10.1111/asj.12320](https://doi.org/10.1111/asj.12320).
- 317 [17] K. Honda, T. Saneyasu, K. Sugimoto, H. Kurachi, K. Takagi, S. Kamisoyama, Role of
 318 peroxisome proliferator-activated receptor alpha in the expression of hepatic fatty acid
 319 oxidation-related genes in chickens, *Anim. Sci. J.* (2016) 61-66. [https://doi:](https://doi.org/10.1111/asj.12392)
 320 [10.1111/asj.12392](https://doi.org/10.1111/asj.12392).
- 321 [18] S. Koduru, S. Yadavalli, S.K. Nadimpalli, Mannose 6-phosphate receptor (MPR 300)
 322 proteins from goat and chicken bind human IGF-II, *Biosci. Rep.* 26 (2006) 101-112.

- 323 [https://doi: 10.1007/s10540-006-9013-0](https://doi.org/10.1007/s10540-006-9013-0).
- 324 [19] H.S. Kang, M.Y. Kim, S.J. Kim, J.H. Lee, Y.D. Kim, Y.K. Seo, J.H. Bae, G.T. Oh, D.K. Song,
 325 Y.H. Ahn, S.S. Im. Regulation of IGFBP-2 expression during fasting. *Biochem. J.* 467
 326 (2015) 453-460. [http://doi: 10.1042/BJ20141248](http://doi.org/10.1042/BJ20141248).
- 327 [20] K. Kita, K. Hangsanet, T. Shibata, M.A. Conlon, T. Sasaki, N. Saito, J. Okumura, Refeeding
 328 increases hepatic insulin-like growth factor-I (IGF-I) gene expression and plasma IGF-I
 329 concentration in fasted chicks, *Br. Poult. Sci.* 39 (1998) 679-682. [https://doi:](https://doi.org/10.1080/00071669888566)
 330 10.1080/00071669888566.
- 331 [21] T.J. Lauterio, L. Marson, W.H. Daughaday, C.A. Baile, Evidence for the role of insulin-like
 332 growth factor II (IGF-II) in the control of food intake, *Physiol. Behav.* 40 (1987) 755-758.
 333 [https://doi: 10.1016/0031-9384\(87\)90279-4](https://doi.org/10.1016/0031-9384(87)90279-4).
- 334 [22] S. Leili, F.C. Buonomo, C.G. Scanes, The effects of dietary restriction on insulin-like growth
 335 factor (IGF)-I and II, and IGF-binding proteins in chickens, *Proc. Soc. Exp. Biol. Med.*
 336 216 (1997) 104-111. [https://doi: 10.3181/00379727-216-44162a](https://doi.org/10.3181/00379727-216-44162a).
- 337 [23] Y. Liu, W. Guo, Z. Pu, X. Li, X. Lei, J. Yao, X. Yang, Developmental changes of insulin-like
 338 growth factors in the liver and muscle of chick embryos, *Poult. Sci.* 95 (2016) 1396-1402.
 339 [https://doi: 10.3382/ps/pew043](https://doi.org/10.3382/ps/pew043).
- 340 [24] H. Lu, B. Martinez-Nieves, K. Lapanowski, J. Dunbar, Intracerebroventricular insulin-like
 341 growth factor-1 decreases feeding in diabetic rats, *Endocrine* 14 (2001) 349-352.
 342 [https://doi: 10.1385/ENDO:14:3:349](https://doi.org/10.1385/ENDO:14:3:349).
- 343 [25] J.W. Lu, J.P. McMurtry, C.N. Coon, Developmental changes of plasma insulin, glucagon,
 344 insulin-like growth factors, thyroid hormones, and glucose concentrations in chick
 345 embryos and hatched chicks, *Poult. Sci.* 86 (2007) 673-683. [https://doi:](https://doi.org/10.1093/ps/86.4.673)
 346 10.1093/ps/86.4.673.
- 347 [26] M. Mahagna, I. Nir, Comparative development of digestive organs, intestinal disaccharidases
 348 and some blood metabolites in broiler and layer-type chicks after hatching, *Br. Poult. Sci.*

349 37 (1996) 359-371. <https://doi: 10.1080/00071669608417867>.

350 [27] U. Matzner, A. Hille-Rehfeld, K. von Figura, R. Pohlmann, Expression of mannose
351 6-phosphate receptors in chicken, *Dev. Dyn.* 207 (1996) 11-24. [https://doi:](https://doi: 10.1002/(SICI)1097-0177(199609)207:1<11::AID-AJA2>3.0.CO;2-Z)
352 10.1002/(SICI)1097-0177(199609)207:1<11::AID-AJA2>3.0.CO;2-Z.

353 [28] J.P. McMurtry, Nutritional and developmental roles of insulin-like growth factors in poultry,
354 *J. Nutr.* 128 (1998) 302S-305S. <https://doi: 10.1093/jn/128.2.302S>.

355 [29] J.P. McMurtry, G.L. Francis, Z. Upton, Insulin-like growth factors in poultry, *Domest. Anim.*
356 *Endocrinol.* 14 (1997) 199-229. [https://doi: 10.1016/s0739-7240\(97\)00019-2](https://doi: 10.1016/s0739-7240(97)00019-2).

357 [30] J.P. McMurtry, G.L. Francis, Z. Upton, P.E. Walton, G. Rosselot, T.J. Caperna, D.M. Brocht,
358 Plasma clearance and tissue distribution of labelled chicken and human IGF-I and IGF-II
359 in the chicken, *J. Endocrinol.* 150 (1996) 149-160. <https://doi: 10.1677/joe.0.1500149>.

360 [31] M.T. Nakamura, B.E. Yudell, J.J. Loo. Regulation of energy metabolism by long-chain fatty
361 acids. *Prog. Lipid Res.* 53 (2014) 124-144. [http:// doi: 10.1016/j.plipres.2013.12.001](http://doi: 10.1016/j.plipres.2013.12.001).

362 [32] R.R. Reinhardt, C.A. Bondy, Insulin-like growth factors cross the blood-brain barrier,
363 *Endocrinology* 135 (1994) 1753-1761. <https://doi: 10.1210/endo.135.5.7525251>.

364 [33] M.P. Richards, M. Proszkowiec-Weglarz, Mechanisms regulating feed intake, energy
365 expenditure, and body weight in poultry, *Poult. Sci.* 86 (2007) 1478-1490. [https://doi:](https://doi: 10.1093/ps/86.7.1478)
366 10.1093/ps/86.7.1478.

367 [34] T. Saneyasu, M. Shiragaki, K. Nakanishi, H. Kamisoyama, K. Honda. Effects of short term
368 fasting on the expression of genes involved in lipid metabolism in chicks. *Comp. Biochem.*
369 *Physiol. B: Biochem. Mol. Biol.* 165 (2013) 114-118. [http://doi:](http://doi: 10.1016/j.cbpb.2013.03.005)
370 10.1016/j.cbpb.2013.03.005.

371 [35] R. Saxena, V.K. Saxena, V. Tripathi, N.A. Mir, K. Dev, J. Begum, R. Agarwal, A. Goel,
372 Dynamics of gene expression of hormones involved in the growth of broiler chickens in
373 response to the dietary protein and energy changes, *Gen. Comp. Endocrinol.* 288 (2020)
374 113377. <https://doi: 10.1016/j.ygcen.2019.113377>.

- 375 [36] M. Shimizu, W.W. Dickhoff, Circulating insulin-like growth factor binding proteins in fish:
376 Their identities and physiological regulation, *Gen. Comp. Endocrinol.* 252 (2017) 150-161.
377 [https://doi: 10.1016/j.ygcen.2017.08.002](https://doi.org/10.1016/j.ygcen.2017.08.002).
- 378 [37] J. Shiraishi, K. Yanagita, F. Nishikawa, Y. Tahara, M. Fujita, J.P. McMurtry, T. Bungo, A
379 comparison of the anorexigenic effects of chicken, porcine, human and bovine insulin on
380 the central nervous system of chicks, *J. Poult. Sci.* 46 (2009) 144-147. [https://doi:](https://doi.org/10.2141/jpsa.46.144)
381 [10.2141/jpsa.46.144](https://doi.org/10.2141/jpsa.46.144).
- 382 [38] T. Tachibana, T. Takagi, S. Tomonaga, A. Ohgushi, R. Ando, D.M. Denbow, M. Furuse,
383 Central administration of cocaine- and amphetamine-regulated transcript inhibits food
384 intake in chicks, *Neurosci. Lett.* 337 (2003) 131-134. [https:// doi:](https://doi.org/10.1016/s0304-3940(02)01321-6)
385 [10.1016/s0304-3940\(02\)01321-6](https://doi.org/10.1016/s0304-3940(02)01321-6).
- 386 [39] F.Z. Upton, G.L. Francis, K. Kita, J.C. Wallace, F.J. Ballard, Production and characterization
387 of recombinant chicken insulin-like growth factor-II from *Escherichia coli*, *J. Mol.*
388 *Endocrinol.* 14 (1995) 79-90. [https://doi: 10.1677/jme.0.0140079](https://doi.org/10.1677/jme.0.0140079).
- 389 [40] S. Versteyhe, B. Klaproth, R. Borup, J. Palsgaard, M. Jensen, S.G. Gray, P. de Meyts, IGF-I,
390 IGF-II, and insulin stimulate different gene expression responses through binding to the
391 IGF-I receptor, *Front. Endocrinol.* 4 (2013) 98. [https://doi: 10.3389/fendo.2013.00098](https://doi.org/10.3389/fendo.2013.00098).
- 392 [41] X. Wan, S. Wang, J. Xu, L. Zhuang, K. Xing, M. Zhang, X. Zhu, L. Wang, P. Gao, Q. Xi, J.
393 Sun, Y. Zhang, T. Li, G. Shu, Q. Jiang. Dietary protein-induced hepatic IGF-1 secretion
394 mediated by PPARgamma activation. *PLoS One* 12 (2017) e0173174. [http://doi:](http://doi.org/10.1371/journal.pone.0173174)
395 [10.1371/journal.pone.0173174](http://doi.org/10.1371/journal.pone.0173174).
- 396 [42] S.C. Woods, The control of food intake: behavioral versus molecular perspectives, *Cell*
397 *Metab.* 9 (2009) 489-498. [https://doi: 10.1016/j.cmet.2009.04.007](https://doi.org/10.1016/j.cmet.2009.04.007).

400 **Figure captions**

401 Fig. 1. Effects of central administration of insulin-like growth factor-2 on food intake in chicks.
402 A: IGF-2 was administered after three hours of fasting. B: IGF-2 was administered under an *ad*
403 *libitum* feeding condition. Data represent means \pm S.E.M. The number of chicks used is shown in
404 parentheses. * indicates significant difference with respect to the 0 pmol group ($P < 0.05$).

405

406 Fig. 2. Effects of six hours of fasting on the mRNA levels of insulin-like growth factor-related
407 genes in the livers of chicks. Data were normalized to the respective average of each feeding group.
408 Data represent means \pm S.E.M. of six chicks. ** indicates significant difference with respect to the
409 feeding group ($P < 0.01$).

410

411 Fig. 3. Effects of WY14643 on the mRNA levels of insulin-like growth factor-related genes in
412 chicken hepatoma cells. Data represent means \pm S.E.M. of four wells in each group. * indicates
413 significant difference with respect to the control group ($P < 0.05$).

414

Table 1. Effects of six hours of fasting on the mRNA levels of insulin-like growth factor-related genes in the hypothalamus of chicks

	Feeding	Fasting
<i>IGF-2</i>	1.00 ± 0.08	0.85± 0.10
<i>IGF-1R</i>	1.00 ± 0.06	0.87± 0.05
<i>IGF-2R</i>	1.00 ± 0.03	1.01 ± 0.07
<i>IGFBP-1</i>	1.00 ± 0.02	1.01 ± 0.04
<i>IGFBP-2</i>	1.00 ± 0.08	0.82 ± 0.16
<i>IGFBP-3</i>	1.00 ± 0.03	0.89 ± 0.04
<i>IGFBP-4</i>	1.00 ± 0.11	0.75 ± 0.06
<i>IGFBP-5</i>	1.00 ± 0.01	0.99 ± 0.06

Data were normalized to respective average of each feeding group. Data represent means ± S.E.M. of six chicks.

Figure 1





Figure 2




Figure 3

