

PDF issue: 2025-07-06

## Ultrasound assessment of muscle mass has potential to identify patients with low muscularity at intensive care unit admission: A retrospective study

Arai, Yuta ; Nakanishi, Nobuto ; Ono, Yuko ; Inoue, Shigeaki ; Kotani, Joji ; Harada, Masafumi ; Oto, Jun

(Citation) Clinical Nutrition ESPEN, 45:177-183

(Issue Date) 2021-10

(Resource Type) journal article

(Version) Accepted Manuscript

## (Rights)

© 2021 European Society for Clinical Nutrition and Metabolism. Published by Elsevier Ltd.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/

## (URL)

https://hdl.handle.net/20.500.14094/90008912



Ultrasound assessment of muscle mass has potential to identify patients with low muscularity at intensive care unit admission: A retrospective study

Yuta Arai, MD<sup>1,2,\*</sup>, Nobuto Nakanishi, MD<sup>3,4,\*</sup>, Yuko Ono, MD, PhD<sup>3</sup>, Shigeaki Inoue, MD, PhD

<sup>3</sup>, Joji Kotani, MD, PhD<sup>3</sup>, Masafumi Harada, MD, PhD<sup>2</sup>, Jun Oto, MD, PhD<sup>4</sup>

\*Contributed equally as co-first authors

 Emergency and Disaster Medicine, Tokushima University Hospital, 2-50-1 Kuramoto, Tokushima 770-8503, Japan

 Department of Radiology, Tokushima University Hospital, 2-50-1 Kuramoto, Tokushima 770-8503, Japan

3. Department of Disaster and Emergency Medicine, Graduate School of Medicine, Kobe University, 7-5-2 Kusunoki, Chuo-ward, Kobe, 650-0017, Japan

4. Emergency and Critical Care Medicine, Tokushima University Hospital, 2-50-1 Kuramoto, Tokushima 770-8503, Japan

Correspondence to: Nobuto Nakanishi, MD

Department of Disaster and Emergency Medicine, Graduate School of Medicine, Kobe University,

7-5-2 Kusunoki, Chuo-ward, Kobe, 650-0017, Japan

Emergency and Critical Care Medicine, Tokushima University Hospital, 2-50-1 Kuramoto, Tokushima 770-8503, Japan

TEL: 088-633-9347

FAX: 088-633-9339

Email: nobuto\_nakanishi@yahoo.co.jp

#### Abstract

Background & aims: Muscle mass is an important biomarker of survival from a critical illness; however, there is no widely accepted method for routine assessment of low muscularity at intensive care unit (ICU) admission. We hypothesize that ultrasound-based partial muscle mass assessments can reflect the trunk muscle mass. Therefore, we aimed to investigate whether ultrasound muscle mass measurements could reflect trunk muscle mass and identify patients with low muscularity. Methods: We performed a retrospective analysis of prospectively obtained ultrasound data at ICU admission. We included patients who underwent computed tomography (CT) imaging at the third lumbar vertebra (L3) within 2 days before and 2 days after ICU admission. Primary outcomes included the correlation between the femoral muscle mass measurements using ultrasound and the cross-sectional area (CSA) at L3 obtained by CT. Low muscularity was defined as a skeletal muscle index of 29.0  $\text{cm}^2/\text{m}^2$  for males and 36.0  $\text{cm}^2/\text{m}^2$  for females. Secondary outcomes included the correlation with the ultrasound measurements of the biceps brachii muscle mass and diaphragm thickness. Results: Among 133 patients, 89 underwent CT imaging, which included the L3. The patient mean age was  $72 \pm 13$  years, and 60 patients (67%) were male. The correlation between the femoral muscle ultrasound and CT was  $\rho = 0.57$  (p < 0.01, n = 89) and  $\rho = 0.48$  (p < 0.01, n = 89) for quadriceps muscle layer thickness and rectus femoris muscle CSA, and these had the discriminative power to assess low muscularity, with the areas under the curve of 0.84 and 0.76, respectively. The ultrasound measurements of the biceps brachii muscle mass and diaphragm thickness were correlated with CT imaging [ $\rho = 0.57-0.60$  (p < 0.01, n = 52) and  $\rho = 0.35$  (p < (0.01, n = 79)]. Conclusions: Ultrasound measurements of muscle mass were correlated with CT measurements, and the measurements of femoral muscle mass were useful to assess low muscularity at ICU admission.

Trial registration: UMIN000044032 (retrospectively registered on April 25, 2021)

Keywords: rectus femoris muscle, muscle mass, ultrasound, computed tomography, intensive care unit

## Abbreviations

ICU: intensive care unit; CT: computed tomography; L3: the third lumbar vertebra; CSA: crosssectional area; AUC: area under the receiver operating characteristic curve; IQR: interquartile range; CI: confidence interval Introduction

Muscle mass is an important biomarker of survival from a critical illness because low muscularity at intensive care unit (ICU) admission is associated with higher mortality and ICU-acquired weakness [1, 2]. The notion that a state of low muscularity is representative of sarcopenia has recently been gaining increased attention [3]. The identification of low muscularity is important in planning nutrition and rehabilitation management strategies. Both are needed to maintain muscle mass during ICU hospitalization. However, to date, there is no widely accepted method for routine assessment of low muscularity at ICU admission.

Several methods have been used to assess muscle mass [4]. In general, dual-energy X-ray absorptiometry and bioelectrical impedance analyses have been used to assess whole-body muscle mass [5]. However, these indirect muscle mass assessments are inaccurate in critically ill patients because these are influenced by dynamic fluid changes [6-8]. Furthermore, it is challenging to transport patients to a dual-energy X-ray absorptiometry machine. During a critical illness, computed tomography (CT) is considered the gold standard to assess muscle mass because it can visually separate muscle mass from other tissues [9]. Although CT is a reliable method to measure muscle mass, prospective CT evaluation is infeasible because of patient transportation risks and radiation exposure [10]. By contrast, ultrasound is an emerging tool used to measure muscle mass noninvasively at the bedside [11]. Although ultrasound is used to assess limb muscles, it is unclear whether partial muscle mass assessments reflect trunk muscle mass in critically ill patients. To validate the ultrasound assessments of muscularity, it is important to show the measurement correlation between ultrasound and CT.

The femoral muscle is assessed commonly using ultrasound, in which muscle thickness or cross-sectional area (CSA) measurements are performed [4]. However, it is unclear whether these mass measurements reflect the trunk or partial muscle mass in critically ill patients. Given that a previous study reported that the CSA of the rectus femoris is preferable than thickness for physical functions [12, 13], we hypothesized that CSA is associated with the trunk muscle mass. We retrospectively evaluated the muscle mass area at the level of the third lumbar vertebra (L3) using CT, and compared it with those obtained from prospectively obtained ultrasound data at ICU admission. This study aimed to investigate whether ultrasound muscle mass measurements were correlated with CT measurements, and could identify patients with low muscularity at ICU admission.

#### Materials & Methods

#### Study design

This two-center retrospective study was conducted in the mixed medical/surgical ICUs of Tokushima University Hospital and Tokushima Prefectural Central Hospital. The study complied with the principles of the Declaration of Helsinki and was approved by the clinical research ethics committees of Tokushima University Hospital (approval number 2593) and Tokushima Prefectural Central Hospital (approval number 1739). Prospectively obtained data from May 2016 to June 2020 were retrospectively analyzed. This study was retrospectively registered at UMIN-Clinical Trials Registry (UMIN000044032). At the time of data acquisition, written informed consent was obtained from patients or their relatives. A part of this study was previously published [7, 14, 15].

#### Study population

We included patients who met the following criteria: (1) adults ( $\geq 18$  years) admitted to ICU; (2) those expected to stay in ICU for >5 days; (3) those who underwent the ultrasound assessments of

the rectus femoris muscle at the day of ICU admission; and (4) those who underwent the CT assessments of the L3 within 2 days before and after ICU admission. The following patients were excluded from the studies: those with (1) primary neuromuscular disease and (2) obstacles at the ultrasound measurement site.

#### Ultrasound

We used a linear transducer and performed B-mode imaging. The details of equipment were shown in Table S1. The measurements were performed at the dominant limb or right limb under no information. Elbows and knees were extended in the supine position, and the transducer was placed perpendicular to the long axis of the limbs. The measurements included the quadriceps muscle layer thickness and rectus femoris muscle CSA. Measurements were performed midway between the anterior superior iliac spine and the proximal end of the patella as a common landmark [4]. The quadriceps muscle layer thickness, including the rectus femoris and vastus intermedius muscles, was measured from the superficial fascia of the rectus femoris to the uppermost part of the femur (Figure S1). CSA was measured by outlining the area shown in the transverse plane (Figure S2). The elbow flexor muscle thickness and biceps brachii muscle CSA were measured at a distance equal to two-thirds of the distance from the acromion to the antecubital crease as a common landmark [16, 17]. The elbow flexor muscle thickness, including the biceps brachii and brachialis muscles, was measured from the superficial fascia of the biceps brachii muscle to the uppermost part of the humerus. The diaphragm was measured at the end expiration on the right chest wall at the zones of apposition 0.5-2 cm below the costophrenic sinus between the anteroaxillary and the midaxillary lines (Figure S3) [18]. The ultrasound measurements were performed three times by a physician (N.N.), and the median value was used for evaluation. Generous

amounts of contact gel was used for minimal compression because probe compression strength affects measurement accuracy [19]. The reliability of measurements was confirmed by another ICU physician, as reported previously (Table S2) [14, 15].

CT

CT was used to evaluate muscle mass at the L3 level, which has been reported to correlate with the trunk muscle mass in patients with cancer [20]. A board-certified radiologist (A.Y.) retrospectively measured the total muscle mass in the CT image at the middle point of the L3 where transverse processes were visualized (Figure S4). At this slice level, the total muscle area included the psoas, quadratus lumborum, transversus abdominis, external and internal obliques, and rectus abdominis muscles. CT images acquired within 2 days before and after ICU admission were included in the analyses, and examinations performed closest to the day of ICU admission were used for comparisons in patients with multiple CT examinations. The radiologist was blinded to all clinical characteristics. All images were analyzed using ImageJ software (National Institutes of Health, Bethesda, MD, USA) [21]. The reliability of measurements was confirmed in 10 patients by 2 examiners (Y.A. and N.N.). The intraclass correlation coefficient was  $\rho = 0.98$  (p < 0.01), and the interclass value was  $\rho = 0.94$  (p < 0.01). The Bland–Altman plot yielded a bias of  $-1.24 \pm 1.58$  and -4.83 to 2.34 at the 95% confidence interval (CI) regarding intraobserver reproducibility and a bias of  $-0.94 \pm 2.67$  and -6.98 to 5.10 at the 95% CI regarding interobserver reproducibility.

Low muscularity was identified using the skeletal muscle index, which is calculated by dividing the CT image CSA at L3 by height squared. The sex-specific cutoff point was set to 29.0  $\text{cm}^2/\text{m}^2$  for males and 36.0  $\text{cm}^2/\text{m}^2$  for females, which are one of the commonly used cutoff points for low muscularity in the Asian population [22]. These cutoff points were previously reported to

be important in Japanese patients with cancer [23].

#### Outcomes

The primary outcome was the relationship between the ultrasound assessments of the quadriceps muscle layer thickness or rectus femoris muscle CSA and the CT assessment of L3 CSA. We also assessed whether these ultrasound assessments can predict low muscularity in the same manner as that assessed by CT. Secondary outcomes of this study included the relationship with the ultrasound measurements of elbow flexor muscle thickness, biceps brachii muscle CSA, the sum of rectus femoris and biceps brachii muscle CSA, and diaphragm thickness.

#### Statistical analyses

Continuous variables were presented as the mean (standard deviations) or median values [interquartile ranges (IQRs)], whereas categorical data were presented as counts and proportions. Variables were compared using either the t-test or the Mann–Whitney U test. The Spearman correlation coefficient was used to investigate relationships in primary and secondary outcomes. The area under the receiver operating characteristic curve (AUC) was generated to determine the cutoff values of ultrasound assessments for low muscularity. Sample size was not calculated a priori and was based as per feasibility because of the retrospective nature of the study. For reproducibility, the Spearman correlation coefficient and the Bland–Altman plot were determined using JMP statistical software, version 13.1.0 (SAS Institute Inc., Cary, NC, USA).

#### Results

In total, 133 patients underwent ultrasound measurements of the femoral muscle mass. Among

them, 89 patients underwent CT imaging at the L3 level within 2 days of ICU admission (Figure 1). Fifty-nine patients underwent CT examinations immediately after ICU admission. CT examinations were performed in 10 and 4 patients on days 1 and 2 after ICU admission, respectively, and in 13 and 3 patients on days 1 and 2 days before ICU admission, respectively. The time between the ultrasound and CT assessments was 0 day (IQR, 0–1 day).

The patient characteristics are summarized in Table 1. The patient mean age was  $72 \pm 13$  years, and 60 patients (67%) were male. The median Acute Physiology and Chronic Health Evaluation II score was 27 (IQR, 24–30), and the median length of ICU stay was 7 (IQR, 5–14) days. Seventy-eight (88%) patients were mechanically ventilated, and 15 (16%) were admitted postoperatively. Low muscularity, assessed by CT, was observed in 24 (27%) patients. The median thickness of the quadriceps muscle layer and CSA of the rectus femoris muscle were 2.4 (IQR, 1.8–3.1) cm and 4.9 (IQR, 3.9–6.5) cm<sup>2</sup>, respectively. There was a significant difference in the body mass index (p < 0.01), quadriceps muscle layer thickness (p < 0.01), rectus femoris muscle CSA (p < 0.01), and muscle mass on CT between low and normal muscularity (p < 0.01).

The ultrasound measurements of the femoral muscle at ICU admission were correlated with the muscle mass at the L3 measured by CT (Figure 2). The correlations of the quadriceps muscle layer thickness and rectus femoris muscle CSA were  $\rho = 0.57$  (p < 0.01, n = 89) and  $\rho = 0.48$  (p < 0.01, n = 89), respectively. The thickness of the quadriceps muscle layer thickness had the discriminative power to assess low muscularity at the AUC of 0.84 (95% CI, 0.74–0.94), in which the cutoff value of 2.0 cm had a sensitivity of 83.3% and specificity of 78.5% (Figure 3). By contrast, the CSA of the rectus femoris muscle had the discriminative power to assess low muscularity at the AUC of 0.76 (95% CI, 0.65–0.88), in which the cutoff value of 4.7 cm<sup>2</sup> had a

sensitivity of 79.2% and specificity of 66.2%.

Among the secondary outcomes, the elbow flexor muscle thickness, biceps brachii muscle CSA, the sum of rectus femoris and biceps brachii muscle CSA, and diaphragm muscles were assessed in 52, 52, 52, and 79 of the 89 patients, respectively (Figure 4). The ultrasound measurements of these at ICU admission were correlated with the muscle mass at the L3 measured by CT. The elbow flexor muscle thickness and biceps brachii muscle CSA yielded the correlations of  $\rho = 0.57$  (p < 0.01, n = 52) and  $\rho = 0.60$  (p < 0.01, n = 52), respectively. The sum of rectus femoris and biceps brachii muscle CSA yielded the correlation of  $\rho = 0.57$  (p < 0.01, n = 52). The thickness of the diaphragm yielded a correlation of  $\rho = 0.35$  (p < 0.01, n = 79).

#### Discussion

In this study on 89 critically ill patients, we investigated whether ultrasound-based quadriceps muscle layer thickness and rectus femoris muscle CSA measurements are indicators of low muscularity. Contrary to our hypothesis, both the thickness and CSA were good indicators of low muscularity at ICU admission. Moreover, the biceps brachii muscle and diaphragm were weak-to-moderate indicators of trunk muscle mass. Given that CT is not routinely available to critically ill patients, the ultrasound measurement of the quadriceps muscle layer thickness and rectus femoris muscle CSA can be an alternative for the noninvasive assessment of low muscularity at ICU admission.

This study provides several important intellectual data. First, we found that both the quadriceps muscle layer thickness and rectus femoris muscle CSA are good indicators of the trunk muscle mass. Previous studies investigated the quadriceps muscle thickness and showed that its thickness was correlated with muscle mass measured using CT [24, 25]. However, differences

between the thickness and CSA had not been clarified in previous studies. Although thickness measurement is not correlated with functional impairments [12, 13], the thickness measurement reflects the trunk muscle mass. Second, this study sets the standard values of the quadriceps muscle layer thickness and rectus femoris muscle CSA at ICU admission. Few studies have investigated subject groups using ultrasound at ICU admission. The quadriceps muscle layer thickness of 2.0 cm and the rectus femoris muscle CSA of 4.7 cm<sup>2</sup> can be set as cutoff values for low muscularity at ICU admission. Hida et al. investigated the cutoff value of the rectus femoris muscle for low muscularity in healthy volunteers; however, the values in critically ill patients have not been reported [26]. We showed the standard value of quadriceps muscle layer thickness and rectus femoris muscle CSA at ICU admission in all 133 patients (Table S3), which may contribute to the clinical application of ultrasound muscle mass assessment in critically ill patients.

Of note, the biceps brachii muscle mass was correlated with the trunk muscle mass. This finding is important because the muscle mass assessments of the biceps brachii muscle may replace the femoral muscle for muscularity assessments. The extension of the lower limb for femoral muscle measurements requires critically ill patients to be placed in a supine position because the bed angle affects lower limb extension [27]. During a critical illness, particularly at ICU admission, the flat position may be risky in some patients. By contrast, the biceps brachii muscle can be extended, regardless of the bed angle. Furthermore, the upper limb can be measured more easily because the biceps brachii muscle is exposed to the outside, whereas the femoral muscle measurements need preparation pertaining to the removal of the patient's clothing. The upper limb circumference has been used to calculate the upper limb muscle mass; however, it is not accurate to evaluate the muscle mass in critically ill patients [28]. This is reasonable because the circumference indirectly evaluates the upper limb muscle mass, whereas ultrasound evaluates it

directly. Therefore, the ultrasound biceps brachii muscle measurement is a promising method for muscle mass assessments.

Contrary to previous studies [29, 30], the sum of rectus femoris and biceps brachii muscle CSA did not improve the correlation with the trunk muscle mass. Campbell et al. reported that the sum of elbow flexor, anterior forearm, and quadriceps femoris thicknesses was useful to evaluate muscularity [29]. Likewise, Lambell et al. reported that the sum of elbow flexor, bilateral quadriceps femoris thicknesses, and patient characteristics improved the correlation with the trunk muscle mass [30]. The utility to add muscle mass may depend on the measurement site and the content to be added. We did not analyze which combination improves the correlation because this was not our primary purpose. Therefore, this result needs further investigation. However, at the very least, this result may indicate that a single site muscle mass measurement is sufficient to assess the trunk muscle mass in critically ill patients.

In addition to the biceps brachii muscle, the diaphragm was also correlated with muscularity. We found that the diaphragm thickness differs among critically ill patients. Sklar et al. reported that a diaphragm thickness of <2.3 mm is associated with prolonged mechanical ventilation and mortality [31]. Our finding suggests that patients with low muscularity are likely to have low diaphragm thickness. Hence, preventing further diaphragm atrophy is an urgent matter in such patients. Diaphragm atrophy may be prevented by avoiding excessive ventilatory support and inflammation [32, 33]. Furthermore, diaphragm muscle training can preserve diaphragm muscle thickness [34]. The prevention and treatment of diaphragm atrophy are being investigated globally; however, few treatments exist [35]. Physical rehabilitation, such as mobilization, may contribute to treating diaphragm muscle atrophy as well as trunk muscle mass because of the associated relationship [36].

Based on the outcomes of the present study, we propose using ultrasound to assess muscularity at the time of ICU admission. The recognition of low muscularity at ICU admission is important for nutritional and rehabilitation intervention. Low muscularity at ICU admission harbors a risk of mortality and physical disability [37]. Early enteral nutrition and early mobilization are recommended in critically ill patients [38, 39], and we propose that personalized management based on muscularity could prevent further muscle loss. Given that we provided the cutoff value of ultrasound assessments, it may be possible to assess low muscularity using ultrasound. Additional studies are, however, needed to confirm that ultrasound muscle mass assessments can be used to improve patient management in ICU.

#### Limitations

First, this was a retrospective analysis of an observational study. Therefore, prospective studies should be conducted to validate these findings. Second, the reliable cutoff value of low muscularity is unclear for the entire ICU population. Therefore, we used a cutoff value for Japanese patients with cancer to avoid ethnicity differences. Third, it is still unclear whether muscle CSA at the L3 assessed by CT is related to whole-body muscularity in ICU population. Fourth, CT and ultrasound examinations were performed within 2 days before and after admission and not on the same day. However, these examinations were performed in the difference of 0 days (IQR, 0–1 days). Thus, temporal differences were not considered influential.

#### Conclusions

We retrospectively evaluated the relationship between ultrasound and CT muscle mass assessments and found that ultrasound measurements of the quadriceps muscle layer thickness and rectus femoris muscle CSA can serve as indicators of low muscularity. Furthermore, the elbow flexor muscle thickness, biceps brachii muscle CSA, and diaphragm thickness had weak-to-moderate correlation to trunk muscle mass. At ICU admission, the ultrasound assessment of muscle mass can be a promising method to identify low muscularity.

#### Acknowledgements

The authors are grateful for the cooperation and support of the ICU staff during the study. The authors thank all those who supported the Muscle Atrophy Zero Project, which aims to prevent muscle atrophy in patients.

#### Funding

This study was partly supported by a crowdfunding project entitled the Muscle Atrophy Zero Project using the platform "Otsucle" <a href="https://otsucle.jp/cf/project/2553.html">https://otsucle.jp/cf/project/2553.html</a>. This study was partially supported by JSPS KAKENHI Grant Numbers JP20K17899 and JP20K17899.

#### Competing interests

The authors declare that they have no competing interests

#### Authors' contributions

A.Y. participated in data acquisition and manuscript drafting. N.N. participated in study design, data acquisition, analysis, and manuscript drafting. The first two authors contributed equally to this study as first authors. Y.O. participated in manuscript revision. Profs. S.I., J.K., M.H., and J.O. supervised all aspects of this study. All authors read and approved the final manuscript.

References

[1] Weijs PJ, Looijaard WG, Dekker IM, Stapel SN, Girbes AR, Oudemans-van Straaten HM, et al. Low skeletal muscle area is a risk factor for mortality in mechanically ventilated critically ill patients. Crit Care. 2014;18:R12.

[2] Mitobe Y, Morishita S, Ohashi K, Sakai S, Uchiyama M, Abeywickrama H, et al. Skeletal muscle index at intensive care unit admission is a predictor of intensive care unit-acquired weakness in patients with sepsis. J Clin Med Res. 2019;11:834-41.

[3] Akan B. Influence of sarcopenia focused on critically ill patients. Acute Crit Care. 2021;36:15-21.

[4] Nakanishi N, Okura K, Okamura M, Nawata K, Shinohara A, Tanaka K, et al. Measuring and monitoring skeletal muscle mass after stroke: A review of current methods and clinical applications. J Stroke Cerebrovasc Dis. 2021.

[5] Chen LK, Woo J, Assantachai P, Auyeung TW, Chou MY, Iijima K, et al. Asian working group for sarcopenia: 2019 consensus update on sarcopenia diagnosis and treatment. J Am Med Dir Assoc. 2020;21:300-7.e2.

[6] Kim D, Sun JS, Lee YH, Lee JH, Hong J, Lee JM. Comparative assessment of skeletal muscle mass using computerized tomography and bioelectrical impedance analysis in critically ill patients. Clin Nutr. 2018. [7] Nakanishi N, Tsutsumi R, Okayama Y, Takashima T, Ueno Y, Itagaki T, et al. Monitoring of muscle mass in critically ill patients: Comparison of ultrasound and two bioelectrical impedance analysis devices. J Intensive Care. 2019;7:61.

[8] St-Onge MP, Wang Z, Horlick M, Wang J, Heymsfield SB. Dual-energy X-ray absorptiometry lean soft tissue hydration: Independent contributions of intra- and extracellular water. Am J Physiol Endocrinol Metab. 2004;287:E842-7.

[9] Dong V, Karvellas CJ. Using technology to assess nutritional status and optimize nutrition therapy in critically ill patients. Curr Opin Clin Nutr Metab Care. 2021;24:189-94.

[10] Okada Y, Kiguchi T, Okada A, Iizuka R, Iwami T, Ohtsuru S. Predictive value of sarcopenic findings in the psoas muscle on CT imaging among patients with sepsis. Am J Emerg Med. 2021;47:180-6.

[11] Weinel LM, Summers MJ, Chapple L-A. Ultrasonography to measure quadriceps muscle in critically ill patients: A literature review of reported methodologies. Anaesthesia and Intensive Care. 2019;47:423-34.

[12] Puthucheary ZA, McNelly AS, Rawal J, Connolly B, Sidhu PS, Rowlerson A, et al. Rectus femoris cross-sectional area and muscle layer thickness: comparative markers of muscle wasting and weakness. Am J Respir Crit Care Med. 2017;195:136-8.

[13] Palakshappa JA, Reilly JP, Schweickert WD, Anderson BJ, Khoury V, Shashaty MG, et al.

Quantitative peripheral muscle ultrasound in sepsis: Muscle area superior to thickness. J Crit Care. 2018;47:324-30.

[14] Nakanishi N, Oto J, Tsutsumi R, Akimoto Y, Nakano Y, Nishimura M. Upper limb muscle atrophy associated with in-hospital mortality and physical function impairments in mechanically ventilated critically ill adults: A two-center prospective observational study. J Intensive Care. 2020;8:87.

[15] Nakanishi N, Oto J, Ueno Y, Nakataki E, Itagaki T, Nishimura M. Change in diaphragm and intercostal muscle thickness in mechanically ventilated patients: A prospective observational ultrasonography study. J Intensive Care. 2019;7:56.

[16] Barotsis N, Tsiganos P, Kokkalis Z, Panayiotakis G, Panagiotopoulos E. Reliability of muscle thickness measurements in ultrasonography. Int J Rehabil Res. 2020;43:123-8.

[17] Perkisas S, Bastijns S, Baudry S, Bauer J, Beaudart C, Beckwée D, et al. Application of ultrasound for muscle assessment in sarcopenia: 2020 SARCUS update. European Geriatric Medicine. 2021;12:45-59.

[18] Tuinman PR, Jonkman AH, Dres M, Shi ZH, Goligher EC, Goffi A, et al. Respiratory muscle ultrasonography: methodology, basic and advanced principles and clinical applications in ICU and ED patients-a narrative review. Intensive Care Med. 2020;46:594-605.

[19] Nakanishi N, Inoue S, Tsutsumi R, Akimoto Y, Ono Y, Kotani J, et al. Rectus femoris

mimicking ultrasound phantom for muscle mass assessment: Design, research, and training application. Journal of Clinical Medicine. 2021;10:2721.

[20] Mourtzakis M, Prado CM, Lieffers JR, Reiman T, McCargar LJ, Baracos VE. A practical and precise approach to quantification of body composition in cancer patients using computed tomography images acquired during routine care. Appl Physiol Nutr Metab. 2008;33:997-1006.

[21] Gomez-Perez SL, Haus JM, Sheean P, Patel B, Mar W, Chaudhry V, et al. Measuring abdominal circumference and skeletal muscle from a single cross-sectional computed tomography image: A step-by-step guide for clinicians using national institutes of health imageJ. JPEN J Parenter Enteral Nutr. 2016;40:308-18.

[22] Su H, Ruan J, Chen T, Lin E, Shi L. CT-assessed sarcopenia is a predictive factor for both long-term and short-term outcomes in gastrointestinal oncology patients: A systematic review and meta-analysis. Cancer Imaging. 2019;19:82.

[23] Iritani S, Imai K, Takai K, Hanai T, Ideta T, Miyazaki T, et al. Skeletal muscle depletion is an independent prognostic factor for hepatocellular carcinoma. J Gastroenterol. 2015;50:323-32.
[24] Fetterplace K, Corlette L, Abdelhamid YA, Presneill JJ, Paris MT, Stella D, et al. Assessment of muscle mass using ultrasound with minimal versus maximal pressure compared with computed tomography in critically ill adult patients. Aust Crit Care. 2020.

[25] Paris MT, Mourtzakis M, Day A, Leung R, Watharkar S, Kozar R, et al. Validation of bedside

ultrasound of muscle layer thickness of the quadriceps in the critically ill patient (VALIDUM study). JPEN J Parenter Enteral Nutr. 2017;41:171-80.

[26] Hida T, Ando K, Kobayashi K, Ito K, Tsushima M, Kobayakawa T, et al. Ultrasound measurement of thigh muscle thickness for assessment of sarcopenia. Nagoya J Med Sci. 2018;80:519-27.

[27] Hacker ED, Peters T, Garkova M. Ultrasound assessment of the rectus femoris cross-sectional area: Subject position implications. West J Nurs Res. 2016;38:1221-30.

[28] Lambell KJ, Earthman CP, Tierney AC, Goh GS, Forsyth A, King SJ. How does muscularity assessed by bedside methods compare to computed tomography muscle area at intensive care unit admission? A pilot prospective cross-sectional study. J Hum Nutr Diet. 2020.

[29] Campbell IT, Watt T, Withers D, England R, Sukumar S, Keegan MA, et al. Muscle thickness, measured with ultrasound, may be an indicator of lean tissue wasting in multiple organ failure in the presence of edema. Am J Clin Nutr. 1995;62:533-9.

[30] Lambell KJ, Tierney AC, Wang JC, Nanjayya V, Forsyth A, Goh GS, et al. Comparison of ultrasound-derived muscle thickness with computed tomography muscle cross-sectional area on admission to the Intensive Care Unit: A pilot cross-sectional study. JPEN J Parenter Enteral Nutr. 2021;45:136-45.

[31] Sklar MC, Dres M, Fan E, Rubenfeld GD, Scales DC, Herridge MS, et al. Association of low

baseline diaphragm muscle mass with prolonged mechanical ventilation and mortality among critically ill adults. JAMA Netw Open. 2020;3:e1921520.

[32] Zambon M, Beccaria P, Matsuno J, Gemma M, Frati E, Colombo S, et al. Mechanical ventilation and diaphragmatic atrophy in critically ill patients: An ultrasound study. Crit Care Med. 2016;44:1347-52.

[33] Hadda V, Kumar R, Tiwari P, Mittal S, Kalaivani M, Madan K, et al. Decline in diaphragm thickness and clinical outcomes among patients with sepsis. Heart Lung. 2021;50:284-91.

[34] O'Rourke J, Soták M, Curley GF, Doolan A, Henlín T, Mullins G, et al. Initial assessment of the percutaneous electrical phrenic nerve stimulation system in patients on mechanical ventilation. Crit Care Med. 2020;48:e362-e70.

[35] Schepens T, Dres M, Heunks L, Goligher EC. Diaphragm-protective mechanical ventilation.Curr Opin Crit Care. 2019;25:77-85.

[36] Hickmann CE, Castanares-Zapatero D, Deldicque L, Van den Bergh P, Caty G, Robert A, et al. Impact of very early physical therapy during septic shock on skeletal muscle: A randomized controlled trial. Crit Care Med. 2018;46:1436-43.

[37] Jaitovich A, Dumas CL, Itty R, Chieng HC, Khan MMHS, Naqvi A, et al. ICU admission body composition: skeletal muscle, bone, and fat effects on mortality and disability at hospital discharge—A prospective, cohort study. Crit Care. 2020;24:566.

[38] Lang JK, Paykel MS, Haines KJ, Hodgson CL. Clinical practice guidelines for early mobilization in the ICU: A systematic review. Crit Care Med. 2020;48:e1121-e8.

[39] Reintam Blaser A, Starkopf J, Alhazzani W, Berger MM, Casaer MP, Deane AM, et al. Early enteral nutrition in critically ill patients: ESICM clinical practice guidelines. Intensive Care Med. 2017;43:380-98. Figure Legends

Figure 1. Flowchart of patients included in this study.

Among the 133 patients screened, 89 underwent computed tomography (CT) examinations at the level of the third lumbar vertebra. Secondary outcomes included the ultrasound assessments of the biceps brachii (n = 52) and diaphragm muscles (n = 79) (CT, computed tomography; L3, third lumbar vertebra).

Figure 2. Relationships between the ultrasound measurements of the quadriceps muscle layer thickness or rectus femoris muscle CSA and CT measurements of the cross-sectional area at the third lumbar vertebra.

(A) Quadriceps muscle layer thickness and (B) Rectus femoris muscle CSA. The Spearman correlation coefficient was used to investigate the relationships. CSA: cross-sectional area, CT: computed tomography

Figure 3. Areas under the receiver operating characteristic curves (AUCs) were estimated to determine the cutoff values of ultrasound assessments for low muscularity.

The Youden index was used to identify the optimal cutoff value. (A) Quadriceps muscle layer thickness and (B) Rectus femoris muscle CSA. (A) cutoff value was 2.0 cm at the sensitivity of 83.3% and the specificity of 78.5% and (B) cutoff value was 4.66 cm<sup>2</sup> at the sensitivity of 79.2% and the specificity of 66.2%. AUC: areas under the receiver operating characteristic curves, CSA: cross-sectional area, CI: confidence interval

Figure 4. Relationships with the ultrasound measurements of elbow flexor muscle thickness,

biceps brachii muscle CSA, sum of rectus femoris and biceps brachii muscle CSA, and diaphragm thickness.

(A) elbow flexor muscle thickness, (B) biceps brachii muscle CSA, (C) sum of rectus femoris and biceps brachii muscle CSA, (D) diaphragm thickness. CSA: cross-sectional area. Correlation between ultrasound and computed tomography measurements were evaluated. The Spearman correlation coefficient was used to investigate the relationships.













## Table 1. Patient characteristics

	All patients	Low muscularity*	Normal muscularity	
Variables	(n = 89)	(n = 24)	(n = 65)	p-value
Age, mean $\pm$ SD, y	$72 \pm 13$	$76 \pm 11$	$70\pm13.6$	0.07
Male/Female	60/29	16/8	44/21	0.93
Body mass index, mean $\pm$ SD, kg/m <sup>2</sup>	$22.2\pm4.4$	$18.6\pm3.0$	$23.5\pm4.0$	< 0.01
APACHE II score	27 (24–30)	28 (25–32)	27 (23–30)	0.31
SOFA	8 (6–11)	8 (6–11)	10 (6–12)	0.66
Sepsis (Sepsis-3 criteria), n (%)	40 (49)	24 (56)	16 (41)	0.27
Postoperative admissions, n (%)	14 (16)	1 (4)	13 (20)	0.07
Mechanical ventilation, n (%)	78 (88)	19 (79.2)	59 (90.8)	0.16
Length of ICU stay, d	7 (5–14)	6 (4–11)	7 (5–17)	0.13
Length of hospital stay, d	29 (18–51)	22 (15–44)	30 (21–51)	0.39
Mortality in the ICU, n (%)	16 (18.0)	4 (17)	12 (19)	0.85
Mortality in the hospital, n (%)	25 (28.1)	10 (41.7)	15 (23.1)	0.08
Ultrasound				
Quadriceps muscle layer thickness (cm)	2.4 (1.8–3.1)	1.7 (1.4–2.0)	2.8 (2.1–3.2)	< 0.01
Rectus femoris muscle CSA (cm <sup>2</sup> )	4.9 (3.9–6.5)	3.9 (2.9–4.6)	5.4 (4.1–7.1)	< 0.01
Elbow flexor muscle thickness (cm)†	2.8 (2.3–3.1)	2.3 (2.0-2.7)	3.2 (2.8–3.4)	< 0.01
Biceps brachii muscle CSA (cm <sup>2</sup> )†	6.1 (4.6-8.6)	5.1 (3.6–5.4)	6.8 (5.1–9.0)	< 0.01
Diaphragm thickness (mm);	1.8 (1.4–2.1)	1.7 (1.3–2.0)	1.8 (1.5–2.2)	0.28
Computed tomography				
CSA at 3 <sup>rd</sup> lumbar vertebra (cm <sup>2</sup> )	102.8 (77.1–133.2)	69.2 (54.1-80.5)	118.9 (97.0–143.5)	< 0.01

APACHE: Acute Physiology and Chronic Health Evaluation; SOFA: Sequential Organ Failure Assessment; SD: standard deviation; ICU:

intensive care unit; IQR: interquartile range; CSA: Cross-sectional area

Data were presented as median (IQR) unless otherwise indicated.

\* Low muscularity was defined as skeletal muscle index  $< 29.0 \text{ cm}^2/\text{m}^2$  for males and  $36.0 \text{ cm}^2/\text{m}^2$  for female.

<sup>†</sup> Included number of patients was 52 with 13 of low muscularity and 39 of normal muscularity.

‡ Included number of patients was 79 with 22 of low muscularity and 57 of normal muscularity.

## Supplemental File

## <u>Ultrasound assessment of muscle mass has potential to identify patients with low muscularity at</u> intensive care unit admission: A retrospective study

	Tokushima University Hospital	Tokushima Prefectural Central Hospital	
Ultrasounds	HI VISION Preirus, Hitachi	LOGIQ P9, GE healthcare, WI, USA	
	Medical Corporation, Tokyo, Japan		
Transducers	EUP-L73S liner transducer (4–9	12L-RS liner transducer (5–13 MHz),	
	MHz), Hitachi Medical	GE healthcare, WI, USA	
	Corporation, Tokyo, Japan		
Computed Tomography	Aquilion 16, Canon Medical	Philips Brilliance iCT, Philips	
	Systems, Tochigi, Japan	Healthcare, OH, USA	

## Table S1. Equipment used in this study

Figure S1 Ultrasound measurement images of limb thickness



A. Elbow flexor muscle thickness was measured from the superficial fascia of the biceps brachii muscle to the uppermost part of the humerus. B. Quadriceps muscle layer thickness was measured from the superficial fascia of the rectus femoris to the uppermost part of the femur.

Figure S2 Ultrasound measurement images of limb cross-sectional area



A. Biceps brachii muscle cross-sectional area was measured by tracking the muscle area shown in the transverse plane. B. Rectus femoris muscle ross-sectional area was measured by tracking the muscle area shown in the transverse plane.

Figure S3 Image of diaphragm thickness measurement



Diaphragm is observed as hypoechogenic muscular layer bordered by the echogenic layer of peritoneum and diaphragmatic pleurae.

### Figure S4 Image of a computed tomography measurement



The white line tracing includes the total muscle area in the third lumbar vertebra, which includes the psoas, quadratus lumborum, transversus abdominis, external and internal obliques, and rectus abdominis muscles.

	Correlation coefficient		Bland-Altman 95% CI	
Variables	r	р	Bias	95% CI
Intra-observer reproducibility				
Quadriceps muscle layer thickness	0.99	< 0.01	$-0.190 \pm 0.091$	-0.396 to 0.016
Rectus femoris muscle CSA	0.99	< 0.01	$0.113\pm0.067$	-0.039 to 0.265
Elbow flexor muscle thickness	0.99	< 0.01	$-0.230\pm0.126$	-0.514 to 0.054
Biceps brachii muscle CSA	0.96	< 0.01	$0.028\pm0.094$	-0.184 to 0.240
Diaphragm thickness	0.98	< 0.01	$0.045\pm0.036$	-0.030 to 0.120
Inter-observer reproducibility				
Quadriceps muscle layer thickness	0.99	< 0.01	$-0.290 \pm 0.307$	-0.985 to 0.405
Rectus femoris muscle CSA	0.99	< 0.01	$0.059\pm0.076$	-0.113 to 0.231
Elbow flexor muscle thickness	0.99	< 0.01	$0.020\pm0.251$	-0.548 to 0.588
Biceps brachii muscle CSA	0.99	< 0.01	$0.004\pm0.075$	-0.165 to 0.173
Diaphragm thickness	0.97	< 0.01	$0.040\pm0.027$	-0.016 to 0.096

## Table S2 Reproducibility of measurements

CI: confidence interval, CSA: cross-sectional area

Reproducibility was assessed for 10 patients in limb and for 20 patients in diaphragm. The Pearson correlation coefficient and Bland-Altman plot were determined by using JMP statistical software version 13.1.0 (SAS Institute Inc., Cary, NC, USA).

Variables	n median (IQR)		
Quadriceps muscle layer thickness (cm)			
All	133	2.4 (1.9–3.2)	
Male	85	2.6 (2.0-3.3)	
Female	48	2.1 (1.8–2.9)	
Young (< 70 years)	59	2.9 (2.1–3.5)	
Older ( $\geq$ 70 years)	74	2.1 (1.7–2.7)	
Rectus femoris muscle cross-sectional area (cm <sup>2</sup> )			
All	133	5.0 (3.9–7.0)	
Male	85	5.8 (4.1–7.9)	
Female	48	4.5 (3.7–5.7)	
Young (< 70 years)	59	6.5 (4.3-8.8)	
Older ( $\geq$ 70 years)	74	4.5 (3.5–5.6)	

**Table S3** The values of quadriceps muscle layer thickness and rectus femoris muscle crosssectional area in various populations

IQR: interquartile range

We presented the data of quadriceps muscle layer thickness and rectus femoris muscle crosssectional area in all 133 patients who had the ultrasound assessment at the ICU admission. The data was also shown in sex and age (young or older).