



# Suppression of the genes responsible for transporting hydrophobic pollutants leads to the production of safer crops

Fujita, Kentaro  
Haga, Yuki  
Yoshihara, Ryouhei  
Matsumura, Chisato  
Inui, Hideyuki

---

## (Citation)

Science of The Total Environment, 741:140439

## (Issue Date)

2020-11-01

## (Resource Type)

journal article

## (Version)

Accepted Manuscript

## (Rights)

© 2020 Elsevier B.V. All rights reserved.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>

## (URL)

<https://hdl.handle.net/20.500.14094/90009053>



# **Suppression of the genes responsible for transporting hydrophobic pollutants leads to the production of safer crops**

Kentaro Fujita<sup>1</sup>, Yuki Haga<sup>2</sup>, Ryouhei Yoshihara<sup>3, a</sup>, Chisato Matsumura<sup>2</sup>, Hideyuki Inui<sup>1,3,\*</sup>

<sup>1</sup>Graduate School of Agricultural Science, Kobe University, 1-1 Rokkodaicho, Nada-ku, Kobe, Hyogo 657-8501, Japan

<sup>2</sup>Hyogo Prefectural Institute of Environmental Sciences, 3-1-18 Yukihirocho, Suma-ku, Kobe, Hyogo 654-0037, Japan

<sup>3</sup>Biosignal Research Center, Kobe University, 1-1 Rokkodaicho, Nada-ku, Kobe, Hyogo 657-8501, Japan

<sup>a</sup>Present address: Faculty of Science, Saitama University, 255 Shimo-okubo, Sakura-ku, Saitama 338-8570, Japan.

\*Corresponding Author Hideyuki Inui, Biosignal Research Center, Kobe University, 1-1 Rokkodaicho, Nada-ku, Kobe, Hyogo, 657-8501, Japan

E-mail: hinui@kobe-u.ac.jp, Telephone number: +81-78-803-5863

## Abstract

Hydrophobic pollutants have become widely distributed across the world. From an agricultural perspective, their accumulation in crops from contaminated soil threatens food security and quality, leading to many diseases in humans. The Cucurbitaceae family can accumulate high concentrations of hydrophobic pollutants in their aerial parts. The Cucurbitaceae family contains major latex-like proteins (MLPs) as transporting factors for hydrophobic pollutants. *MLP* genes are expressed in the roots in which the MLPs bind hydrophobic pollutants. MLPs transport these hydrophobic pollutants to the aerial parts of the plant through the xylem vessels. As a result, hydrophobic pollutant contamination occurs in the Cucurbitaceae family. In this study, we suppressed the expression of *MLP* genes in the roots and reduced the amounts of MLPs with pesticide treatments. First, the fungicides Benlate and Daconil that decreased the hydrophobic pollutant, perylene, concentration in the xylem sap of zucchini plants were selected. Daconil suppressed the transcription activity of *MLP* in the roots. In the Daconil treatment, the amount of MLPs in the roots and xylem sap of zucchini plants was decreased, and the concentrations of the hydrophobic pollutants, pyrene and dieldrin, were significantly decreased. Our research contributes to the production of safer crops.

## Key words

*Cucurbita pepo*; hydrophobic pollutant; major latex-like protein; pesticide; gene suppression

## 1. Introduction

Some industrial materials, pesticides, and unintentional products have been recognized as hydrophobic pollutants, such as persistent organic pollutants (POPs) and polycyclic aromatic hydrocarbons. Since their toxicity was revealed, their use and production have been prohibited. However, hydrophobic pollutant contamination is still detected worldwide (Bartrons et al., 2016). Because of their high bioaccumulation ability, living organisms at the top of the food chain, such as human beings, are likely to accumulate high concentrations of hydrophobic pollutants. The intake of contaminated crops is a major route in the accumulation of hydrophobic pollutants. Agricultural soil can be contaminated by hydrophobic pollutants through biochar treatment (Fabbri et al., 2013), the incomplete combustion of straw (Jenkins et al., 1996), and POP-insecticide treatment, such as dieldrin, dichloro-diphenyl-trichloroethane, and  $\beta$ -hexachlorocyclohexanes (Namiki et al., 2013). The intake of hydrophobic pollutants by human beings occurs through the uptake of these pollutants by crops. The accumulation of hydrophobic pollutants in the human body leads to many diseases, such as respiratory syndromes (Bortey-Sam et al., 2017), digestive tract cancer (Abdur Rehman et al., 2017), and Alzheimer's (Yan et al., 2016). Therefore, it is critically important to inhibit the uptake of hydrophobic pollutants into crops at the first step of the food chain.

The Cucurbitaceae family, which includes cucumbers (*Cucumis sativus*), squashes (*Cucurbita maxima*), and zucchinis (*Cucurbita pepo*), can accumulate high concentrations of hydrophobic pollutants in their fruits (Parrish et al., 2006; Otani et al., 2007). It has been reported that the Cucurbitaceae family accumulates high concentrations of drins (dieldrin and endrin) (Otani et al., 2007), polychlorinated dibenzo-*p*-dioxins, polychlorinated dibenzofurans (Hülster et al., 1994; Inui et al., 2008), polychlorinated biphenyls (Inui et al., 2008), dichlorodiphenyldichloroethylene (Lunney et al., 2004), chlordane (Mattina et al., 2004), and pyrene (Parrish et al., 2006). Various approaches for the reduction of contamination through soil remediation were attempted to inhibit the accumulation of

73 hydrophobic pollutants in crops. For example, POPs are adsorbed by active carbon (Murano et al.,  
74 2009), and POP-degrading microorganisms were screened (Yamazaki et al., 2014). However, their  
75 POP removal efficiency was low. Therefore, we developed a new approach, using plant functions to  
76 produce low-contamination crops in contaminated soil.

77 It has been reported that major latex-like proteins (MLPs) play a crucial role in the accumulation of  
78 POPs in the Cucurbitaceae family (Inui et al., 2013). MLPs belong to Bet v 1 family (Fernandes et al.,  
79 2013) and are thought to function as resistance against pathogens (Yang et al., 2015) and as  
80 transporters of hydrophobic compounds using an internal hydrophobic cavity. *MLP* genes were highly  
81 expressed in the roots of the Cucurbitaceae family, and MLPs were detected in the roots and xylem  
82 sap (Goto et al., 2019). Furthermore, MLPs bind POPs in the root cells using an internal hydrophobic  
83 cavity (Goto et al., 2019). The resultant MLP-POP complexes are translocated into the xylem vessels,  
84 and the POPs are transported to the aerial parts of the plant (Goto et al., 2019). As a result,  
85 contamination by hydrophobic pollutants occurs. Therefore, the expression of *MLP* genes in the roots  
86 is a crucial step of contamination in the Cucurbitaceae family.

87 In this study, we focused on the expression of *MLP* genes in the roots. We attempt to develop a strategy  
88 for the cultivation of low-contamination crops, even in soil contaminated by hydrophobic pollutants.  
89 Previously, the amount of MLPs in zucchini plants was decreased by environmental factors, such as  
90 cultivation temperature and day length, leading to a decrease in pyrene concentration (Inui et al., 2020).  
91 Therefore, it is possible to reduce hydrophobic pollutant contamination in the Cucurbitaceae family  
92 by reducing the amount of MLPs. We attempted to suppress the expression of *MLP* genes in the roots  
93 using pesticide treatments in zucchini plants with the highest ability to accumulate the hydrophobic  
94 pollutants in the Cucurbitaceae family (Otani et al., 2007). Pesticides are essential products in  
95 agriculture and subject to safety tests. To accomplish our purpose, we selected the fungicide Daconil  
96 with the ability to suppress *MLP* gene expression and inhibit the accumulation of the hydrophobic

pollutants, pyrene and dieldrin, in the Cucurbitaceae family. This study enables farmers to produce safer crops in the contaminated farmland with the hydrophobic pollutants, although farmers have to apply the pesticide to reduce crop contamination even if they do not need it. The pesticide application is confirmed in safety and low cost compared with conventional manners such as the application of the active carbon and microorganisms.

## 2. Materials & Methods

### 2.1 Plant materials & pesticides

The seeds of the *C. pepo* subspecies *pepo* cultivars ‘Magda’ (MG) and ‘Raven’ (RA) were purchased from Johnny’s Selected Seeds (Albion, ME, USA). The insecticides Guardbait (Sankei Chemical Co., Ltd., Kagoshima, Japan), Starkle (Hokko Chemical Industry Co., Ltd., Tokyo, Japan), and Diazinon (Sankei Chemical Co., Ltd.) and the fungicides Benlate (Sumika Agrotech Co., Ltd., Osaka, Japan) and Daconil (Kumiai Chemical Industry Co., Ltd., Tokyo, Japan) were purchased from a market (Table S1).

### 2.2 Collection of xylem sap

The soil (Hyponex Japan Corp., Ltd., Osaka, Japan) was autoclaved for 15 min at 120 °C and completely dried. Then, 1 kg of dry soil was spiked with 500 mL of 2.5 mM perylene, 2.5 mM pyrene, or 25 µM dieldrin in acetone, and the acetone completely evaporated in a draft chamber. The seed coats of the MG and RA cultivar seeds were removed, and the seeds soaked in tap water overnight at 4 °C. Two or three seeds were sown in glass jars supplemented with 200 g of the contaminated soil. They were cultivated at 26 °C under a 16/8 h light/dark cycle. One healthy seedling was selected, and the others removed around 1 week after sowing. During cultivation, pesticides were applied to the soil with the recommended number of times according to the manufacturer's instructions, either at the

recommended dose or triple the recommended dose (Table S3 and S4). The powder of Guardbait, Starkle, Diazinon, and Benlate was applied, and Daconil was applied after 1,000 times dilution with tap water. After 27 days, xylem sap was collected, as previously reported (Inui et al., 2013). The stem was cut 1 cm below the cotyledon. A glass tube, washed with acetone or hexane, was set against the stem, and around 500  $\mu$ L of xylem sap was collected. The collected xylem sap was stored at 4 °C until use. After the collection of xylem sap, the roots were carefully washed with tap water to remove soil particles. The washed roots were stored at –80 °C until use.

### 2.3 Cloning of the promoter region of the MLP gene

The promoter region of the *MLP-GR3* gene was cloned by thermal asymmetric interlaced (TAIL) PCR (Liu et al., 1995). First, a 751-bp upstream region of the *MLP-GR3* gene was amplified with a random primer and iPCR-MLP-as1. The other two specific antisense primers, TAIL-MLP-P-as2 and TAIL-MLP-P-as3, were designed by referring 751-bp regions to amplify further upstream regions. The sequences of these primers were listed in Supplementary Table 2. Genomic DNA extracted from *C. pepo* ssp. *ovifera* cv. ‘Patty Green’ was used as the template, and the PCR reaction was performed with 2xQuick Taq HS DyeMix (Toyobo. Co., Ltd., Osaka, Japan). Although the reaction conditions of the TAIL-PCR followed the PCR conditions described by Liu and colleagues (Liu et al., 1995), the annealing temperature of the random hexamer and specific primers were changed to 33 °C and 53 to 56 °C, respectively.

A 1-month-old leaf of the MG cultivar was sampled, and genomic DNA was extracted using an Isoplant kit (Nippon Gene Co., Tokyo, Japan). The DNA fragment, including the promoter region of the *MLP* gene, was amplified by KOD FX Neo polymerase (Toyobo Co., Ltd.) with 100 ng of genomic DNA and the primers *MG-Pro-s* and *MG-Pro-as* under the following conditions: 5 min at 94 °C; 32 cycles of 30 s at 94 °C, 30 s at 56 °C, and 2 min at 72 °C; and 7 min at 72 °C (Table S2). The amplified

fragment was purified using the Gel/PCR DNA Isolation System (VIOGENE BIOTEK Co., New Taipei City, Taiwan). The 2,005-bp fragment was finally amplified and then inserted into the vector pBI221, digested with *Hind* III and *Xba* I using an In-Fusion HD Cloning Kit (Takara Bio Inc., Shiga, Japan). After sequencing, the fragment containing the *MLP* gene promoter and  $\beta$ -glucuronidase (*GUS*) gene cut by *Hind* III and *Sac* I was ligated with the plant expression vector pGWB402 $\Omega$  (provided by Dr. Tsuyoshi Nakagawa at Shimane University), digested by both restriction enzymes (Nakagawa et al., 2007).

#### 2.4 Transformation of tobacco plants (*Nicotiana tabacum*)

The plant expression plasmid containing the *MLP* gene promoter and *GUS* gene was introduced into the *Rhizobium radiobacter* strain LBA4404, and the leaf disk method was employed to produce transgenic tobacco lines because tobacco plants are commonly employed for GUS activity assay (Horsch et al., 1985). The T<sub>4</sub> generation of the transgenic tobacco plants was used for the GUS assay under the pesticide treatments.

#### 2.5 GUS assay

The seeds of transgenic tobacco plants containing the *pMLP-GR3::GUS* fusion were sown on Murashige and Skoog medium and incubated for 1 day at 4 °C in the dark. Then, they were incubated at 24 °C under a 16-h light/8-h dark cycle. After two weeks, the seedlings were transferred to water and incubated at 24 °C for another day. They were transferred to water containing 0.22  $\mu$ M, 0.43  $\mu$ M, or 0.58  $\mu$ M benomyl (Wako Pure Chemical Industries, Ltd., Osaka, Japan), an active compound in Benlate, or 0.019  $\mu$ M, 0.038  $\mu$ M, or 0.11  $\mu$ M tetrachloroisophthalonitrile (TPN) (Wako Pure Chemical Industries, Ltd.), an active compound in Daconil, with 1% dimethyl sulfoxide (DMSO), and cultivated at 24 °C under a 16-h light/8-h dark cycle. The concentration gradients for pesticides were selected



because the concentrations of benomyl and TPN influencing physiological changes in transgenic tobacco plants were unknown. After two days, the roots of each plant were collected, frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$  until use. Root samples in a protein extraction buffer [50 mM sodium phosphate buffer (pH 7.0), 10 mM EDTA (pH 8.0), 0.1% (v/v) Triton X-100, 0.1% (w/v) *N*-lauroylsarcosine sodium salt, and 0.072% (v/v) 2-mercaptoethanol] were homogenized. After centrifugation at  $4^{\circ}\text{C}$  at  $20,700 \times g$  for 5 min, the supernatants were collected and centrifuged again under the same conditions. The supernatants were subjected to GUS assay, as reported previously (Kodama et al., 2007).

## 2.6 Western blot analysis

Roots from the MG cultivar were ground in liquid nitrogen, and an extraction buffer [50 mM sodium phosphate buffer (pH 7.0), 10 mM EDTA, 0.1% (v/v) Triton X-100, 0.1% (v/v) *N*-lauroylsarcosine sodium salt] supplemented with 10 mM 2-mercaptoethanol was added to the root powder. The suspension was vigorously mixed, and the supernatant was collected after centrifugation at  $20,700 \times g$  for 15 min at  $4^{\circ}\text{C}$ , as reported previously (Goto et al., 2019). The concentrations of proteins in the xylem sap and of the extracted root proteins were quantified using the Bradford method (Bradford, 1976). Sample buffer solution with reducing reagent for SDS-PAGE (Nacalai Tesque, Inc., Kyoto, Japan) was added to the samples, and they were subjected to SDS-PAGE on 15% acrylamide gel. Western blot analysis was performed with anti-MLP-PG1 and anti-MLP-GR3 antibodies, as reported previously (Goto et al., 2019). Briefly, polyvinylidene difluoride membranes with proteins transferred from gels were reacted with the anti-MLP-PG1 and anti-MLP-GR3 antibodies (Medical & Biological Laboratories Co., Ltd., Aichi, Japan) in TTBS buffer [20 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.05% (v/v) Tween 20]. After the reaction with alkaline phosphatase-conjugated goat anti-rabbit IgG

(Sigma-Aldrich, St. Louis, MO) in TTBS buffer, MLPs were detected. The band intensities were quantified using ImageJ software (Schneider et al., 2012).

## 2.7 Quantification of perylene, pyrene, and dieldrin

The obtained xylem sap (50  $\mu$ L) and 50  $\mu$ L of DMSO were mixed in a 96-well black microtiter plate. The mixture was scanned using a microplate reader (Microplate Reader SH-9000, Corona Electric Co., Ltd., Hitachinaka, Ibaraki, Japan). The fluorescences of perylene and pyrene were measured at a 410 nm excitation wavelength and 445 nm emission wavelength, and a 330 nm excitation wavelength and at 390 nm emission wavelength, respectively.

The xylem sap (250  $\mu$ L) was mixed with 20  $\mu$ L of 200 ng/mL  $^{13}$ C-labeled dieldrin (Cambridge Isotope Laboratories, Inc., Tewksbury, MA, USA), offered by the Tohoku Ryokka Kankyohozen Co., Ltd. (Sendai, Japan) as an internal standard. Dieldrin was extracted twice using 1 mL of hexane. After dehydration with sodium sulfate dehydrate, 50  $\mu$ L of nonane and 10  $\mu$ L of 50 ng/mL MBP-15, 70, 101, 153 (Wellington Laboratories Inc., Guelph, Canada) as a syringe spike was added, and the samples were concentrated using nitrogen gas at 35  $^{\circ}$ C until the total amount reached approximately 100  $\mu$ L. The concentration of dieldrin was quantified by high-resolution gas chromatography and high-resolution mass spectrometry system [HRGC/HRMS: GC, 6890 N (Agilent Technologies, Tokyo, Japan); MS, JMS-800D (JEOL Ltd., Tokyo, Japan)] equipped with a DB-5MS column (Kanto Chemical Co. Inc., Tokyo, Japan).

## 2.8 Statistical analysis

One-way analysis of variance was performed, and Dunnett's multiple comparison test was applied to judge significant differences.

### 3. Results

#### 3.1 Selection of pesticides to decrease perylene concentration in the xylem sap

The insecticides Guardbait, Starkle, and Diazinon and the fungicides Benlate and Daconil were selected as applicable pesticides for Cucurbitaceae family plants. The fresh weights of the aerial parts of the plants did not significantly differ between treatments after 27 days of incubation (Figure S1A). The perylene concentration in the xylem sap was not significantly lower in any of the tested pesticide treatments, but seemed likely to decrease in the Benlate and Daconil treatments (Figure 1A). Furthermore, the perylene concentrations of the plants in the triple-dose pesticide treatments were also investigated. The fresh weight of the aerial parts did not differ between treatments (Figure S1B). Perylene concentrations were significantly decreased in the triple-dose Diazinon, Benlate, and Daconil treatments (Figure 1B). Benlate and Daconil were selected as pesticides with the ability to suppress the expression of *MLP* genes.

#### 3.2 Decrease in the transcription activity of the MLP gene

GUS activity in the roots treated with benomyl did not significantly decrease, but tended towards decreasing after treatment with 0.43  $\mu$ M benomyl (Figure 2). In contrast, GUS activity was significantly decreased, by 45% and 36%, after the treatment with 0.019  $\mu$ M and 0.11  $\mu$ M TPN, respectively (Figure 2). Thus, Daconil was selected as an *MLP* gene-suppressing pesticide.

#### 3.3 Decrease of pyrene concentration in the xylem sap

The MG cultivar was cultivated in soil contaminated with pyrene and treated with Daconil. In the Daconil treatment with the recommended dose, the pyrene concentration in the xylem sap was significantly decreased by 36% (Figure 3). There were no significant differences in the fresh weights of the aerial parts between treatments (Figure S2).

### 3.4 Decrease in the amount of MLPs in the roots and xylem sap

In the roots, a band of over 17 kDa was detected after the reaction with the anti-MLP-PG1 antibody. In the Daconil treatment, the relative band intensity of MLP-PG1 was decreased by 84% (Figure 4A). Two bands of over 17 kDa were detected in the roots after the reaction with the anti-MLP-GR3 antibody. In the Daconil treatment, both band intensities were decreased, and the relative band intensity of the lower band of MLP-GR3 was decreased by 93% (Figure 4A). Thus, the amount of MLP-PG1 and MLP-GR3 in the roots was decreased in the Daconil treatment. Two bands of over 17 kDa were detected in the xylem sap after the reaction with the anti-MLP-PG1 antibody. In the Daconil treatment, both bands disappeared (Figure 4B). A band below 17 kDa was detected in the xylem sap after the reaction with the anti-MLP-GR3 antibody. In the Daconil treatment, the relative band intensity of MLP-GR3 was decreased by 18% (Figure 4B). Thus, the amount of MLP-PG1 and MLP-GR3 in the xylem sap was also decreased in the Daconil treatment (Figure 4B).

### 3.5 Decrease of dieldrin concentration in xylem sap

The RA cultivar was cultivated in soil contaminated with dieldrin and treated with Daconil. In the Daconil treatment with the triple dose, the dieldrin concentration was significantly decreased by 52% (Figure 5). In the Daconil treatment with the recommended dose, there was no significant decrease, but the value was 13% lower (Figure 5). There were no significant differences in the fresh weights of the aerial parts between treatments (Figure S3).

## 4. Discussion

The purpose of the present study was to reduce crop contamination by hydrophobic pollutants through the suppression of the expression of *MLP* genes. The binding of POPs by MLPs in the roots is a crucial

step for the POP contamination in the Cucurbitaceae family (Inui et al., 2013; Goto et al., 2019).

Therefore, suppression of the expression of *MLP* genes in the roots leads to a reduction of contamination. First, we focused on identifying pesticides that suppress *MLP* gene expression. Pesticides are widely used for crop cultivation and cause physiological changes in plants. This study proposes the novel use of pesticides to reduce contamination. Thus, the pesticides are not utilized for their insecticidal and fungicidal actions.

In this study, pesticides with the ability to suppress *MLP* gene expression were selected from five pesticides that can be used on the Cucurbitaceae family. Diazinon, Benlate, and Daconil caused a significant decrease in perylene concentration in the xylem sap in the triple-dose treatments. Benlate and Daconil caused a decrease of perylene in the xylem sap even with the recommended dose. Thus, these pesticides were used for further analysis (Figure 1A). Pesticides contain active compounds for insecticidal and fungicidal action, spreading agents, and detergent. Transgenic tobacco plants, containing the *GUS* gene downstream of the promoter of the *MLP-GR3* gene, were treated with active compounds to confirm that an active compound suppresses *MLP* gene expression. TPN, an active compound in Daconil, significantly decreased GUS activity in the roots. The results clearly show that TPN suppressed *MLP* gene transcription in the roots. We did not clarify the mechanisms underlying the suppression of *MLP* gene expression. The detailed mechanisms underlying this gene suppression should be clarified, as it could lead to the production of safer crops.

There are several reports that pesticides changed the expression level of genes in plants (Lu et al., 2016). TPN up-regulated more than 500 genes, including *cytochrome P450* and *glutathione S-transferase*, and down-regulated more than 400 genes in the leaves of tomato plants (Zhou et al., 2015). Some pesticides induce the expression of genes involved in resistance against pathogens. For example, the pesticides probenazole and benzothiadiazole induced the genes for UDP: glucose salicylic acid glucosyltransferase and WRKY45, respectively, in rice (Shimono et al., 2007; Umemura et al., 2009).

Therefore, it is possible that *MLP* gene expression was suppressed by the pesticide treatments. MLPs have been identified in many plants, such as *Arabidopsis thaliana* (Lytle et al., 2009), soybeans (Strömvik et al., 1999), cotton (Chen and Dai, 2010), grapes (Zhang et al., 2018), and mulberries (Gai et al., 2018). Thus, the expression of *MLP* in various plant species could probably be controlled using pesticide treatments.

Our results clearly show that a decrease in the amount of MLP-PG1 in the roots led to a decrease in the amount of MLP-PG1 in the xylem sap translocated from the roots. However, the amount of MLP-GR3 in the xylem sap was only slightly decreased, although that of the MLP-GR3 in the roots was remarkably decreased. One possible reason for this is that the stability of MLP-GR3 is relatively high in xylem sap (the accumulation of MLP-GR3 was found even in the early growth stage of *C. pepo*), whereas the expression level of the *MLP-GR3* gene is low (Goto et al., 2019). It is thought that the decrease of two MLPs in the xylem sap additively produced a significant decrease in the concentrations of pyrene and dieldrin. Since xylem vessels are the main pathway connecting the roots and aerial parts of the plant, such as the leaves and fruits, the concentration of these pollutants in the fruits would also be decreased.

## 5. Conclusion

Based on these results, in the fungicide Daconil treatment, the transcription of *MLP* in the roots was suppressed, and the amount of MLPs in the roots decreased. Subsequently, the amount of MLPs translocated into the xylem vessels from the roots also decreased. As a result, the concentration of hydrophobic pollutants in the xylem sap of zucchini plants was significantly decreased, leading to a decrease in pollutant concentrations in the aerial parts of the plant. We are now trying to clarify the mechanisms underlying the suppression of *MLP* in the fungicide Daconil treatment. This is the first report to reduce crop contamination through the suppression of *MLP* expression using pesticide

treatment. This study proposes a new usage of pesticides. Since it is easy and low-cost, this approach is applicable and could potentially replace the remediation of soil by biological, chemical, and physical methods. It is notable that farmers can potentially cultivate crops in contaminated soil and produce safe crops using pesticide treatments.

### **Conflicts of interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### **Acknowledgments**

We thank Dr. Tsuyoshi Nakagawa of Shimane University and Tohoku Ryokka Kankyohozen Co., Ltd. (Sendai, Japan) for kindly providing us the plant expression plasmid pGWB402Ω and <sup>13</sup>C-labeled dieldrin, respectively.

### **Funding**

This work was, in part, supported by a grant from the Japan Crop Protection Association in 2016 and a Sasakawa Scientific Research Grant from The Japan Science Society (No. 2019-5004) to KF.

### **References**

- Abdur Rehman, M.Y., Kamal, A., Taqi, M.M., Malik, R.N., 2017. Tracing biomarker of PAH-exposure and susceptibility factor (GSTM-polymorphism) among cancer patients in Pakistan. *Chemosphere* 178, 384–390. <https://doi.org/10.1016/j.chemosphere.2017.03.017>
- Bartrons, M., Catalan, J., Penuelas, J., 2016. Spatial And Temporal Trends Of Organic Pollutants In Vegetation From Remote And Rural Areas. *Sci. Rep.* 6, 1–10. <https://doi.org/10.1038/srep25446>

336 Bortey-Sam, N., Ikenaka, Y., Akoto, O., Nakayama, S.M.M., Asante, K.A., Baidoo, E., Obirikorang, C.,  
 337 Saengtienchai, A., Isoda, N., Nimako, C., Mizukawa, H., Ishizuka, M., 2017. Oxidative stress and  
 338 respiratory symptoms due to human exposure to polycyclic aromatic hydrocarbons (PAHs) in  
 339 Kumasi, Ghana. *Environ. Pollut.* 228, 311–320. <https://doi.org/10.1016/j.envpol.2017.05.036>  
 340 Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of  
 341 protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72, 248–254.  
 342 [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)  
 343 Chen, J.-Y., Dai, X.-F., 2010. Cloning and characterization of the *Gossypium hirsutum* major latex  
 344 protein gene and functional analysis in *Arabidopsis thaliana*. *Planta* 231, 861–73.  
 345 <https://doi.org/10.1007/s00425-009-1092-2>  
 346 Fabbri, D., Rombolà, A.G., Torri, C., Spokas, K.A., 2013. Determination of polycyclic aromatic  
 347 hydrocarbons in biochar and biochar amended soil. *J. Anal. Appl. Pyrolysis* 103, 60–67.  
 348 <https://doi.org/10.1016/j.jaap.2012.10.003>  
 349 Fernandes, H., Michalska, K., Sikorski, M., Jaskolski, M., 2013. Structural and functional aspects of PR-  
 350 10 proteins. *FEBS J.* 280, 1169–1199. <https://doi.org/10.1111/febs.12114>  
 351 Gai, Y.P., Yuan, S.S., Liu, Z.Y., Zhao, H.N., Liu, Q., Qin, R.L., Fang, L.J., Ji, X.L., 2018. Integrated  
 352 phloem sap mRNA and protein expression analysis reveals phytoplasma-infection responses in  
 353 mulberry. *Mol. Cell. Proteomics* 17, 1702–1719. <https://doi.org/10.1074/mcp.RA118.000670>  
 354 Goto, J., Iwabuchi, A., Yoshihara, R., Kodama, N., Matsui, T., Hirota, M., Eun, H., Inui, H., 2019.  
 355 Uptake mechanisms of polychlorinated biphenyls in *Cucurbita pepo* via xylem sap containing  
 356 major latex-like proteins. *Environ. Exp. Bot.* 162, 399–405.  
 357 <https://doi.org/10.1016/j.envexpbot.2019.03.019>



358 Horsch, R.B., Fry, J.E., Hoffmann, N.L., Eichholtz, D., Rogers, S.G., Fraley, R.T., 1985. A Simple and  
 359 General Method for Transferring Genes into Plants. *Science* (80-. ). 227, 1229–1231.  
 360 <https://doi.org/10.1210/jcem-10-10-1361>  
 361 Hülster, A., Müller, J.F., Marschner, H., 1994. Soil–Plant Transfer of Polychlorinated Dibenzo-*p*-dioxins  
 362 and Dibenzofurans to Vegetables of the Cucumber Family (*Cucurbitaceae*). *Environ. Sci. Technol.*  
 363 28, 1110–1115. <https://doi.org/10.1021/es00055a021>  
 364 Inui, H., Katte, N., Goto, J., Iwabuchi, A., 2020. High temperatures promote the uptake of hydrophobic  
 365 pollutants by *Cucurbita pepo* via altered gene expression levels of major latex-like proteins. *J.*  
 366 *Pestic. Sci.* 45, 75–80. <https://doi.org/10.1584/jpestics.d19-065>  
 367 Inui, H., Sawada, M., Goto, J., Yamazaki, K., Kodama, N., Tsuruta, H., Eun, H., 2013. A Major Latex-  
 368 Like Protein Is a Key Factor in Crop Contamination by Persistent Organic Pollutants. *Plant Physiol.*  
 369 161, 2128–2135. <https://doi.org/10.1104/pp.112.213645>  
 370 Inui, H., Wakai, T., Gion, K., Kim, Y.S., Eun, H., 2008. Differential uptake for dioxin-like compounds by  
 371 zucchini subspecies. *Chemosphere* 73, 1602–1607.  
 372 <https://doi.org/10.1016/j.chemosphere.2008.08.013>  
 373 Jenkins, B.M., Jones, A.D., Turn, S.Q., Williams, R.B., 1996. Emission factors for polycyclic aromatic  
 374 hydrocarbons from biomass burning. *Environ. Sci. Technol.* 30, 2462–2469.  
 375 <https://doi.org/10.1021/es950699m>  
 376 Kodama, S., Okada, K., Inui, H., Ohkawa, H., 2007. Aryl hydrocarbon receptor (AhR)-mediated reporter  
 377 gene expression systems in transgenic tobacco plants. *Planta* 227, 37–45.  
 378 <https://doi.org/10.1007/s00425-007-0592-1>  
 379 Liu, Y. -G, Mitsukawa, N., Oosumi, T., Whittier, R.F., 1995. Efficient isolation and mapping of  
 380 *Arabidopsis thaliana* T-DNA insert junctions by thermal asymmetric interlaced PCR. *Plant J.*  
 381 <https://doi.org/10.1046/j.1365-313X.1995.08030457.x>

382 Lu, Y.C., Feng, S.J., Zhang, J.J., Luo, F., Zhang, S., Yang, H., 2016. Genome-wide identification of DNA  
 383 methylation provides insights into the association of gene expression in rice exposed to pesticide  
 384 atrazine. *Sci. Rep.* 6, 1–15. <https://doi.org/10.1038/srep18985>  
 385 Lunney, A.I., Zeeb, B.A., Reimer, K.J., 2004. Uptake of weathered DDT in vascular plants: Potential for  
 386 phytoremediation. *Environ. Sci. Technol.* 38, 6147–6154. <https://doi.org/10.1021/es030705b>  
 387 Lytle, B.L., Song, J., De La Cruz, N.B., Peterson, F.C., Johnson, K.A., Bingman, C.A., Phillips, G.N.,  
 388 Volkman, B.F., 2009. Structures of two *Arabidopsis thaliana* major latex proteins represent novel  
 389 helix-grip folds. *Proteins Struct. Funct. Bioinforma.* 76, 237–243.  
 390 <https://doi.org/10.1002/prot.22396>  
 391 Mattina, M.I., Eitzer, B.D., Iannucci-Berger, W., Lee, W.Y., White, J.C., 2004. Plant uptake and  
 392 translocation of highly weathered, soil-bound technical chlordane residues: Data from field and  
 393 rhizotron studies. *Environ. Toxicol. Chem.* 23, 2756–2762. <https://doi.org/10.1897/03-570>  
 394 Murano, H., Otani, T., Makino, T., Seike, N., 2009. Effects of the application of carbonaceous adsorbents  
 395 on pumpkin (*Cucurbita maxima*) uptake of heptachlor epoxide in soil. *Soil Sci. Plant Nutr.* 55, 325–  
 396 332. <https://doi.org/10.1111/j.1747-0765.2009.00361.x>  
 397 Nakagawa, T., Suzuki, T., Murata, S., Nakamura, S., Hino, T., Maeo, K., Tabata, R., Kawai, T., Tanaka,  
 398 K., Niwa, Y., Watanabe, Y., Nakamura, K., Kimura, T., Ishiguro, S., 2007. Improved gateway  
 399 binary vectors: High-performance vectors for creation of fusion constructs in transgenic analysis of  
 400 plants. *Biosci. Biotechnol. Biochem.* 71, 2095–2100. <https://doi.org/10.1271/bbb.70216>  
 401 Namiki, S., Otani, T., Seike, N., 2013. Fate and plant uptake of persistent organic pollutants in soil. *Soil*  
 402 *Sci. Plant Nutr.* 59, 669–679. <https://doi.org/10.1080/00380768.2013.813833>  
 403 Otani, T., Seike, N., Sakata, Y., 2007. Differential uptake of dieldrin and endrin from soil by several plant  
 404 families and *Cucurbita* genera. *Soil Sci. Plant Nutr.* 53, 86–94. [https://doi.org/10.1111/j.1747-](https://doi.org/10.1111/j.1747-0765.2007.00102.x)  
 405 [0765.2007.00102.x](https://doi.org/10.1111/j.1747-0765.2007.00102.x)

406 Parrish, Z.D., White, J.C., Isleyen, M., Gent, M.P.N., Iannucci-Berger, W., Eitzer, B.D., Kelsey, J.W.,  
 407 Mattina, M.I., 2006. Accumulation of weathered polycyclic aromatic hydrocarbons (PAHs) by plant  
 408 and earthworm species. *Chemosphere* 64, 609–618.  
 409 <https://doi.org/10.1016/j.chemosphere.2005.11.003>  
 410 Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image analysis.  
 411 *Nat. Methods* 9, 671–675. <https://doi.org/10.1038/nmeth.2089>  
 412 Shimono, M., Sugano, S., Nakayama, A., Jiang, C.J., Ono, K., Toki, S., Takatsuji, H., 2007. Rice  
 413 WRKY45 plays a crucial role in benzothiadiazole-inducible blast resistance. *Plant Cell* 19, 2064–  
 414 2076. <https://doi.org/10.1105/tpc.106.046250>  
 415 Strömvik, M. V., Sundararaman, V.P., Vodkin, L.O., 1999. A novel promoter from soybean that is active  
 416 in a complex developmental pattern with and without its proximal 650 base pairs. *Plant Mol. Biol.*  
 417 41, 217–231.  
 418 Umemura, K., Satou, J., Iwata, M., Uozumi, N., Koga, J., Kawano, T., Koshiba, T., Anzai, H., Mitomi,  
 419 M., 2009. Contribution of salicylic acid glucosyltransferase, OsSGT1, to chemically induced  
 420 disease resistance in rice plants. *Plant J.* 57, 463–472. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-313X.2008.03697.x)  
 421 313X.2008.03697.x  
 422 Yamazaki, K., Takagi, K., Kataoka, R., Kotake, M., Yamada, T., Kiyota, H., 2014. Novel  
 423 phosphorylation of aldrin-trans-diol by dieldrin-degrading fungus *Mucor racemosus* strain DDF.  
 424 *Int. Biodeterior. Biodegrad.* 92, 36–40. <https://doi.org/10.1016/j.ibiod.2014.04.005>  
 425 Yan, D., Zhang, Y., Liu, L., Yan, H., 2016. Pesticide exposure and risk of Alzheimer’s disease: A  
 426 systematic review and meta-analysis. *Sci. Rep.* 6, 1–9. <https://doi.org/10.1038/srep32222>  
 427 Yang, C.L., Liang, S., Wang, H.Y., Han, L.B., Wang, F.X., Cheng, H.Q., Wu, X.M., Qu, Z.L., Wu, J.H.,  
 428 Xia, G.X., 2015. Cotton major latex protein 28 functions as a positive regulator of the ethylene

responsive factor 6 in defense against *Verticillium dahliae*. Mol. Plant 8, 399–411.

<https://doi.org/10.1016/j.molp.2014.11.023>

Zhang, N., Li, R., Shen, W., Jiao, S., Zhang, J., Xu, W., 2018. Genome-wide evolutionary

characterization and expression analyses of major latex protein (MLP) family genes in *Vitis*

*vinifera*. Mol. Genet. Genomics 293, 1061–1075. <https://doi.org/10.1007/s00438-018-1440-7>

Zhou, Y., Xia, X., Yu, G., Wang, J., Wu, J., Wang, M., Yang, Y., Shi, K., Yu, Y., Chen, Z., Gan, J., Yu,

J., 2015. Brassinosteroids play a critical role in the regulation of pesticide metabolism in crop

plants. Sci. Rep. 5, 1–7. <https://doi.org/10.1038/srep09018>

## Figure legends

Figure 1. Perylene concentration in xylem sap from *Cucurbita pepo* 'Magda' (MG) after treatment with pesticides at the recommended (A) and triple the recommended (B) dose.

The MG cultivar was cultivated in soil contaminated with perylene (1.25 mmol/kg) under a 16/8 h light/dark cycle at 26 °C. During cultivation, pesticides were applied under various dosage conditions (Table S3). After 27 days, xylem sap was collected from each plant, and the perylene concentration quantified by measuring the fluorescence at 410 nm excitation and 445 nm emission. –, cultivation in soil not contaminated with perylene or no pesticide treatment; +, cultivation in soil contaminated with perylene; GB, SK, DZ, BL, and DN, Guardbait, Starkle, Diazinon, Benlate, and Daconil treatment, respectively. Error bars indicate standard deviations ( $n = 3-4$ ). Asterisks indicate significant differences compared to perylene concentration in samples cultivated in soil contaminated with perylene without pesticide treatment (\*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ).

Figure 2. Suppression of reporter gene expression in the roots of transgenic tobacco plants after treatment with the active compounds of pesticides.

Two-week-old transgenic tobacco plants expressing the  $\beta$ -glucuronidase (*GUS*) gene downstream of the *MLP-GR3* promoter were incubated in water containing 0.22  $\mu$ M (+), 0.43  $\mu$ M (++), or 0.58  $\mu$ M (+++) benomyl or 0.019  $\mu$ M (+), 0.038  $\mu$ M (++), or 0.11  $\mu$ M (+++) TPN under a 16/8 h light/dark cycle at 24 °C for 2 days. The proteins from the roots were extracted, and GUS assay was performed.

Error bars indicate standard deviation ( $n = 4-5$ ). Asterisks indicate significant differences compared to GUS activity in samples incubated in water without pesticides (\*,  $p < 0.05$ ).

Figure 3. Pyrene concentration in the xylem sap of *Cucurbita pepo* 'Magda' (MG) after Daconil treatment.

The MG cultivar was cultivated in soil contaminated with pyrene (1.25 mmol/kg) under a 16/8 h light/dark cycle at 26 °C. The fungicide Daconil was applied at the recommended dose. During cultivation, pesticides were applied under various dosage conditions (Table S4). After 27 days, xylem sap was collected from each plant, and the pyrene concentration was quantified by measuring the fluorescence at 330 nm excitation and 390 nm emission. –, cultivation in soil not contaminated with pyrene or no Daconil treatment; +, cultivation in soil contaminated with pyrene or treatment of Daconil at the recommended dose. Error bars indicate standard deviation ( $n = 7-8$ ). Asterisks indicate significant differences compared to the pyrene concentration of samples cultivated in soil contaminated with pyrene without Daconil treatment. (\*\*,  $p < 0.01$ ).

Figure 4. Major latex-like proteins in the roots (A) and xylem sap (B) of *Cucurbita pepo* 'Magda' (MG) after Daconil treatment.

The MG cultivar was cultivated in soil contaminated with pyrene (1.25 mmol/kg) under a 16/8 h light/dark cycle at 26 °C. The fungicide Daconil was applied at the recommended dose. During cultivation, pesticides were applied under various dosage conditions (Table S4). After 27 days, the roots and xylem sap were collected, and the proteins in the roots were extracted. Root extracts and xylem sap were subjected to SDS-PAGE. MLP-PG1 and MLP-GR3 were detected by western blot analysis using the anti-MLP-PG1 antibody and anti-MLP-GR3 antibody, respectively. –, no Daconil treatment; +, Daconil treatment. Band intensities were quantified by ImageJ.

Figure 5. Dieldrin concentration in the xylem sap of *Cucurbita pepo* ‘Raven’ (RA) after Daconil treatment.

The RA cultivar was cultivated in soil contaminated with dieldrin (12.5  $\mu\text{mol/kg}$ ) under a 16/8 h light/dark cycle at 26 °C. The fungicide Daconil was applied at the recommended dose and triple the recommended dose. During cultivation, pesticides were applied under various dosage conditions (Table S4). After 27 days, xylem sap was collected, and dieldrin concentration was quantified by high-resolution gas chromatography and high-resolution mass spectrometry. –, cultivation in soil not contaminated with dieldrin or no Daconil treatment; +, cultivation in soil contaminated with dieldrin or Daconil treatment at the recommended dose; +++, Daconil treatment at triple the recommended dose. Error bars indicate standard deviation ( $n = 5-8$ ). Asterisks indicate significant differences compared to the dieldrin concentration of samples cultivated in soil contaminated with dieldrin with no Daconil treatment (\*\*\*,  $p < 0.001$ ).

Supplementary Figure 1. Fresh weight of the aerial parts of *Cucurbita pepo* ‘Magda’ (MG) after pesticide treatment at recommended (A) and triple (B) doses.

The MG cultivar was cultivated in soil contaminated with perylene (1.25 mmol/kg) under a 16/8 h light/dark cycle at 26 °C. During cultivation, pesticides were applied under various dosage conditions (Table S3). After 27 days, the fresh weights of the aerial parts were measured. –, cultivation in soil not contaminated with perylene or no pesticide treatment; +, cultivation in soil contaminated with perylene; GB, SK, DZ, BL, and DN, Guardbait, Starkle, Diazinon, Benlate, and Daconil treatment, respectively. Error bars indicate standard deviation ( $n = 3-4$ ).

Supplementary Figure 2. Fresh weight of the aerial parts of *Cucurbita pepo* ‘Magda’ (MG) after Daconil treatment.

The MG cultivar was cultivated in soil contaminated with pyrene (1.25 mmol/kg) under a 16/8 h light/dark cycle at 26 °C. The fungicide Daconil was applied at the recommended dose. During cultivation, pesticides were applied under various dosage conditions (Table S4). After 27 days, the fresh weights of the aerial parts were measured. –, cultivation in soil not contaminated with pyrene or no Daconil treatment; +, cultivation in soil contaminated with pyrene or Daconil treatment at the recommended dose. Error bars indicate standard deviation ( $n = 7-8$ ).

Supplementary Figure 3. Fresh weight of the aerial parts of *Cucurbita pepo* ‘Raven’ (RA) after Daconil treatment.


The RA cultivar was cultivated in soil contaminated with dieldrin (12.5 µmol/kg) under a 16/8 h light/dark cycle at 26 °C. The fungicide Daconil was applied at the recommended dose and triple the recommended dose. During cultivation, pesticides were applied under various dosage conditions (Table S4). After 27 days, the fresh weights of the aerial parts were measured. –, cultivation in soil not contaminated with dieldrin or no Daconil treatment; +, cultivation in soil contaminated with dieldrin or Daconil treatment at the recommended dose; +++, Daconil treatment at triple the recommended dose. Error bars indicate standard deviation ( $n = 5-8$ ).



**Pesticides**



***MLP* genes**



**Toxic**




**Zucchini**  
**(*Cucurbita pepo*)**

**Safe**



(A)



(B)




Figure 1




Figure 2





Figure 3

**(A)**



**(B)**




Figure 4






Figure 5


(A)



(B)




Supplementary Figure 1



Supplementary Figure 2





Supplementary Figure 3

Supplementary Table 1. Physicochemical and toxic properties of pesticides.

Property	Pesticide				
	GB	SK	DZ	BL	DN
$\log K_{ow}$	6.5	-0.549	3.81	2.12	3.05
LD <sub>50</sub> (mg/kg)	430 - 4000	2000 - 2800	1340	>9590	>10000

The value for  $\log K_{ow}$  and LD<sub>50</sub> of GB, DZ, BL, and DN were from web service (<https://pubchem.ncbi.nlm.nih.gov/>). Those of SK were from papers (Corbel et al., 2004; Li et al., 2018).

Supplementary Table 2. Sequences of primers used in this study.

Primer name	Sequence
MG-Pro-s	5'-TGATTACGCCAAGCTTAAGCATTCAATAAGTTGTT-3'
MG-Pro-as	5'-CCGGGGATCCTCTAGTTTCTTTCGATGTGATACAA-3'
iPCR-MLP-as1	5'-CCTAAAATCTTCTCCAGAGA-3'
TAIL-MLP-P-as2	5'-ATCTTGTCTCCCTCTCCAAC-3'
TAIL-MLP-P-as3	5'-CTTCGTGAACTAAATGGGGGC-3'

Supplementary Table 3. Applied amount of pesticides at the recommended dose during the cultivation of *Cucurbita pepo* ‘Magda’ (MG).

Cultivation period (day)	Pesticide				
	GB	SK	DZ	BL	DN
10	9 mg	0.45 mg	24 mg	300 mg	9 µL
13	9 mg	—	—	300 mg	9 µL
17	9 mg	—	—	300 mg	—
20	9 mg	—	—	—	—
24	9 mg	—	—	—	—

The MG cultivar was cultivated in soil contaminated with perylene (1.25 mmol/kg) under a 16/8 h light/dark cycle at 26°C. After 27 days, xylem sap was collected from each plant. The triple dose of the pesticides was applied at three times the amount described in the table. —, no pesticide treatment; GB, SK, DZ, BL, and DN, Guardbait, Starkle, Diazinon, Benlate, and Daconil treatment, respectively. The powder of GB, SK, DZ, and BL was applied, and DN was applied after 1,000 times dilution in tap water.

Supplementary Table 4. Applied amount of Daconil during the cultivation of *Cucurbita pepo* ‘Magda’ (MG) and ‘Raven’ (RA) at the recommended dose.

Cultivation period (day)	Cultivar	
	MG	RA
11	9 µL	–
12	–	9 µL
14	9 µL	–
15	–	9 µL
20	–	9 µL

The MG and RA cultivars were cultivated in soil contaminated with pyrene (1.25 mmol/kg) and dieldrin (12.5 µmol/kg) under a 16/8 h light/dark cycle at 26°C. After 27 days, xylem sap was collected from each plant. The triple dose of pesticides was applied at three times the amount described in the table. –, no pesticide treatment. DN was applied after 1,000 times dilution in tap water.

## References

Corbel, V., Duchon, S., Zaim, M., Hougard, J.-M., 2004. Dinotefuran: A Potential Neonicotinoid

Insecticide Against Resistant Mosquitoes. *J. Med. Entomol.* 41, 712–717.

<https://doi.org/10.1603/0022-2585-41.4.712>

Li, Y., Long, L., Yan, H., Ge, J., Cheng, J., Ren, L., Yu, X., 2018. Comparison of uptake, translocation

and accumulation of several neonicotinoids in komatsuna (*Brassica rapa* var. *perviridis*) from

contaminated soils. *Chemosphere* 200, 603–611.

<https://doi.org/10.1016/j.chemosphere.2018.02.104>