



# Almond Skin Polyphenol Extracts Stimulate the Activation of Diacylglycerol Kinase alpha via a 67 kDa Laminin Receptor

Chinthammit, Boonyaporn  
Okamoto, Seika  
Shirai, Yasuhito

---

**(Citation)**

Functional Foods in Health and Disease, 12(4):151-160

**(Issue Date)**

2022-04

**(Resource Type)**

journal article

**(Version)**

Version of Record

**(Rights)**

©FFC 2022.

This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

**(URL)**

<https://hdl.handle.net/20.500.14094/90009259>





## Almond Skin Polyphenol Extracts Stimulate the Activation of Diacylglycerol Kinase alpha via a 67 kDa Laminin Receptor

Boonyaporn Chinthammit<sup>1</sup>, Seika Okamoto<sup>1</sup>, Yasuhito Shirai<sup>1\*</sup>

<sup>1</sup> Department of Agrobioscience, Graduate School of Agriculture, Kobe University, Japan

**\*Corresponding Author:** Yasuhito Shirai, Ph.D. Department of Agrobioscience, Graduate School of Agriculture, Kobe University, Japan

**Submission Date:** January 19<sup>th</sup>, 2022; **Acceptance Date:** March 18<sup>th</sup>, 2022; **Publication Date:** April 1<sup>st</sup>, 2022

**Please cite this article as:** Chinthammit B., Okamoto S., Shirai Y. Almond Skin Polyphenol Extracts Stimulate the Activation of Diacylglycerol Kinase alpha via a 67 kDa Laminin Receptor. *Functional Foods in Health and Disease* 2022; 12(4):151-160. DOI: <https://www.doi.org/10.31989/ffhd.v12i3.894>

### ABSTRACT


**Background:** Almond skins are the byproduct of the almond industry that are rich in dietary fibers and polyphenols. Recently, there has been an increasing interest in utilizing the phenolic compounds from almond skins for their functional benefits. Galloylated catechins activate diacylglycerol kinase  $\alpha$  (DGK $\alpha$ ) which is involved in the amelioration of diabetic nephropathy. Therefore, in this study, we investigated whether almond skin polyphenol extracts can also induce the activation of DGK $\alpha$ .

**Methods:** Phenolic contents in the almond skin polyphenol extracts were identified by liquid chromatography (LC)-time of flight mass spectrometry (TOFMS). Using confocal microscopy, the translocation of green fluorescent protein (GFP)-DGK $\alpha$  to the cell membrane was observed upon stimulation with almond skin polyphenol extracts. To check the involvement of 67 kDa laminin receptor (67LR), pre-treatment of anti-67LR antibody was used.

**Results:** We identified that naringenin and flavanone (2,3-dihydroflavone), which are among the phenolic contents in the almond skin polyphenol extracts identified, can also induce the activation of DGK $\alpha$ . In addition, we also investigated whether the pathway involves the same receptor as that of epigallocatechin-3-gallate (EGCg); the 67LR. Naringenin stimulated through the 67LR, while flavanone mainly used the 67LR-independent pathways.

**Conclusion:** These findings were additional function of almond skin polyphenol extracts and may implicate the benefits of the intake of nuts in daily diets. Polyphenols can be extracted from almond skins as an inexpensive source. The various health benefits of polyphenols can be applied to functional foods and supplements.

**Keywords:** flavonoid, lipid kinase, imaging, catechin, vitamin E, diabetic nephropathy



©FFC 2022. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

## INTRODUCTION

Diabetic nephropathy is one of the major vascular complications that affects approximately 35% of patients with diabetes mellitus and is the leading cause of end-stage kidney disease [1]. The pathogenesis of the disease involves complex interplay between both genetic and environmental factors, leading to the stimulation of inflammatory responses and abnormal renal functions [2]. Hyperglycemia has been extensively studied as the major cause of glomerular hyperfiltration, and one of the underlying mechanisms is the protein Kinase C (PKC)-dependent pathways. PKC has diverse functions ranging from controlling the polyol pathway to the production of advanced glycation end products. It is activated by

diacylglycerol (DAG) which increases during hyperglycemia through many pathways such as, de novo synthesis and conversion from phosphatidic acid (PA). The excess activation of PKC by DAG may enhance the activity of angiotensin II and the production of extracellular matrix (ECM), contributing towards diabetic nephropathy [3]. Recently, the amelioration of diabetic nephropathy has been focusing on reducing the activity of PKC through inhibition [4] or reducing the amount of its activator, DAG, by DAG kinase (DGK) [5].

DGK phosphorylates DAG forming PA and thus, attenuates the activity of PKC [6]. Upon activation, DGK translocates from the cytoplasm to the plasma membrane. Among the 5 subtypes of mammalian DGK

expressed in the glomeruli, DGK $\alpha$  has been shown to be stimulated by d- $\alpha$ -tocopherol [7] and the oral administration of d- $\alpha$ -tocopherol significantly reduced the loss of podocytes, which make slits for renal filtration [8]. The chroman ring of d- $\alpha$ -tocopherol is required for the activation and translocation of DGK $\alpha$  to the plasma membrane [9]. Subsequently, galloylated catechins have also been confirmed to activate DGK $\alpha$ , especially epigallocatechin-3-gallate (EGCg) which has been shown to induce the activation through the cell surface receptor 67-kDa laminin receptor (67LR) [10]. 67LR is the receptor for EGCg, the predominant green tea polyphenol, and is responsible for the anticancer activities of EGCg such as, cell growth inhibition and apoptosis [11]. The requirement for 67LR by galloylated catechins to activate DGK $\alpha$  may be the underlying mechanism for the biological effects of galloylated catechins. However, our understanding of the interactions between DGK $\alpha$  and 67LR is limited.

Almonds are rich in monounsaturated fats, proteins, and dietary fibers. Recently, along with almonds' highly nutritious profile, almond skin polyphenols have been extensively characterized as the source of antioxidants and bioactives. The polyphenols identified in almonds include proanthocyanidins, hydrolysable tannins and flavonoids [12]. Blanched almond skins are the by-product of the food processing industry which have a potential to be used as a functional ingredient, due to their high dietary fibers and polyphenol composition. A growing number of evidence supports the health benefits of almond consumption in improving metabolic syndrome [13]. Almond consumption has been shown to improve postprandial glycemia and hormonal control of patients with type 2 diabetes [14]. However, less is known about the molecular mechanism underlying these health benefits. Therefore, in this study, we investigated whether almond skin polyphenols stimulate the activation of DGK $\alpha$  via a 67LR.

## MATERIALS AND METHODS

**2.1 Almond skin polyphenol extraction:** Almond skin polyphenol crude (ASP-A), 30% (ASP-B) and 50% (ASP-C) extracts were obtained following the same extraction but different purification procedures. Blanched almond skin (provided by Ezaki Glico Co., Ltd., Osaka, Japan) was dried at 60°C until the moisture was < 5%. Extraction was conducted by refluxing 1:10 (w/v) skin in 70% (v/v) aqueous ethanol for 1 hr at 80°C. The extract was filtrated through No. 5C filter paper (Advantech MFS, CA, USA) to remove the skin. For ASP-A, ethanol was evaporated, and the extract was passed through Whatman Grade 2 qualitative filter paper (GE Healthcare, NJ, USA) in the presence of 20% Radiolite 300 diatomite filter aid (Showa Chemical Industry, Tokyo, Japan).

For ASP-B and ASP-C, the extract was passed through Whatman Grade 2 qualitative filter paper with 1% Radiolite 300 diatomite filter aid. Then, ethanol was removed, and the filtrate was centrifuged at 17,000  $\times$  g for 15 min at 4°C to remove insoluble particles. Column separation was performed using Diaion HP20 column (Mitsubishi Chemical, Tokyo, Japan). The column was washed with 10% (v/v) aqueous ethanol and eluted with 30% and 50% (v/v) aqueous ethanol to obtain ASP-B and ASP-C, respectively. The extracts were freeze-dried for storage.

**2.2 Total phenolic content quantification:** The total phenolic content was determined based on Folin-Ciocalteu method [15]. Stock solutions of almond skin polyphenol extracts were prepared at 10 mg/ml in methanol and diluted with Milli-Q water. 100  $\mu$ l sample was mixed with 500  $\mu$ l 10% Folin-Ciocalteu reagent (Nacalai tesque, Kyoto, Japan) for 3 min, then 400  $\mu$ l 7.5% (w/v) aqueous sodium carbonate was added. The mixture was incubated in the dark for 1 hr, and absorbance was measured at 765 nm. The total phenolic contents were calculated in terms of Gallic Acid Equivalents (GAE).

**2.3 LC-TOFMS analysis:** Approximately 5 mg of almond skin polyphenol extract was mixed with 300  $\mu$ l 1% (v/v) formic acid in acetonitrile containing internal standard solution (H3304-1002, Human Metabolome Technologies Inc., Tsuruoka, Japan). The mixture was homogenized three times at 1,500 rpm for 120 s using a tissue homogenizer (Micro Smash MS100R, Tomy Digital Biology, Tokyo, Japan). The mixture was homogenized again after adding 100  $\mu$ l Milli-Q water, followed by centrifugation at  $2,300 \times g$  for 5 min at 4°C. The supernatant was filtrated through 3-kDa cut-off filter (Nanocap 3K Omega, Pall Corporation, Michigan, USA) to remove proteins, and further filtrated by using Hybrid SPE phospholipid 55261-U (Supelco, PA, USA) to remove phospholipids. The filtrate was desiccated and then dissolved in 100  $\mu$ l 50% (v/v) aqueous isopropanol prior to analysis. LC-TOFMS was carried out through a facility service (Human Metabolome Technologies Inc., Tsuruoka, Japan).

**2.4 GFP-DGK $\alpha$  translocation assay:** Smooth muscle cells, DDT1-MF2, were cultured in high glucose DMEM (Wako, Osaka, Japan) supplemented with 2 mM L-glutamine, 5% fetal bovine serum (FBS) and 1% antibiotics (100 U/ml penicillin and 100  $\mu$ g/ml streptomycin). At sub-confluence, cells were sub-cultured to 35 mm poly-lysine coated glass bottom dish (Matsunami, Osaka, Japan) at  $1.0 \times 10^5$  cells/dish and incubated for 24 hr at 37°C. Then, lipofection was conducted following the protocol of FuGENE HD transfection reagent (Promega, WI, USA) with the plasmid encoding GFP-DGK $\alpha$  [9]. Stock solutions of almond skin polyphenol extracts, Epicatechin gallate (ECg) (Wako, Osaka, Japan), EGCg (LKT laboratories Inc., MN, USA), naringenin and flavanone (Sigma Aldrich, MO, USA) were prepared in methanol and diluted with Ringer's solution composed of Krebs-Ringer phosphate buffer (120 mM NaCl, 4.8 mM KCl, 1 mM CaCl<sub>2</sub>, 1.2 mM MgSO<sub>4</sub>, 16.5 mM NaH<sub>2</sub>PO<sub>4</sub>/Na<sub>2</sub>HPO<sub>4</sub>, pH 7.4, and 5 mM  $\beta$ -D-glucose). Mixture sample contained ECg, flavanone


and naringenin in the ratio of 16:740:8.1, respectively. 67LR blocking was performed prior to sample treatment where cells were treated with 20  $\mu$ g/ml anti-67LR antibody (clone name MLC5) (GeneTex, CA, USA) for 2 hr at 37°C. Transfected cells were treated with various concentrations of samples for 5 min at 37°C and fixed with 4% PFA. Images were taken with confocal fluorescence microscope (FV-1000, Olympus, PA, USA). Translocation rate was calculated as the average percentage of the number of cells where GFP-DGK $\alpha$  translocation was observed to the total number of GFP-DGK $\alpha$  expressing cells.

**2.5 Statistical analysis:** All values are expressed as mean (SD). Statistical analysis was performed using R 3.6.1 (R Core Team, 2019). The data were analyzed using one-way analysis of variance (ANOVA) with a post-hoc Tukey HSD test. \* and \*\* means  $p < 0.05$  and  $p < 0.01$ , respectively.


## RESULTS AND DISCUSSION

### 3.1 Effects of almond skin polyphenol extracts on DGK $\alpha$

**localization:** The extract contained mainly polyphenols and the effects of almond polyphenols on DGK $\alpha$  activation were investigated through the translocation of GFP-DGK $\alpha$  to the plasma membrane. After stimulation with different almond skin polyphenol extracts, GFP-DGK $\alpha$  was observed to aggregate as spots on the cell membrane, like the pattern of EGCg (Fig. 1A). At 50  $\mu$ g/ml, both ASP-B and ASP-C showed higher translocation rates than the EGCg control (Fig. 1B). Fig. 2 shows the total phenolic contents of respective almond skin polyphenol extracts, indicating that the total phenolic contents of almond skin polyphenol extract remarkably increased after purification. Corresponding to the amount of total phenolic contents, ASP-A induced the lowest translocation rate compared to the other fractions.



**Fig 1.** Effects of almond skin polyphenol extracts on GFP-DGK $\alpha$  localization. (A) Typical images of GFP-DGK $\alpha$  after stimulation with almond skin polyphenol extracts. (B) Comparison of the translocation rates of GFP-DGK $\alpha$  induced by different almond skin polyphenol extracts, compared to EGCg positive control, at 50  $\mu$ g/ml. (\*\*  $p < 0.01$ )




**Fig 2.** Total phenolic contents of almond skin polyphenol extracts, expressed as gallic acid equivalents (GAE) per g extract (dry weight). (\*\*  $p < 0.01$ )

**3.2 Identification of DGK $\alpha$  stimulating flavanones in almond skin polyphenol extracts:** To identify the bioactive compounds that induced DGK $\alpha$  activation, ASP-B and ASP-C were subjected to polyphenol composition analysis by LC-TOFMS (Fig. 3). 34 phenolic compounds

were identified; 26 of which were flavonoids. Interestingly, EGCg was not detected in the almond skin polyphenol extracts. ASP-B contained 15 flavonoids, while ASP-C contained 23 flavonoids. The structural basis behind the activation of DGK $\alpha$  by catechins requires both

the chroman ring and the gallate moieties [10]. However, DGK $\alpha$  activation was still observed for ASP-C which lacked galloylated catechins. Therefore, other active compounds should be present in ASP-C that induced the activation of DGK $\alpha$ . It has been reported that only the chroman ring was sufficient for the action of d- $\alpha$ -tocopherol [9]. There were 14 unmodified flavonoids identified in the almond skin polyphenol extracts; 4 of

which belonged to the flavanone subgroup and only flavanone (2,3-dihydroflavone) and naringenin were detected in both extracts. ASP-C contained higher flavanone than ASP-B of  $1.3 \times 10^{-1}$  and  $7.4 \times 10^{-2}$  relative concentration units, respectively. Naringenin was also higher in ASP-C than ASP-B of  $1.7 \times 10^{-2}$  and  $8.1 \times 10^{-4}$  relative concentration units, respectively. Thus, we focused on the flavanone and the naringenin.




**Fig 3.** Classification of the phenolic compounds found in almond skin polyphenol extracts by LC-TOFMS. Comparisons of the relative concentrations of unmodified flavonoids in ASP-B (filled) and ASP-C (unfilled) are also shown



Upon stimulation with flavanone and naringenin, GFP-DGK $\alpha$  was observed to accumulate around the cell membrane (Fig. 4A). The dose-dependent effect of flavanone was comparable to that of ECg (Fig. 4Bi). On the other hand, naringenin showed a moderate effect of just approximately 26% translocation rate at 50  $\mu$ g/ml. Although the results showed a variable effectiveness of different flavanones, they suggest that, at least, flavanone and naringenin are bioactive compounds contained in almond skins to activate DGK $\alpha$ . This was supported by the observation that the mixture sample, containing the same ratio of ECg: flavanone: naringenin as the ASP-B, was effective in inducing the translocation

of DGK $\alpha$  (Fig. 4Bii).

Interestingly, the dose-dependent effect of the mixture sample was lower than that of ASP-B. The difference indicates the presence of other DGK $\alpha$  activating phenolic compounds in the almond skin polyphenol extracts, apart from ECg, flavanone and naringenin. Further experiments are required to analyze the phenolic contents of the almond skin polyphenol extracts to identify other DGK $\alpha$  activating compounds. In addition, we had better check if the other small components contained in the extract can activate DGK $\alpha$  or not.




**Fig 4.** Effects of different phenolic compounds and almond skin polyphenol extracts on GFP-DGK $\alpha$  localization. (A) Typical images of GFP-DGK $\alpha$  after stimulated with different flavanones and a mixture sample, containing the same ratio of ECg:Naringenin:Flavanone as ASP-B. ECg was used as a positive control. (B) Dose-dependent effects on GFP-DGK $\alpha$  translocation induced by (i) flavanones and ECg; (ii) almond skin polyphenol extracts and a mixture sample.



**3.3 Involvement of 67LR in the activation of DGK $\alpha$ :** 67LR is involved in the activation of DGK $\alpha$  by both EGCg and  $\alpha$ -tocopherol. To assess whether 67LR is also required for the activation of DGK $\alpha$  by flavanones, we investigated the effect of the anti-67LR antibody on the translocation. The extent of the anti-67LR antibody inhibition on GFP-DGK $\alpha$  activation varied depending on the extracts' phenolic compositions (Fig. 5). Blocking the 67LR with antibody significantly attenuated the translocation of GFP-DGK $\alpha$  induced by naringenin and ECg, by approximately 27% and 83%, respectively. This shows that naringenin DGK $\alpha$  activates via 67LR. On the other hand, the flavanone-induced translocation showed the tendency to be inhibited by the presence of anti-67LR antibody, yet the difference was not statistically significant. This highlights the presence of 67LR-dependent and 67LR-independent pathways utilized by different phenolic compounds in activating DGK $\alpha$ . Due to the lipophilic structures of the flavanones, they can readily enter the cell through passive diffusion and bind

to cytoplasmic receptors to induce signal transductions. Flavanone has been shown to bind to peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) [16]. Therefore, it may be possible that flavanones activate DGK $\alpha$  via an intracellular receptor, rather than binding to the cell surface 67LR. However, the different side chains of the flavanones influence their hydrophobicity and affect the reactivity and the mechanism of action. There is some evidence supporting the binding of flavanones to cell surface receptors, such as the human  $\beta$ 2-adrenergic receptor [17] and FMS-like tyrosine kinase 3 (FLT3) [18]. Although evidence for naringenin binding to cell surface receptors is limited, the presence of hydroxyl side chains on naringenin reduces its lipophilicity, and thus it may be possible that naringenin preferentially binds to the cell surface 67LR. The low DGK $\alpha$  activation effect of naringenin may highlight the low affinity interaction with the 67LR. The binding of naringenin to 67LR will need to be further elucidated and whether the binding occurs at the same motif as EGCg [11]



**Fig5.** Effects of pretreatment with the 67LR antibody on the translocation of GFP-DGK $\alpha$  upon stimulation with different compounds. (A) ECg; (B) Flavanone; (C) Naringenin; (\*  $p < 0.05$ , \*\*  $p < 0.01$ )

The activation of DGK $\alpha$  may be related to alterations to the cell membrane through lipid raft modifications. D- $\alpha$ -tocopherol has been reported to associate with lipid rafts, modulating the membrane lipid composition and influencing signal transductions [19]. 67LR has also been identified to associate with lipid rafts which affected the

activity of EGCg [20]. Previously, the activities of DGK $\alpha$  and DGK $\zeta$  have been shown to be modulated by membrane properties and compositions [21]. Therefore, compounds that can alter the properties of the cell membrane are likely to affect the activation of DGK $\alpha$ .

Recently, focus has been on utilizing the beneficial

almond skin polyphenols for their antioxidant capacity [22], anti-inflammation [23] and anti-viral effects [24]. Here, we demonstrated that almond skin polyphenols stimulate the activation of DGK $\alpha$  via a 67LR. DGK $\alpha$  has also been reported to be involved in insulin secretion by pancreatic  $\beta$  cells [25]. 67LR has been identified as cell surface receptor, and there has been reported several signal pathways through this receptor. Thus, there are potentials in almond polyphenols in activating DGK $\alpha$  or signaling pathway through 67LR. For example, there is a potential in ameliorating diabetes through the synergistic effects of both improving the kidney function and regulating insulin secretion, although the effect will have to be further validated through clinical trials involving patients with diabetic nephropathy. The findings in this study provide additional functions of almond skin polyphenol extracts and may implicate the benefits of the intake of nuts in daily diets.

## CONCLUSION

Almond skin polyphenol extracts induced the activation of DGK $\alpha$ . The effect was dependent on the phenolic acid compositions of the extracts. ASP-B contained fewer types of flavonoids than ASP-C, yet its effect was stronger. Among the phenolic compounds identified in the almond skin polyphenol extracts, flavanone and naringenin were also found to activate DGK $\alpha$ . The 67LR was required for the induction of DGK $\alpha$  activation by ECg, and naringenin. However, flavanone stimulated DGK $\alpha$  via a 67LR-independent pathway.

**List of Abbreviations:** ASP: Almond skin polyphenol crude; DAG: diacylglycerol; DGK: diacylglycerol kinase, ECg: epicatechin gallate; EGCg: epigallocatechin-3-gallate; ECM: extracellular matrix; FBS: fetal bovine serum; GAE: gallic acid equivalents; GFP: green fluorescent protein; LC: liquid chromatograph; MS : mass

spectrometry; PKC: Protein Kinase C; 67LR: 67 kDa laminin receptor; PA: phosphatidic acid; PPAR $\gamma$ : peroxisome proliferator-activated receptor  $\gamma$

**Author Contributions:** B. Chinthammit and S. Okamoto designed and conducted the experiments. B. Chinthammit interpreted the results and drafted the manuscript. Y. Shirai provided supervision throughout this work.

**Conflicts of Interest:** The authors declare no conflict of interest.

**Acknowledgements and Funding:** This work was supported and funded by joint research with Ezaki Glico Co., Ltd. Specially, we thank Dr. Hiroko Takumi and Dr. Hiroshi Kamasaka at Ezaki Glico Co., Ltd. for their constructive advice.

## REFERENCES

1. Khoury CC, Chen S, Ziyadeh FN: Pathophysiology of diabetic nephropathy. In Chronic Renal Disease. Edited by Paul L. Kimmel and Mark E. Rosenberg. 2015:151-162. <https://doi.org/10.1016/B978-0-12-411602-3.00013-5>
2. Navarro-González J, Mora-Fernández C, de Fuentes M, García-Pérez J: Inflammatory molecules and pathways in the pathogenesis of diabetic nephropathy. Nat Rev Nephrol 2011, 7:327-340. <https://doi.org/10.1038/nrneph.2011.51>
3. Lz J, Gobe G: Protein kinase C activation and its role in kidney disease. Nephrology 2006, 11:428-434. <https://doi.org/10.1111/j.1440-1797.2006.00673.x>
4. Kelly DJ, Zhang Y, Hepper C, Gow RM, Jaworski K, Kemp BE, Wilkinson-Berka JL, Gilbert RE: Protein kinase C beta inhibition attenuates the progression of experimental diabetic nephropathy in the presence of continued hypertension. Diabetes 2003, 52:512-518. <https://doi.org/10.2337/diabetes.52.2.512>
5. Hayashi D, Ueda S, Yamanoue M, Ashida H, Shirai Y: Amelioration of diabetic nephropathy by oral administration of d- $\alpha$ -tocopherol and its mechanisms. Biosci Biotechnol Biochem 2018, 82: 65-73. <https://doi.org/10.1080/09168451.2017.1411184>
6. Mérida I, Arranz-Nicolás J, Rodríguez-Rodríguez C, Ávila-Flores A: Diacylglycerol kinase control of protein kinase C.

- Biochem J 2019, 476:1205–1219.  
<https://doi.org/10.1042/BCJ20180620>
7. Hayashi D, Shirai Y: Protective role of alpha-tocopherol in diabetic nephropathy. In Nutritional and Therapeutic Interventions for Diabetes and Metabolic Syndrome. 2nd edition. Edited by Debasis Bagchi and Sreejayan Nair: 2018:375-383. <https://doi.org/10.1016/B978-0-12-812019-4.00030-1>
  8. Hayashi D, Yagi K, Song C, Ueda S, Yamanoue M, Topham M, Suzuki T, Saito N, Emoto N, Shirai Y: Diacylglycerol kinase alpha is involved in the vitamin E-induced amelioration of diabetic nephropathy in mice. Sci Rep 2017, 7:2597. <https://doi.org/10.1038/s41598-017-02354-3>
  9. Fukunaga-Takenaka R, Shirai Y, Yagi K, Adachi N, Sakai N, Merino E, Merida I, Saito N: Importance of chroman ring and tyrosine phosphorylation in the subtype-specific translocation and activation of diacylglycerol kinase  $\alpha$  by d- $\alpha$ -tocopherol. Genes Cells 2005, 10: 311-319. <https://doi.org/10.1111/j.1365-2443.2005.00842>
  10. Hayashi D, Ueda S, Yamanoue M, Saito N, Ashida H, Shirai Y: Epigallocatechin-3-gallate activates diacylglycerol kinase alpha via a 67 kDa laminin receptor: A possibility of galloylated catechins as functional food to prevent and/or improve diabetic renal dysfunctions. J Funct Foods 2015, 15:561-569. <https://dx.doi.org/10.1016/j.jff.2015.04.005>
  11. Fujimura Y, Sumida M, Sugihara K, Tsukamoto S, Yamada K, Tachibana H: Green tea polyphenol EGCG sensing motif on the 67-kDa laminin receptor. PLoS ONE 2012, 7: e37942. <https://doi.org/10.1371/journal.pone.0037942>
  12. Bolling B: Almond polyphenols: methods of analysis, contribution to food quality, and health promotion. Compr Rev Food Sci Food Saf 2017, 16: 346-368. <https://doi.org/10.1111/1541-4337.12260>
  13. Kamil A, Chen CO: Health benefits of almonds beyond cholesterol reduction. J Agric Food Chem 2012, 60: 6694-6702. <https://doi.org/10.1021/jf2044795>
  14. Bodnaruc A, Prud'homme D, Giroux I: Acute effects of an isocaloric macronutrient-matched breakfast meal containing almonds on glycemic, hormonal, and appetite responses in men with type 2 diabetes: a randomized crossover study. Appl Physiol Nutr Metab 2020, 45: 520-529. <https://doi.org/10.1139/apnm-2019-0559>
  15. International Organization for Standardization: Determination of substances characteristic of green and black tea - Part 1: Content of total polyphenols in tea - Colorimetric method using Folin-Ciocalteu reagent (ISO Standard No. 14502-1:2005).  
<https://www.iso.org/standard/31356.html>
  16. Saito T, Abe D, Sekiya K: Flavanone exhibits PPAR $\gamma$  ligand activity and enhances differentiation of 3T3-L1 adipocytes. Biochem Biophys Res Commun 2009, 380:281-285. <https://doi.org/10.1016/j.bbrc.2009.01.058>
  17. Eniafe GO, Metibemu DS, Omotuyi OI, Ogunleye AJ, Inyan, OK, Adelakun NS, Adeniran YO, Adewumi B, Enejoh OA, Osunmuyiwa JO, Shodehinde SA, Oyeneyin OE: Agemone mexicana flavanones; apposite inverse agonists of the  $\beta_2$ -adrenergic receptor in asthma treatment. Bioinformation 2018, 14: 60-67. <https://doi.org/10.6026/97320630014060>
  18. Chin Y, Kong JY, Han S: Flavonoids as receptor tyrosine kinase FLT3 inhibitors. Bioorganic Med Chem Lett 2013, 23:1768-1770. <https://doi.org/10.1016/j.bmcl.2013.01.049>
  19. Zingg J: Vitamin E: A role in signal transduction. Annu Rev Nutr 2015, 35:135-173. <https://doi.org/10.1146/annurev-nutr-071714-034347>
  20. Fujimura Y, Yamada K, Tachibana H: A lipid raft-associated 67kDa laminin receptor mediates suppressive effect of epigallocatechin-3-O-gallate on Fc $\epsilon$ RI expression. Biochem Biophys Res Commun 2005, 336:674-681. <https://doi.org/10.1016/j.bbrc.2005.08.146>
  21. Fanani ML, Topham MK, Walsh JP, Epand RM: Lipid modulation of the activity of diacylglycerol kinase  $\alpha$ - and  $\zeta$ -isoforms: Activation by Phosphatidylethanolamine and Cholesterol. Biochemistry 2004, 43:14767-14777. <https://doi.org/10.1021/bi049145z>
  22. Chen CO, Blumberg JB: In vitro activity of almond skin polyphenols for scavenging free radicals and inducing quinone reductase. J Agric Food Chem 2008, 56: 4427-4434. <https://doi.org/10.1021/jf800061z>
  23. Mandalari G, Bisignano C, Genovese T, Mazzon E, Wickham M, Paterniti I, Cuzzocrea S: Natural almond skin reduced oxidative stress and inflammation in an experimental model of inflammatory bowel disease. Int Immunopharmacol 2011, 11:915-924. <https://doi.org/10.1016/j.intimp.2011.02.003>
  24. Bisignano C, Mandalari G, Smeriglio A, Trombetta D, Pizzo M, Pennisi R, Sciortino M: Almond skin extracts abrogate HSV-1 replication by blocking virus binding to the cell. Viruses 2017, 9: 178. <https://doi.org/10.3390/v9070178>
  25. Kaneko YK, Kobayashi Y, Motoki K, Nakata K, Miyagawa S, Yamamoto M, Hayashi D, Shirai Y, Sakane F, Ishikawa T: Depression of type I diacylglycerol kinases in pancreatic  $\beta$ -cells from male mice results in impaired insulin secretion. Endocrinology 2013, 154: 4089-4098. <https://doi.org/10.1210/en.2013-1356>