



# Regulation of Pancreatic $\beta$ -Cell Mass by Gene-Environment Interaction

Asahara, Shun-ichiro

Inoue, Hiroyuki

Kido, Yoshiaki

---

## (Citation)

Diabetes & Metabolism Journal, 46(1):38-48

## (Issue Date)

2022-01

## (Resource Type)

journal article

## (Version)

Version of Record

## (Rights)

© 2022 Korean Diabetes Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any...

## (URL)

<https://hdl.handle.net/20.500.14094/90009435>



# Regulation of Pancreatic $\beta$ -Cell Mass by Gene-Environment Interaction

Shun-ichiro Asahara<sup>1</sup>, Hiroyuki Inoue<sup>2</sup>, Yoshiaki Kido<sup>1,2</sup>

<sup>1</sup>Division of Diabetes and Endocrinology, Department of Internal Medicine, Kobe University Graduate School of Medicine, Kobe,

<sup>2</sup>Division of Medical Chemistry, Department of Metabolism and Diseases, Kobe University Graduate School of Health Sciences, Kobe, Japan

The main pathogenic mechanism of diabetes consists of an increase in insulin resistance and a decrease in insulin secretion from pancreatic  $\beta$ -cells. The number of diabetic patients has been increasing dramatically worldwide, especially in Asian people whose capacity for insulin secretion is inherently lower than that of other ethnic populations. Causally, changes of environmental factors in addition to intrinsic genetic factors have been considered to have an influence on the increased prevalence of diabetes. Particular focus has been placed on “gene-environment interactions” in the development of a reduced pancreatic  $\beta$ -cell mass, as well as type 1 and type 2 diabetes mellitus. Changes in the intrauterine environment, such as intrauterine growth restriction, contribute to alterations of gene expression in pancreatic  $\beta$ -cells, ultimately resulting in the development of pancreatic  $\beta$ -cell failure and diabetes. As a molecular mechanism underlying the effect of the intrauterine environment, epigenetic modifications have been widely investigated. The association of diabetes susceptibility genes or dietary habits with gene-environment interactions has been reported. In this review, we provide an overview of the role of gene-environment interactions in pancreatic  $\beta$ -cell failure as revealed by previous reports and data from experiments.


**Keywords:** Epigenomics; Fetal growth retardation; Gene-environment interaction; Genome-wide association study; Insulin-secreting cells

## INTRODUCTION

The incidence of diabetes has been increasing rapidly for several decades worldwide. There were more than 400 million patients with diabetes in 2019, and this number is estimated to reach 700 million by 2045 [1]. In the Western Pacific region, including East Asia, there was an especially large increase in the number of diabetic patients after World War II because of dietary changes and a decrease in physical activity. East Asian individuals have lower insulin secretory capacity than Caucasians; thus, East Asians are considered to have a tendency to develop pancreatic  $\beta$ -cell failure owing to genetic factors [2-4].

Previous work has shown the importance of insulin signaling in pancreatic  $\beta$ -cells using  $\beta$ -cell-specific transgenic mice. In 2006, Hashimoto et al. [5] clarified that a deficiency of

3-phosphoinositide dependent protein kinase 1 (Pdk1), an insulin signaling-related molecule, resulted in severe pancreatic  $\beta$ -cell failure, leading to hyperglycemia. In addition, it has been reported that mice deficient for tuberous sclerosis complex 2, a downstream molecule of Pdk1, exclusively in pancreatic  $\beta$ -cells develop hyperinsulinemia as a result of increased  $\beta$ -cell mass at younger ages, but show downregulation of insulin signaling induced by autophagic impairment and negative feedback, resulting in pancreatic  $\beta$ -cell failure at older ages [6-8]. To date, many investigators have used genetically modified mice as the main tool for the study of  $\beta$ -cell failure. These studies have revealed the pathogenesis of diabetes and the important role of various molecules in the onset of pancreatic  $\beta$ -cell failure. However, diabetes is a multifactorial disease. There are only a few monogenic forms of diabetes, such as maturity onset dia-

Corresponding author: Yoshiaki Kido  <https://orcid.org/0000-0003-2433-5799>  
Department of Metabolism and Diseases, Kobe University Graduate School of Health Sciences, 7-10-2 Tomogaoka, Suma-ku, Kobe 654-0142, Japan  
E-mail: kido@med.kobe-u.ac.jp

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.


betes of the young, and it is considered that pancreatic  $\beta$ -cell failure as well as diabetes generally occur through a combination of genetic and environmental factors. The rapid increase in the number of diabetic patients in East Asia must be the result of genetic factors inherent in Asian people and environmental factors that have changed recently. Furthermore, for the combination of genetic and environmental factors, synergistic effects are considered to be more important than additive effects for the onset of diabetes and pancreatic  $\beta$ -cell failure. Thus, environmental factors increase the influence of some genetic factors synergistically, resulting in the development of diabetes and pancreatic  $\beta$ -cell failure (Fig. 1). Understanding gene-environment interactions is expected to lead to the personalization of diet and exercise therapy. In this review, we describe gene-environment interactions according to the findings of previous studies.

## PHENOTYPIC VARIATIONS AMONG MOUSE STRAINS AND HUMAN ETHNIC GROUPS

First, we discuss the effects of diet in different mouse strains as a simple example of gene-environment interactions. Currently,

C57BL/6J is the most widely used mouse strain in animal experiments because this strain readily shows phenomena such as obesity with a high-fat diet (HFD) [9,10]. C57BL/6J mice are lean with a normal chow diet, but they develop hyperglycemia, hyperinsulinemia, and hyperleptinemia with a HFD [11–13]. On the other hand, strains such as 129/Sv or A/J are resistant to obesity and hyperglycemia under a HFD [14]. When C57BL/6J and 129/Sv mice are fed an HFD, 129/Sv mice show an increase in thermogenesis and maintain insulin sensitivity. When the genetic backgrounds of both strains were compared by a genome-wide scan, there was a significant difference in a locus associated with insulin resistance located on chromosome 14 [15]. Namely, the genetic differences between these strains are considered to produce their divergent phenotypic responses to a HFD.

Such phenomena are also observed in humans, with differences in body mass index, energy consumption, insulin resistance, and insulin secretion between East Asian and Caucasian populations [16–20]. Moreover, it has been recently reported that the regulation of pancreatic  $\beta$ -cell mass varies among racial groups [21–24]. Saisho et al. [21] found that pancreatic  $\beta$ -cell mass was increased in an obese group of healthy Cauca-



**Fig. 1.** Model of the associations between gene-environment interactions and pancreatic  $\beta$ -cell mass. In type 1 diabetes mellitus, an autoimmune attack is induced by the combination of genetic factors (human leukocyte antigen [HLA] haplotypes) and environmental factors (obesity, endoplasmic reticulum [ER] stress, dietary habits, and viral infection), resulting in a decrease in pancreatic  $\beta$ -cell mass. In type 2 diabetes mellitus, pancreatic  $\beta$ -cell mass is mainly regulated by the proliferation and apoptosis of pancreatic  $\beta$ -cells. Epigenetic modifications caused by intrauterine growth restriction (IUGR) are an important mechanism as a genetic factor. Environmental factors such as lipotoxicity, obesity, and ER stress caused by high-fat diet (HFD) feeding induce pancreatic  $\beta$ -cell failure in combination with genetic factors in a synergistic manner. Green boxes show genetic factors (non-modifiable factors) and orange boxes show environmental factors (modifiable factors).

sians when compared with a lean group, which was considered to be due to a compensatory mechanism for insulin resistance. Conversely, when pancreatic  $\beta$ -cell mass was quantified in healthy Japanese subjects and compared between obese and lean groups, there was no significant difference in pancreatic  $\beta$ -cell mass between the groups, but there was a slight decrease in the obese group compared with the lean group [22]. These results indicate that the compensatory mechanisms for pancreatic  $\beta$ -cell mass under conditions of insulin resistance, such as obesity, are different between Caucasian and Japanese individuals. It is considered that the phenotype may differ according to genetic factors, even if the dietary habits are the same.

## DEVELOPMENTAL ORIGINS OF HEALTH AND DISEASE

Environmental factors can affect genetic factors in adulthood, but the most important gene-environment interactions occur during the fetal period. In 1986, Barker and Osmond [25] proposed a theory that malnutrition in early life, including fetal life, causes susceptibility to ischemic heart disease and diabetes in adulthood (Barker's hypothesis). This hypothesis, which is also referred to as the Developmental Origins of Health and Disease theory, suggests that the embryonic or neonatal environment significantly contributes to the onset of future metabolic diseases [26-30]. A 50-year epidemiological survey reported that children of mothers who experienced the "Dutch famine" at the end of World War II have a high incidence of hypertension, obesity, or glucose intolerance [31,32]. Furthermore, the mechanism underlying this phenomenon has been elucidated using animal models. Yura et al. [33] generated intrauterine growth restriction (IUGR) mice with uterine artery ligation as a model of undernutrition. They showed that IUGR mice have significantly lower birth weight than control mice, but there is no significant difference in body weight in adulthood between both groups because IUGR mice show "catch-up" growth after weaning. In addition, when both groups were fed a HFD, IUGR mice exhibit significantly more severe obesity and glucose intolerance [33]. This is considered to be due to a "neonatal leptin surge," that is, an increase in serum leptin levels during the neonatal period, which is considered to lead to the onset of obesity by inducing leptin resistance.

Other reports have indicated that catch-up growth is associated with the development of obesity in later life in humans and mice [34-38]. Catch-up growth is considered to contribute

significantly to the development of metabolic syndrome. It has previously been reported that catch-up growth occurs in pancreatic  $\beta$ -cell mass as well as the whole body in the adult period and insulin signaling has a key mechanistic role in this process [39,40]. In addition, the offspring of female rats lacking vitamin B12 have been reported to be susceptible to insulin resistance in adulthood, and the offspring of pregnant rats fed a low-protein diet also develop insulin resistance through the reduction of protein kinase C zeta (PKC $\zeta$ ) expression in skeletal muscle [41-43].

Recently, an increase in fat intake has become a severe problem in developed countries. The offspring of HFD-fed female monkeys exhibit increased expression of gluconeogenic enzymes and transcription factors in hepatic tissues, resulting in the development of nonalcoholic fatty liver disease [44]. Interestingly, the female offspring of male HFD-fed rats exhibit pancreatic  $\beta$ -cell failure and hyperglycemia [45]. Compared with control mice, the female offspring of HFD-fed fathers show a change in the expression of 642 genes in microarray analysis using pancreatic islets. These changes were considered to induce pancreatic  $\beta$ -cell failure.

As described above, many reports have described the effects of fetal and neonatal environments on genetic factors, leading to type 2 diabetes mellitus (T2DM) and obesity. The most well-known molecular mechanism underlying this process is epigenetics, and we discuss the association between T2DM and epigenetics in the next section.

## EPIGENETICS

Epigenetics is regarded as an important molecular mechanism in not only metabolic diseases but also various other diseases. Epigenetics is a system that changes gene expression via DNA or histone modifications without mutation of the nucleotide sequence [46-48]. As it is considered that environmental changes can affect gene expression, we should not forget this mechanism when we consider gene-environment interactions. In particular, changes in the intrauterine environment can have an impact on epigenetic modifications, and fetal nutritional conditions memorized in embryonic tissue have been found to lead to the development of various diseases in the future (metabolic memory) [49-51]. Park et al. [52] generated an IUGR rat model by ligation of the uterine artery, and these rats developed pancreatic  $\beta$ -cell failure. When they analyzed epigenetic modifications in isolated islets of IUGR rats, they found that

the expression of pancreatic and duodenal homeobox 1 (Pdx1), a transcription factor, was decreased through an increase in DNA methylation and a change of histone modification at the Pdx1 promoter. The offspring of low-protein diet-fed female mice developed T2DM in adulthood, and there was a reduction of hepatocyte nuclear factor 4 alpha (HNF4 $\alpha$ ) expression in the pancreatic islets of those mice. This phenomenon was shown to be due to the inhibition of HNF4 $\alpha$  promoter-enhancer interactions [53]. In addition, in stem cells isolated from cord blood of neonates with IUGR, changes of DNA methylation in multiple genes including HNF4 $\alpha$  were found [54]. IUGR was predicted to influence the expression of various transcription factors, regardless of species or tissue.

The effects of IUGR have also been investigated in other insulin target organs [55,56], for instance, histone modifications at the glucose transporter type 4 (Glut4) promoter contribute to a reduction in Glut4 expression through the suppression of transcription in the skeletal muscle of IUGR rats [57]. Furthermore, in hepatocytes, Ehara et al. [58] clarified that the expression of glycerol-3-phosphate acyltransferase 1 was regulated by DNA methylation of its promoter. Interestingly, the expression of lipogenic genes is affected in the male offspring (F1) of IUGR mice, and even in hepatocytes of F2 mice. The expression of liver X receptor-alpha (Lxra) and DNA methylation at the Lxra promoter in hepatocytes of F2 mice has been reported to be altered significantly [59]. The fact that the effect of IUGR was passed on not only to children but also to grandchildren was a huge surprise. Possibly, the effects of the Dutch famine after World War II may still remain today.

## TYPE 1 DIABETES MELLITUS AND GENE-ENVIRONMENT INTERACTIONS

Type 1 diabetes mellitus (T1DM) is induced through the destruction of pancreatic  $\beta$ -cells by the autoimmune response [60,61], and it is likely to occur during childhood [62]. However, a number of individuals develop T1DM in adulthood [63]. Two large epidemiological studies, Diabetes Mondiale (DIAMOND) and European Diabetes (EURODIAB), reported that the incidence of T1DM was increasing worldwide [64,65]. The reasons given for this included changes in environmental factors. Therefore, in this section, we describe the relationship between gene-environment interactions and T1DM.

Specific human leukocyte antigen (HLA) alleles are genetic risk factors for the onset of T1DM [66] and influence its inci-

dence and age of onset [67,68]. The highest risk HLA haplotypes are *DRB1\*03:01-DQA1\*05:01-DQB1\*02:01* (known as DR3) and *DRB1\*04-DQA1\*03:01-DQB1\*03:02* (known as DR4). Individuals heterozygous for DR3/4 have an odds ratio (OR) of 16.59 for T1DM, while DR3/3 and DR4/4 homozygotes have ORs of 6.32 and 5.68, respectively [66]. These ORs are considerably higher than those of T2DM susceptibility genes. However, it is difficult to explain how the number of T1DM patients is increased by genetic factors alone. In recent years, the number of obese people has increased gradually worldwide [69-71]. In addition, the prevalence of overweight and obesity has risen in children with T1DM [72,73]. The increase in the number of obese subjects may be associated with the increase in the number of patients with T1DM. Interestingly, in Finland and Norway, which are two high-incidence T1DM countries, a rate of increase of T1DM has plateaued in recent years [74], indicating that this phenomenon is due to a plateau in the growth of obesity in both countries [75]. The ratio of obese subjects in a T1DM group was shown to be nearly equivalent to that in the general population or underwent a rapid increase [76]. In recent reports, the risk of developing T1DM was 63% higher in obese children than in lean children [77], and obesity increased susceptibility to insulin resistance, resulting in stress on  $\beta$ -cells and rendering them more vulnerable to an autoimmune attack [78].

Viral infection has been focused on as an important environmental factor of T1DM. It has been previously reported that enteroviruses, coxsackievirus B, cytomegalovirus, rotaviruses, and encephalomyocarditis virus contribute to the pathogenesis of T1DM [79,80]. In addition, dietary habits were found to be associated with the onset of T1DM as well as T2DM. Protein from cow milk and gluten are associated with an increase or decrease in the risk of developing T1DM [81-84].

The mechanisms associated with the destruction of pancreatic  $\beta$ -cells by environmental factors have not been elucidated fully [85]. One possibility is that endoplasmic reticulum (ER) stress contributes to the destruction of  $\beta$ -cells [86]. Inflammatory cytokines released by insulinitis disrupt ER homeostasis, resulting in the induction of the unfolded protein response. During this adaptive phase, activation of the inflammatory response is predicted to play an important role in the pathogenesis of T1DM [87]. Furthermore, the fact that ER stress induces translational errors suggests that aberrant insulin produced by ER stress may become a target of the autoimmune response as an antigenic peptide [88].


## INTERACTIONS BETWEEN DIABETES SUSCEPTIBILITY GENES AND HFD FEEDING

Finally, we discuss recent findings about gene-environment interactions in T2DM. In 2008, the potassium voltage-gated channel subfamily Q member 1 (*KCNQ1*) gene was identified as a T2DM susceptibility gene through large-scale single nucleotide polymorphism analysis of Japanese T2DM patients [89,90]. Subsequent reports from many facilities confirmed that there is an association between *KCNQ1* mutations and reduced insulin secretion [91-94], but the mechanism for the onset of diabetes remains unclear. It was previously reported that decreased expression of the non-coding RNA *Kcnq1ot1*, which is expressed from within the *Kcnq1* genetic region, increases the expression of cyclin-dependent kinase inhibitor 1C (*Cdkn1c*) through epigenetic modifications, resulting in reduced pancreatic  $\beta$ -cell mass [95]. *Kcnq1* is an imprinted gene, and *Kcnq1ot1* expression is decreased in pancreatic  $\beta$ -cells of mice only when a *Kcnq1* genetic mutation is paternally inherited. As a result, pancreatic  $\beta$ -cell mass is decreased by enhancing *Cdkn1c* expression. These results may partially indicate that *Kcnq1* mutation is an underlying pathogenic mechanism of T2DM. However, the OR of developing T2DM was only 1.4, suggesting that T2DM is not induced by genetic mutation alone. Therefore, focus shifted to the fact that there is a CCAAT motif in the *Cdkn1c* promoter region. As the CCAAT motif is a binding site for members of the CCAAT enhancer-binding protein (C/EBP) transcription factor family, it was hypothesized that binding of C/EBP family members to this motif may further enhance *Cdkn1c* expression. C/EBP $\beta$  expression was found to be enhanced in the islets of *db/db* mice, a T2DM mouse model, and reported that the accumulation of C/EBP $\beta$  increased vulnerability to ER stress and resulted in reduced  $\beta$ -cell mass [96,97]. In addition, recent studies revealed that C/EBP $\beta$  also accumulates in the pancreatic islets of HFD-fed mice, which may cause pancreatic  $\beta$ -cell failure through enhanced *Cdkn1c* expression and the induction of ER stress, and is considered to contribute to the development of T2DM.

Therefore, the influence of C/EBP $\beta$  accumulation in pancreatic  $\beta$ -cells was investigated when *Kcnq1ot1* expression is decreased. A significant increase in fed blood glucose levels and a reduction in serum insulin levels were observed in mice with decreased *Kcnq1ot1* expression and C/EBP $\beta$  overexpression exclusively in pancreatic  $\beta$ -cells, compared to control groups,

which was accompanied by a significant decrease in pancreatic  $\beta$ -cell mass. This is presumed to be due to increased binding of C/EBP $\beta$  to the *Cdkn1c* promoter as a result of epigenetic modifications such as loosening chromatin structure under reduced *Kcnq1ot1* expression. These results revealed that the synergistic effect of genetic factors (*Kcnq1* mutation) and environmental factors (C/EBP $\beta$  accumulation) could induce markedly reduced pancreatic  $\beta$ -cell mass.

Furthermore, the role of eukaryotic translation initiation factor 2 $\alpha$  kinase 4 (*EIF2AK4*) was investigated. *EIF2AK4* encodes general control nonderepressible 2 (GCN2), which, in yeast and mammals, is activated as a result of amino acid deficiency and was identified as a T2DM susceptibility gene, in addition to *KCNQ1*, through large-scale single nucleotide polymorphism analysis of Japanese T2DM patients [98-101]. It was found that insulin secretion was reduced in carriers of the *EIF2AK4* risk allele compared with individuals without the risk allele [102]. In addition, HFD-fed GCN2 knockout mice exhibit reduced pancreatic  $\beta$ -cell mass and decreased insulin secretion, while the phenotype of GCN2 knockout mice fed a normal chow diet is comparable to that of control mice [102]. This phenomenon was attributed to the constitutive enhancement of mechanistic target of rapamycin complex 1 (mTORC1) activity in pancreatic  $\beta$ -cells of GCN2 knockout mice. In fact, chronic enhancement of mTORC1 signaling was shown to result in reduced pancreatic  $\beta$ -cell mass [6-8], and this mechanism was considered to contribute to the development of reduced  $\beta$ -cell mass in HFD-fed GCN2 knockout mice. GCN2 is known to be activated under cellular amino acid deficiency [99], and it was found that amino acid concentration was decreased in the pancreatic  $\beta$ -cells of HFD-fed mice [102]. The HFD was shown to increase insulin demand in the whole body, and a large quantity of amino acids was consumed due to insulin production in pancreatic  $\beta$ -cells [103], resulting in reduced amino acid concentrations in these cells [102]. In pancreatic  $\beta$ -cells of HFD-fed GCN2 knockout mice, GCN2 cannot be activated in spite of amino acid deficiency. HFD-fed GCN2 knockout mice were considered to show hyperglycemia accompanied by reduced pancreatic  $\beta$ -cell mass through the constitutive enhancement of mTORC1 due to GCN2 inactivation (Fig. 2). The effect of GCN2 alone is not sufficient to induce pancreatic  $\beta$ -cell failure; however, environmental factors such as a HFD interact with GCN2 synergistically to cause pancreatic  $\beta$ -cell failure. As far as previous studies have shown, pancreatic islets were the only tissue in which cellular amino



**Fig. 2.** Model for general control nonderepressible 2 (GCN2)-dependent regulation of pancreatic  $\beta$ -cell mass during high-fat diet (HFD) feeding. HFD feeding increases insulin demand, resulting in reduced amino acid concentrations due to their consumption in pancreatic  $\beta$ -cells. HFD-fed GCN2 knockout mice develop hyperglycemia accompanied by reduced pancreatic  $\beta$ -cell mass through the constitutive enhancement of mechanistic target of rapamycin complex 1 (mTORC1) due to GCN2 inactivation. GCN2, a genetic factor, and environmental factors, such as a HFD, cause pancreatic  $\beta$ -cell failure in a synergistic manner. “P” indicates the phosphorylation of GCN2. Modified from Kanno et al. [102], with permission from American Society for Clinical Investigation.

acid levels were decreased [102]. This result indicates that the response of pancreatic  $\beta$ -cells to HFD feeding is unique, and that gene-environment interactions may have an especially strong influence on the viability of pancreatic  $\beta$ -cells, as many T2DM susceptibility genes are associated with insulin secretion. A recent report showed that pancreatic  $\beta$ -cell death caused by gene-environment interactions could be examined using human pluripotent stem cells [104]. This method may help to clarify the associations between diabetes susceptibility genes and gene-environment interactions.

## CONCLUSIONS

The explosive increase in the number of diabetic patients worldwide in recent years is mainly due to changes in the environment, given that there have been no significant or consistent changes in genetic factors. However, changes in environmental factors are likely to affect genetic factors. In this review, we documented the evidence for the relationship between gene-environment interactions and IUGR or diabetes susceptibility genes. However, gene-environment interactions are believed to be universally applicable to the contemporary pathology of diabetes, especially reduced  $\beta$ -cell mass. Therefore, further detailed analysis of these gene-environment interactions

may suggest the key for personalized medicine. We look forward to further progress in this field in the future.

## CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

## ORCID

Shun-ichiro Asahara <https://orcid.org/0000-0001-6661-9402>

Yoshiaki Kido <https://orcid.org/0000-0003-2433-5799>

## FUNDING

None

## ACKNOWLEDGMENTS

None

## REFERENCES

1. International Diabetes Federation. IDF Diabetes Atlas. 9th ed.

- Brussels: International Diabetes Federation; 2019.
2. Yabe D, Seino Y. Type 2 diabetes via  $\beta$ -cell dysfunction in East Asian people. *Lancet Diabetes Endocrinol* 2016;4:2-3.
  3. Yabe D, Seino Y, Fukushima M, Seino S.  $\beta$  Cell dysfunction versus insulin resistance in the pathogenesis of type 2 diabetes in East Asians. *Curr Diab Rep* 2015;15:602.
  4. Cho YS, Chen CH, Hu C, Long J, Ong RT, Sim X, et al. Meta-analysis of genome-wide association studies identifies eight new loci for type 2 diabetes in East Asians. *Nat Genet* 2011;44:67-72.
  5. Hashimoto N, Kido Y, Uchida T, Asahara S, Shigeyama Y, Matsuda T, et al. Ablation of PDK1 in pancreatic beta cells induces diabetes as a result of loss of beta cell mass. *Nat Genet* 2006;38:589-93.
  6. Shigeyama Y, Kobayashi T, Kido Y, Hashimoto N, Asahara S, Matsuda T, et al. Biphasic response of pancreatic beta-cell mass to ablation of tuberous sclerosis complex 2 in mice. *Mol Cell Biol* 2008;28:2971-9.
  7. Koyanagi M, Asahara S, Matsuda T, Hashimoto N, Shigeyama Y, Shibutani Y, et al. Ablation of TSC2 enhances insulin secretion by increasing the number of mitochondria through activation of mTORC1. *PLoS One* 2011;6:e23238.
  8. Bartolome A, Kimura-Koyanagi M, Asahara S, Guillen C, Inoue H, Teruyama K, et al. Pancreatic  $\beta$ -cell failure mediated by mTORC1 hyperactivity and autophagic impairment. *Diabetes* 2014;63:2996-3008.
  9. Jo J, Gavrilova O, Pack S, Jou W, Mullen S, Sumner AE, et al. Hypertrophy and/or hyperplasia: dynamics of adipose tissue growth. *PLoS Comput Biol* 2009;5:e1000324.
  10. Kim DH, Gutierrez-Aguilar R, Kim HJ, Woods SC, Seeley RJ. Increased adipose tissue hypoxia and capacity for angiogenesis and inflammation in young diet-sensitive C57 mice compared with diet-resistant FVB mice. *Int J Obes (Lond)* 2013;37:853-60.
  11. Eldar-Finkelman H, Schreyer SA, Shinohara MM, LeBoeuf RC, Krebs EG. Increased glycogen synthase kinase-3 activity in diabetes- and obesity-prone C57BL/6J mice. *Diabetes* 1999;48:1662-6.
  12. Surwit RS, Kuhn CM, Cochrane C, McCubbin JA, Feinglos MN. Diet-induced type II diabetes in C57BL/6J mice. *Diabetes* 1988;37:1163-7.
  13. West DB, Boozer CN, Moody DL, Atkinson RL. Dietary obesity in nine inbred mouse strains. *Am J Physiol* 1992;262(6 Pt 2):R1025-32.
  14. Surwit RS, Feinglos MN, Rodin J, Sutherland A, Petro AE, Opara EC, et al. Differential effects of fat and sucrose on the development of obesity and diabetes in C57BL/6J and A/J mice. *Metabolism* 1995;44:645-51.
  15. Almind K, Kahn CR. Genetic determinants of energy expenditure and insulin resistance in diet-induced obesity in mice. *Diabetes* 2004;53:3274-85.
  16. Chiu KC, Cohan P, Lee NP, Chuang LM. Insulin sensitivity differs among ethnic groups with a compensatory response in beta-cell function. *Diabetes Care* 2000;23:1353-8.
  17. Jensen CC, Cnop M, Hull RL, Fujimoto WY, Kahn SE; American Diabetes Association GENNID Study Group. Beta-cell function is a major contributor to oral glucose tolerance in high-risk relatives of four ethnic groups in the U.S. *Diabetes* 2002;51:2170-8.
  18. Gerstein HC, Anand S, Yi QL, Vuksan V, Lonn E, Teo K, et al. The relationship between dysglycemia and atherosclerosis in South Asian, Chinese, and European individuals in Canada: a randomly sampled cross-sectional study. *Diabetes Care* 2003;26:144-9.
  19. Davis TM, Mulder H, Lokhnygina Y, Aschner P, Chuang LM, Raffo Grado CA, et al. Effect of race on the glycaemic response to sitagliptin: insights from the Trial Evaluating Cardiovascular Outcomes with Sitagliptin (TECOS). *Diabetes Obes Metab* 2018;20:1427-34.
  20. Ma RC, Chan JC. Type 2 diabetes in East Asians: similarities and differences with populations in Europe and the United States. *Ann N Y Acad Sci* 2013;1281:64-91.
  21. Saisho Y, Butler AE, Manesso E, Elashoff D, Rizza RA, Butler PC.  $\beta$ -Cell mass and turnover in humans: effects of obesity and aging. *Diabetes Care* 2013;36:111-7.
  22. Kou K, Saisho Y, Satoh S, Yamada T, Itoh H. Change in  $\beta$ -cell mass in Japanese nondiabetic obese individuals. *J Clin Endocrinol Metab* 2013;98:3724-30.
  23. Roh E, Kim KM, Park KS, Kim YJ, Chun EJ, Choi SH, et al. Comparison of pancreatic volume and fat amount linked with glucose homeostasis between healthy Caucasians and Koreans. *Diabetes Obes Metab* 2018;20:2642-52.
  24. Rahier J, Guiot Y, Goebbels RM, Sempoux C, Henquin JC. Pancreatic beta-cell mass in European subjects with type 2 diabetes. *Diabetes Obes Metab* 2008;10 Suppl 4:32-42.
  25. Barker DJ, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet* 1986;1:1077-81.
  26. Simeoni U, Osmond C, Garay R, Buffat C, Boubred F, Chagnaud C, et al. Leptin and insulin in young adulthood are associated

- ated with weight in infancy. *J Endocrinol* 2020;244:249-59.
27. Huang RC, Prescott SL, Godfrey KM, Davis EA. Assessment of cardiometabolic risk in children in population studies: underpinning developmental origins of health and disease mother-offspring cohort studies. *J Nutr Sci* 2015;4:e12.
  28. Heindel JJ, Blumberg B. Environmental obesogens: mechanisms and controversies. *Annu Rev Pharmacol Toxicol* 2019; 59:89-106.
  29. Tain YL, Hsu CN. Developmental programming of the metabolic syndrome: can we reprogram with resveratrol? *Int J Mol Sci* 2018;19:2584.
  30. Lee WC, Wu KLH, Leu S, Tain YL. Translational insights on developmental origins of metabolic syndrome: focus on fructose consumption. *Biomed J* 2018;41:96-101.
  31. Roseboom TJ, van der Meulen JH, Ravelli AC, Osmond C, Barker DJ, Bleker OP. Effects of prenatal exposure to the Dutch famine on adult disease in later life: an overview. *Mol Cell Endocrinol* 2001;185:93-8.
  32. Painter RC, de Rooij SR, Bossuyt PM, de Groot E, Stok WJ, Osmond C, et al. Maternal nutrition during gestation and carotid arterial compliance in the adult offspring: the Dutch famine birth cohort. *J Hypertens* 2007;25:533-40.
  33. Yura S, Itoh H, Sagawa N, Yamamoto H, Masuzaki H, Nakao K, et al. Role of premature leptin surge in obesity resulting from intrauterine undernutrition. *Cell Metab* 2005;1:371-8.
  34. Ozanne SE, Hales CN. Lifespan: catch-up growth and obesity in male mice. *Nature* 2004;427:411-2.
  35. Stettler N, Zemel BS, Kumanyika S, Stallings VA. Infant weight gain and childhood overweight status in a multicenter, cohort study. *Pediatrics* 2002;109:194-9.
  36. Nordman H, Jaaskelainen J, Voutilainen R. Birth size as a determinant of cardiometabolic risk factors in children. *Horm Res Paediatr* 2020;93:144-53.
  37. Shi H, Yang X, Wu D, Wang X, Li T, Liu H, et al. Insights into infancy weight gain patterns for term small-for-gestational-age babies. *Nutr J* 2018;17:97.
  38. Berends LM, Dearden L, Tung YC, Voshol P, Fernandez-Twinn DS, Ozanne SE. Programming of central and peripheral insulin resistance by low birthweight and postnatal catch-up growth in male mice. *Diabetologia* 2018;61:2225-34.
  39. Yoshida Y, Fuchita M, Kimura-Koyanagi M, Kanno A, Matsuda T, Asahara SI, et al. Contribution of insulin signaling to the regulation of pancreatic beta-cell mass during the catch-up growth period in a low birth weight mouse model. *Diabetol Int* 2014;5:43-52.
  40. Inoue T, Kido Y, Asahara S, Matsuda T, Shibutani Y, Koyanagi M, et al. Effect of intrauterine undernutrition during late gestation on pancreatic beta cell mass. *Biomed Res* 2009;30:325-30.
  41. Stewart CP, Christian P, Schulze KJ, Arguello M, LeClerq SC, Khatry SK, et al. Low maternal vitamin B-12 status is associated with offspring insulin resistance regardless of antenatal micronutrient supplementation in rural Nepal. *J Nutr* 2011;141: 1912-7.
  42. Ozanne SE, Olsen GS, Hansen LL, Tingey KJ, Nave BT, Wang CL, et al. Early growth restriction leads to down regulation of protein kinase C zeta and insulin resistance in skeletal muscle. *J Endocrinol* 2003;177:235-41.
  43. Ozanne SE, Jensen CB, Tingey KJ, Storgaard H, Madsbad S, Vaag AA. Low birthweight is associated with specific changes in muscle insulin-signalling protein expression. *Diabetologia* 2005;48:547-52.
  44. McCurdy CE, Bishop JM, Williams SM, Grayson BE, Smith MS, Friedman JE, et al. Maternal high-fat diet triggers lipotoxicity in the fetal livers of nonhuman primates. *J Clin Invest* 2009;119:323-35.
  45. Ng SF, Lin RC, Laybutt DR, Barres R, Owens JA, Morris MJ. Chronic high-fat diet in fathers programs  $\beta$ -cell dysfunction in female rat offspring. *Nature* 2010;467:963-6.
  46. Gruenert DC, Cozens AL. Inheritance of phenotype in mammalian cells: genetic vs. epigenetic mechanisms. *Am J Physiol* 1991;260(6 Pt 1):L386-94.
  47. Takeshima H, Ushijima T. Accumulation of genetic and epigenetic alterations in normal cells and cancer risk. *NPJ Precis Oncol* 2019;3:7.
  48. Dalfra MG, Burlina S, Del Vescovo GG, Lapolla A. Genetics and epigenetics: new insight on gestational diabetes mellitus. *Front Endocrinol (Lausanne)* 2020;11:602477.
  49. Duran Fernandez-Feijoo C, Carrasco Carrasco C, Villalmazo Francisco N, Cebria Romero J, Fernandez Lorenzo JR, Jimenez-Chillaron JC, et al. Influence of catch up growth on spatial learning and memory in a mouse model of intrauterine growth restriction. *PLoS One* 2017;12:e0177468.
  50. Suter MA, Ma J, Vuguin PM, Hartil K, Fiallo A, Harris RA, et al. In utero exposure to a maternal high-fat diet alters the epigenetic histone code in a murine model. *Am J Obstet Gynecol* 2014;210:463.
  51. Tozour J, Hughes F, Carrier A, Vieau D, Delahaye F. Prenatal hyperglycemia exposure and cellular stress, a sugar-coated view of early programming of metabolic diseases. *Biomole-*

- cules 2020;10:1359.
52. Park JH, Stoffers DA, Nicholls RD, Simmons RA. Development of type 2 diabetes following intrauterine growth retardation in rats is associated with progressive epigenetic silencing of Pdx1. *J Clin Invest* 2008;118:2316-24.
  53. Sandovici I, Smith NH, Nitert MD, Ackers-Johnson M, Uribe-Lewis S, Ito Y, et al. Maternal diet and aging alter the epigenetic control of a promoter-enhancer interaction at the Hnf4a gene in rat pancreatic islets. *Proc Natl Acad Sci U S A* 2011;108:5449-54.
  54. Einstein F, Thompson RF, Bhagat TD, Fazzari MJ, Verma A, Barzilai N, et al. Cytosine methylation dysregulation in neonates following intrauterine growth restriction. *PLoS One* 2010;5:e8887.
  55. Xing Y, Zhang J, Wei H, Zhang H, Guan Y, Wang X, et al. Reduction of the PI3K/Akt related signaling activities in skeletal muscle tissues involves insulin resistance in intrauterine growth restriction rats with catch-up growth. *PLoS One* 2019;14:e0216665.
  56. Dunlop K, Cedrone M, Staples JF, Regnault TR. Altered fetal skeletal muscle nutrient metabolism following an adverse in utero environment and the modulation of later life insulin sensitivity. *Nutrients* 2015;7:1202-16.
  57. Raychaudhuri N, Raychaudhuri S, Thamotharan M, Devaskar SU. Histone code modifications repress glucose transporter 4 expression in the intrauterine growth-restricted offspring. *J Biol Chem* 2008;283:13611-26.
  58. Ehara T, Kamei Y, Takahashi M, Yuan X, Kanai S, Tamura E, et al. Role of DNA methylation in the regulation of lipogenic glycerol-3-phosphate acyltransferase 1 gene expression in the mouse neonatal liver. *Diabetes* 2012;61:2442-50.
  59. Martinez D, Pentinat T, Ribo S, Daviaud C, Bloks VW, Cebria J, et al. In utero undernutrition in male mice programs liver lipid metabolism in the second-generation offspring involving altered Lxra DNA methylation. *Cell Metab* 2014;19:941-51.
  60. Atkinson MA, Eisenbarth GS, Michels AW. Type 1 diabetes. *Lancet* 2014;383:69-82.
  61. Incani M, Serafini C, Satta C, Perra L, Scano F, Frongia P, et al. High prevalence of diabetes-specific autoimmunity in first-degree relatives of Sardinian patients with type 1 diabetes. *Diabetes Metab Res Rev* 2017;33:e2864.
  62. You WP, Henneberg M. Type 1 diabetes prevalence increasing globally and regionally: the role of natural selection and life expectancy at birth. *BMJ Open Diabetes Res Care* 2016;4:e000161.
  63. Monaghan M, Helgeson V, Wiebe D. Type 1 diabetes in young adulthood. *Curr Diabetes Rev* 2015;11:239-50.
  64. DIAMOND Project Group. Incidence and trends of childhood type 1 diabetes worldwide 1990-1999. *Diabet Med* 2006;23:857-66.
  65. Patterson CC, Dahlquist GG, Gyurus E, Green A, Soltesz G; EURODIAB Study Group. Incidence trends for childhood type 1 diabetes in Europe during 1989-2003 and predicted new cases 2005-20: a multicentre prospective registration study. *Lancet* 2009;373:2027-33.
  66. Noble JA, Valdes AM. Genetics of the HLA region in the prediction of type 1 diabetes. *Curr Diab Rep* 2011;11:533-42.
  67. Mishra R, Akerlund M, Cousminer DL, Ahlqvist E, Bradfield JP, Chesi A, et al. Genetic discrimination between LADA and childhood-onset type 1 diabetes within the MHC. *Diabetes Care* 2020;43:418-25.
  68. Buzzetti R, Prudente S, Copetti M, Dauriz M, Zampetti S, Garofolo M, et al. Clinical worthlessness of genetic prediction of common forms of diabetes mellitus and related chronic complications: a position statement of the Italian Society of Diabetology. *Nutr Metab Cardiovasc Dis* 2017;27:99-114.
  69. NCD Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *Lancet* 2016;387:1377-96.
  70. Asahara SI, Miura H, Ogawa W, Tamori Y. Sex difference in the association of obesity with personal or social background among urban residents in Japan. *PLoS One* 2020;15:e0242105.
  71. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 2017;390:2627-42.
  72. De Keukelaere M, Fieuws S, Reynaert N, Vandoorne E, Kerckhove KV, Asscherickx W, et al. Evolution of body mass index in children with type 1 diabetes mellitus. *Eur J Pediatr* 2018;177:1661-6.
  73. Corbin KD, Driscoll KA, Pratley RE, Smith SR, Maahs DM, Mayer-Davis EJ, et al. Obesity in type 1 diabetes: pathophysiology, clinical impact, and mechanisms. *Endocr Rev* 2018;39:629-63.
  74. Patterson CC, Harjutsalo V, Rosenbauer J, Neu A, Cinek O, Skrivarhaug T, et al. Trends and cyclical variation in the incidence of childhood type 1 diabetes in 26 European centres in the 25 year period 1989-2013: a multicentre prospective regis-

- tration study. *Diabetologia* 2019;62:408-17.
75. Public Health Report: Overweight and obesity in Norway. Available from: <https://www.fhi.no/en/op/hin/health-disease/overweight-and-obesity-in-norway---> (updated 2017 Nov 3).
  76. Fellinger P, Fuchs D, Wolf P, Heinze G, Luger A, Krebs M, et al. Overweight and obesity in type 1 diabetes equal those of the general population. *Wien Klin Wochenschr* 2019;131:55-60.
  77. Ferrara CT, Geyer SM, Liu YF, Evans-Molina C, Libman IM, Besser R, et al. Excess BMI in childhood: a modifiable risk factor for type 1 diabetes development? *Diabetes Care* 2017;40:698-701.
  78. Saltiel AR, Olefsky JM. Inflammatory mechanisms linking obesity and metabolic disease. *J Clin Invest* 2017;127:1-4.
  79. Rewers M, Ludvigsson J. Environmental risk factors for type 1 diabetes. *Lancet* 2016;387:2340-8.
  80. Nekoua MP, Bertin A, Sane F, Alidjinou EK, Lobert D, Trauet J, et al. Pancreatic beta cells persistently infected with coxsackievirus B4 are targets of NK cell-mediated cytolytic activity. *Cell Mol Life Sci* 2020;77:179-94.
  81. Lamb MM, Miller M, Seifert JA, Frederiksen B, Kroehl M, Rewers M, et al. The effect of childhood cow's milk intake and HLA-DR genotype on risk of islet autoimmunity and type 1 diabetes: the Diabetes Autoimmunity Study in the Young. *Pediatr Diabetes* 2015;16:31-8.
  82. Lempainen J, Tauriainen S, Vaarala O, Makela M, Honkanen H, Marttila J, et al. Interaction of enterovirus infection and cow's milk-based formula nutrition in type 1 diabetes-associated autoimmunity. *Diabetes Metab Res Rev* 2012;28:177-85.
  83. Ziegler AG, Schmid S, Huber D, Hummel M, Bonifacio E. Early infant feeding and risk of developing type 1 diabetes-associated autoantibodies. *JAMA* 2003;290:1721-8.
  84. Uusitalo U, Lee HS, Andren Aronsson C, Vehik K, Yang J, Hummel S, et al. Early infant diet and islet autoimmunity in the TEDDY study. *Diabetes Care* 2018;41:522-30.
  85. Op de Beeck A, Eizirik DL. Viral infections in type 1 diabetes mellitus: why the  $\beta$  cells? *Nat Rev Endocrinol* 2016;12:263-73.
  86. Eizirik DL, Miani M, Cardozo AK. Signalling danger: endoplasmic reticulum stress and the unfolded protein response in pancreatic islet inflammation. *Diabetologia* 2013;56:234-41.
  87. Thomaidou S, Zaldumbide A, Roep BO. Islet stress, degradation and autoimmunity. *Diabetes Obes Metab* 2018;20(Suppl 2):88-94.
  88. Kracht MJ, van Lummel M, Nikolic T, Joosten AM, Laban S, van der Slik AR, et al. Autoimmunity against a defective ribosomal insulin gene product in type 1 diabetes. *Nat Med* 2017;23:501-7.
  89. Yasuda K, Miyake K, Horikawa Y, Hara K, Osawa H, Furuta H, et al. Variants in KCNQ1 are associated with susceptibility to type 2 diabetes mellitus. *Nat Genet* 2008;40:1092-7.
  90. Unoki H, Takahashi A, Kawaguchi T, Hara K, Horikoshi M, Andersen G, et al. SNPs in KCNQ1 are associated with susceptibility to type 2 diabetes in East Asian and European populations. *Nat Genet* 2008;40:1098-102.
  91. Hu C, Wang C, Zhang R, Ma X, Wang J, Lu J, et al. Variations in KCNQ1 are associated with type 2 diabetes and beta cell function in a Chinese population. *Diabetologia* 2009;52:1322-5.
  92. Jonsson A, Isomaa B, Tuomi T, Taneera J, Salehi A, Nilsson P, et al. A variant in the KCNQ1 gene predicts future type 2 diabetes and mediates impaired insulin secretion. *Diabetes* 2009;58:2409-13.
  93. Tan JT, Nurbaya S, Gardner D, Ye S, Tai ES, Ng DP. Genetic variation in KCNQ1 associates with fasting glucose and beta-cell function: a study of 3,734 subjects comprising three ethnicities living in Singapore. *Diabetes* 2009;58:1445-9.
  94. Rosengren AH, Braun M, Mahdi T, Andersson SA, Travers ME, Shigeto M, et al. Reduced insulin exocytosis in human pancreatic  $\beta$ -cells with gene variants linked to type 2 diabetes. *Diabetes* 2012;61:1726-33.
  95. Asahara S, Etoh H, Inoue H, Teruyama K, Shibutani Y, Ihara Y, et al. Paternal allelic mutation at the Kcnq1 locus reduces pancreatic  $\beta$ -cell mass by epigenetic modification of Cdkn1c. *Proc Natl Acad Sci U S A* 2015;112:8332-7.
  96. Matsuda T, Kido Y, Asahara S, Kaisho T, Tanaka T, Hashimoto N, et al. Ablation of C/EBP $\beta$  alleviates ER stress and pancreatic beta cell failure through the GRP78 chaperone in mice. *J Clin Invest* 2010;120:115-26.
  97. Matsuda T, Takahashi H, Mieda Y, Shimizu S, Kawamoto T, Matsuura Y, et al. Regulation of pancreatic  $\beta$  cell mass by cross-interaction between CCAAT enhancer binding protein  $\beta$  induced by endoplasmic reticulum stress and AMP-activated protein kinase activity. *PLoS One* 2015;10:e0130757.
  98. Deval C, Chaveroux C, Maurin AC, Cherasse Y, Parry L, Carraro V, et al. Amino acid limitation regulates the expression of genes involved in several specific biological processes through GCN2-dependent and GCN2-independent pathways. *FEBS J* 2009;276:707-18.
  99. Zhang P, McGrath BC, Reinert J, Olsen DS, Lei L, Gill S, et al. The GCN2 eIF2 $\alpha$  kinase is required for adaptation to ami-

- no acid deprivation in mice. *Mol Cell Biol* 2002;22:6681-8.
100. Guo F, Cavener DR. The GCN2 eIF2 $\alpha$  kinase regulates fatty-acid homeostasis in the liver during deprivation of an essential amino acid. *Cell Metab* 2007;5:103-14.
101. Miyake K, Yang W, Hara K, Yasuda K, Horikawa Y, Osawa H, et al. Construction of a prediction model for type 2 diabetes mellitus in the Japanese population based on 11 genes with strong evidence of the association. *J Hum Genet* 2009;54:236-41.
102. Kanno A, Asahara SI, Furubayashi A, Masuda K, Yoshitomi R, Suzuki E, et al. GCN2 regulates pancreatic  $\beta$  cell mass by sensing intracellular amino acid levels. *JCI Insight* 2020;5:e128820.
103. Kanno A, Asahara SI, Masuda K, Matsuda T, Kimura-Koyanagi M, Seino S, et al. Compensatory hyperinsulinemia in high-fat diet-induced obese mice is associated with enhanced insulin translation in islets. *Biochem Biophys Res Commun* 2015;458:681-6.
104. Zhou T, Kim TW, Chong CN, Tan L, Amin S, Sadat Badieyan Z, et al. A hPSC-based platform to discover gene-environment interactions that impact human  $\beta$ -cell and dopamine neuron survival. *Nat Commun* 2018;9:4815.