

PDF issue: 2025-06-14

## AIDS-Related Burkitt Lymphoma Presenting Multiple Nodules in the Pancreas

Kohashi, Shinya Sakai, Arata Kodama, Yuzo

(Citation) Clinical Gastroenterology and Hepatology,20(5):e920

(Issue Date) 2022-05

(Resource Type) journal article

(Version) Accepted Manuscript

(Rights)

© 2022 by the AGA Institute. This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International license.

(URL)

https://hdl.handle.net/20.500.14094/90009549



## AIDS-related Burkitt's lymphoma presenting multiple nodules in the pancreas

Shinya Kohashi<sup>1</sup>, Arata Sakai<sup>1</sup> and Yuzo Kodama<sup>1</sup>

<sup>1</sup>Division of Gastroenterology, Department of Internal Medicine, Kobe University Graduate School of Medicine

## **Corresponding Author:**

Arata Sakai, MD, PhD

Division of Gastroenterology, Department of Internal Medicine

Kobe University Graduate School of Medicine

7-5-1 Kusunoki-cho, Chuo-ku, Kobe, Hyogo 650-0017, Japan

Telephone: +81-78-382-6305; Fax: +81-78-382-6309

E-mail: asakai@med.kobe-u.ac.jp

Disclosure statement: The authors have no conflicts of interest to declare.

Grant support: Supported by JSPS KAKENHI grant number JP19K07938 (AS).

## Abbreviations:

AIDS: acquired immune deficiency syndrome

cART: combination antiretroviral therapy

CT: Computed tomography

EUS: endoscopic ultrasonography

sIL-2R: soluble interleukin-2 receptor

A 60-year-old man being treated with combination antiretroviral therapy (cART) for acquired immune deficiency syndrome (AIDS) with suspected pancreatic tumor on computed tomography (CT) was referred to our department. He had general malaise, but no other symptom. Laboratory data were as follows: leukocyte count 1700 /µL (normal range, 3300-8600 /µL), CD4 cell count 86 /µL, lactate dehydrogenase 838 U/L (124-222 U/L), pancreatic amylase 123 U/L (16–52 U/L). Serum soluble interleukin-2 receptor (sIL-2R), used as a diagnostic and prognostic marker for malignant lymphoma, was 2739 U/mL (121-613 U/mL). Abdominal contrast-enhanced CT revealed multiple hypo-vascular nodules in the entire pancreas (Figure A). Endoscopic ultrasonography (EUS) revealed multiple hypoechoic nodules in the entire pancreas (Figure B). EUS-guided fine needle aspiration biopsy from the pancreatic nodule was performed. Histological examination revealed the infiltration of malignant lymphocytes (Figure C). These atypical lymphocytes were positive for CD20, CD10, BCL-6, and negative for BCL-2. Ki-67 displayed a proliferation index >95%. FISH showed c-MYC gene rearrangement. Based on these findings, the diagnosis was AIDS-related Burkitt's lymphoma (BL). CHOP (cyclophosphamide, hydroxydaunorubicin, oncovin and prednisone) chemotherapy was started with continued cART. He had a complete response and was still alive with no sign of recurrence at 9 months after chemotherapy (Figure D).







