



Effects of acute-phase multidisciplinary rehabilitation on unplanned readmissions after cardiac surgery

Ogawa, Masato ; Satomi-Kobayashi, Seimi ; Yoshida, Naofumi ; Tsuboi, Yasunori ; Komaki, Kodai ; Wakida, Kumiko ; Gotake, Yasuko ; Izawa, P. ...

(Citation)

The Journal of Thoracic and Cardiovascular Surgery, 161(5):1853-1860.e2

(Issue Date)

2021-05

(Resource Type)

journal article

(Version)

Accepted Manuscript

(URL)

<https://hdl.handle.net/20.500.14094/90009575>



**Effects of Acute Phase Multidisciplinary Rehabilitation on Unplanned
Readmissions After Cardiac Surgery**

Masato Ogawa,PT,Ph.D,^{a,b} Seimi Satomi-Kobayashi,MD,Ph.D,^c Naofumi
Yoshida,MD,Ph.D,^c Yasunori Tsuboi,PT,Ph.D,^a Kodai Komaki,PT^a, Kumiko
Wakida,RD,MSc,^d Yasuko Gotake, MD,^f Kazuhiro P. Izawa, PT,Ph.D,^b Yoshitada
Sakai,MD,Ph.D,^g Kenji Okada, MD,Ph.D,^f

^aDivision of Rehabilitation Medicine, Kobe University Hospital, Kobe, Japan

^bDepartment of Public Health, Kobe University Graduate School of Health Sciences,
Kobe, Japan

^cDivision of Cardiovascular Medicine, Department of Internal Medicine, Kobe
University Graduate School of Medicine, Kobe, Japan

^dDepartment of Nutrition, Kobe University Hospital, Kobe, Japan

^eDepartment of Nursing, Kobe University Hospital, Kobe, Japan

^fDivision of Cardiovascular Surgery, Department of Surgery, Kobe University Graduate
School of Medicine, Kobe, Japan

^gDivision of Rehabilitation Medicine, Kobe University Graduate School of Medicine,
Kobe, Japan

Corresponding Author: Seimi Satomi-Kobayashi, MD, PhD

Division of Cardiovascular Medicine, Department of Internal Medicine
Kobe University Graduate School of Medicine

Kobe, Japan 7-5-2, Kusunoki-cho, Chuo-ku, Kobe, Hyogo 650-0017, Japan

25 Tel: +81-78-382-5846

26 Fax: +81-382-6499

27 E-mail: seimik@med.kobe-u.ac.jp

28 Word count: 3415 words

29

30 **Funding statement**

31 JSPS KAKENHI supported this study [grant numbers JP17K01500, 18K17669, and
32 JP16K01819].

33

34 **Conflict of interest.**

35 None declared.

36

37 **Abbreviations and Acronyms**

38 AF = atrial fibrillation

39 BMI = body mass index

40 CI = confidence interval

41 HR = hazard ratio

42 QOL = quality of life

43 Conv. CR = conventional exercise-based cardiac rehabilitation

44 Multi. CR = multidisciplinary cardiac rehabilitation

45

46 **Central Picture Legend.**

47 Multidisciplinary cardiac rehabilitation improves unplanned readmission-free survival.

48

49 **Central message**

50 Comprehensive multidisciplinary cardiac rehabilitation during the acute inpatient phase

51 after cardiac surgery is important for patient management and preventing unplanned

52 readmission.

53

54 **Perspective statement**

55 Cardiac rehabilitation after cardiac surgery has been underutilized worldwide.

56 Multidisciplinary cardiac rehabilitation focused on patient education and disease self-

57 management during the early postoperative stage can reduce subsequent

58 hospitalizations. These data suggest the importance of acute phase cardiac rehabilitation
59 and the continuity of rehabilitative care after cardiac surgery.

60

Abstract

Objectives: The provision of inpatient programs that reduce the incidence of readmission after cardiac surgery remains challenging. Investigators have focused on multidisciplinary cardiac rehabilitation (CR) because it reduces the postoperative readmission rate; However, most previous studies utilized outpatient models (phase II CR). We retrospectively investigated the effect of comprehensive multidisciplinary interventions in the acute inpatient phase (phase I CR) on unplanned hospital readmission.

Methods: In a retrospective cohort study, we compared consecutive patients after cardiac surgery. We divided them into the multidisciplinary CR (multi. CR) group or conventional exercise-based CR (conv. CR) group according to their postoperative intervention during phase I CR. Multi. CR included psychological and educational intervention and individualized counseling in addition to conv. CR. The primary outcome was unplanned readmission rates between the groups. A propensity score-matching analysis was performed to minimize selection biases and the differences in clinical characteristics.

Results: In our cohort (n=341), 56 (18.3 %) patients had unplanned readmission during the follow-up period (median, 419 days). Compared to the conv. CR group, the multi. CR group had a significantly lower unplanned readmission rate (multivariable regression analysis; hazard ratio, 0.520; 95% confidence interval, 0.28-0.95; p=0.024). A Kaplan-Meier analysis of our propensity score-matched cohort showed that, compared to the conv. CR group, the multi. CR group had a significantly lower incidence of readmission (stratified log-rank test, p=0.041).

Conclusions: In phase I, compared to conv. CR alone, multi. CR reduced the incidence

85 of unplanned readmission. Early multidisciplinary CR can reduce hospitalizations and
86 improve long-term prognosis after cardiac surgery.

87

88 **Abstract Word Count: 250 words**

Introduction

In recent decades, the increased use of fast-track recovery protocols after cardiac surgery has significantly reduced the postoperative hospital length-of-stay and cost of care without increasing postoperative mortality or morbidity [1, 2]. Ironically, because many patients are discharged without an adequate assessment of residual cardiovascular risks, medication titration, physical rehabilitation, or educational intervention, these protocols also increased the rate of unplanned postoperative hospital readmission [3, 4]. Notably, previous studies showed that approximately 20% of patients who underwent cardiac surgery were readmitted within 30 days after discharge resulting in higher medical costs [4, 5].

Current clinical guidelines address this issue by emphasizing the importance of exercise-based cardiac rehabilitation (CR), which effectively improves symptoms and exercise capacity; consequently, it is expected to decrease the rate of readmission [6]. Currently, in addition to conventional exercise-based CR, comprehensive multidisciplinary interventions are recommended to address the impaired quality of life, depressive symptoms, and other disease-related problems that can occur after cardiac surgery. Multidisciplinary CR includes dedicated care with psycho-educational components, nutritional support, and exercise training. Several randomized controlled trials have identified lifestyle interventions and educational programs that improve health outcomes and reduce the risk of a new cardiac event after cardiac surgery by modifying unhealthy behaviors [7, 8]. However, almost all of these CR interventions were carried out during phase II, which begins after discharge. Considering that re-hospitalization is reached approximately 20% within 30 days after discharge, we believe that it is very important to intervene the patients during an earlier phase. However, to

our knowledge, very few investigators have evaluated the impact of comprehensive multidisciplinary CR administered in the hospital immediately after cardiac surgery (phase I) on readmission rates.

Therefore, we investigated the effects of comprehensive multidisciplinary phase I CR interventions, including educational programs designed to help patients self-manage their disease. We aimed to evaluate whether comprehensive multidisciplinary phase I CR can improve cardiovascular event prevention and reduce unplanned hospital readmissions compared to conventional exercised-based CR alone.

Materials and methods

Study Population

The present study was a retrospective analysis conducted at a single center. The study flow chart was shown in Fig. 1. We retrospectively compared consecutive patients who underwent elective cardiac surgery (e.g., coronary artery bypass surgery [CABG], valve replacement or repair, or CABG with concomitant valve replacement or repair) between December 2015 and April 2017 according to the postoperative CR intervention they had received during phase I CR. We excluded patients with a psychiatric disorder or severe dementia and those who could not complete the CR program due to postoperative complications such as respiratory failure, stroke, or hospital death. The study complied with the principles of the Declaration of Helsinki regarding investigations in human subjects and was approved by the Kobe University Institutional Review Board (Approval No.180182). Written informed consent was obtained from each patient before their cardiac surgery.

Patient Clinical Characteristics

We evaluated baseline characteristics including age; sex; body mass index (BMI); left ventricular ejection fraction; estimated glomerular filtration rate; brain natriuretic peptide, hemoglobin, serum albumin, and C-reactive protein levels; comorbidities; medications; and the European System for Cardiac Operative Risk Evaluation (EuroSCORE) II [9]. Laboratory data were evaluated within one week before cardiac surgery. We recorded operative variables including the type and duration (in minutes) of cardiac surgery and postoperative variables including type of phase I CR, hospital mortality, length of intensive care unit stay, length of hospital stay, and postoperative surgery-related complications. Health-related quality of life (QOL) was investigated with the Japanese version of the Medical Outcomes Study 36-Item Short-Form General Health Survey Version 2.0 (SF-36; v2) [10]. The primary endpoint was unplanned hospital readmission after discharge during the follow-up period [11]. We defined unplanned readmission as any type of emergency re-admission (including emergency fast-track or an urgent admission requested by the general practitioner). Planned readmissions were excluded, which were defined as readmissions that were classified as elective. The patients were followed-up as outpatients, and the date and cause of any reported event were determined during regularly scheduled outpatient visits and confirmed by review of hospital medical records.

Intervention for Postoperative Rehabilitation

All patients received CR beginning the day after surgery until hospital discharge per the Japanese Circulation Society guidelines for rehabilitation of patients with cardiovascular disease as a standard of care [6], but none of the patients received phase II CR after

discharge. We divided them into the multidisciplinary CR (multi. CR) group or conventional exercise-based CR (conv. CR) group according to the postoperative intervention they had received during phase I CR. The multidisciplinary rehabilitation team consisted of dedicated medical practitioners focused on improving care for elderly patients. We implemented a twice weekly multidisciplinary conference to review patients scheduled for cardiac surgery. Our team recruited consenting patients who had a great need for psychoeducational intervention, including those with many comorbidities and unhealthy lifestyle habits. Patients who did not require this treatment based on our team assessment or those who did not consent to multi. CR received conv. CR.

Exercise-based Rehabilitation Program (Conv.CR)

CR programs were implemented during the inpatient period for approximately two weeks and were based on Japanese Circulation Society guidelines [6]. CR mainly focused on exercise training and the prevention of postoperative complications. Physical training included a half hour of personalized aerobic exercise each weekday and daily resistance exercise for approximately one hour. Aerobic exercise sessions involved monitored use of a cycle ergometer or treadmill walking at the intensity of 11 to 13 on the Borg scale. The heart rate and oxygen saturation were measured at each session, and patients used pulse watches during cycle training. Muscle and endurance exercises consisted of sit-to-stand and heel lifting exercises with an increasing number of repetitions. The in-hospital physical training was supervised by a physiotherapist. Because patients had undergone a sternotomy, we did not prescribe upper body strength training to avoid complications such as an unstable sternum and exercises that primarily targeted lower body muscles. If necessary, patients underwent respiratory physiotherapy

consisting of deep breathing exercises.

Multidisciplinary Rehabilitation Program (Multi. CR)

Multi. CR consisted of twice-weekly group education and discussion sessions and individualized counseling in addition to conv. CR. Multi. CR focused on psycho-educational intervention to improve patients' coping strategies, improve disease management, provide information, promote self-monitoring of their heart failure symptoms, and help resume daily life after cardiac surgery. Each educational and counseling session lasted approximately one hour. This component included classes conducted by a cardiologist, cardiology nurses, and a nutritionist. These sessions aimed to change habits that adversely affect the risk of cardiovascular diseases through the use of coping strategies and disease management, provide information, and help patients resume daily activities after cardiac surgery. CR personnel also helped patients incorporate self-care behaviors into their daily routine. The educational program was offered to patients' caregivers as well.

Professional nursing education included teaching patients how to manage their heart failure symptoms, measure their pulse, recognize an arrhythmia or infection, monitor their weight and surgical sites, and optimize their cardioprotective medications. Nurses also provided information about smoking cessation, coping with stress, and daily physical activity.

A nutritionist identified comorbidities including chronic kidney disease, obesity, hypertension, diabetes, and dyslipidemia, provided appropriate meals in the hospital, and explained the necessity of continuing this diet after discharge. The nutritionist also provided education about water management, salt restriction, and daily alcohol

consumption.

Statistical Analysis

We conducted statistical analyses after confirming that the data were normally distributed using the Shapiro-Wilk test. Patients were separated into the conv. CR or multi. CR group, and between-group differences in clinical characteristics were compared using the independent t- and chi-squared tests. We adjusted for baseline characteristics using propensity scores to reduce the risk of bias in treatment selection and other potential confounding factors because patients were not randomly assigned to a CR group [12]. To produce propensity scores, a logistic regression analysis using the 22 variables which all variables listed in Table 1 was performed (c statistic = 0.721). We performed a 1:1 nearest available matching on the logit of the propensity score with a caliper value of 0.2 and no replacement. We used the standardized mean difference to measure covariate balance, whereby an absolute standardized mean difference <0.1 represents a meaningful imbalance between groups. In the matched cohort, paired comparisons were performed with the use of McNemar's test for binary variables and a paired Student's t-test for continuous variables. Association between CR modality and unplanned readmission was evaluated by using a proportional hazards regression model stratified on matched pairs to preserve the benefit of matching. Unplanned readmission was compared using the Fine-Gray model, which adjusted for death as a competing risk [13]. Proportional hazard assumption was checked both graphically and using the Schoenfeld residual test. In the entire cohort, the multivariable proportional-hazards regression model with the stepwise backwards (Wald) method from factors with p-values <0.10 in the univariable analysis of the entire cohort was performed to calculate hazard ratios (HR) and the impact of CR on

unplanned readmission. We estimated overall unplanned readmission-free survival between conventional CR group and multidisciplinary CR group by using Kaplan-Meier curves for all patients and a propensity score-matched cohort. In the matched cohort, we compared overall re-admission free survival in both groups by using the stratified log-rank test. The sample size was calculated with referring to the previous paper [14] and our unpublished data (power = 0.8, significance level = 0.05, mean difference = 14.0%; n = 300 patients). A p-value <0.05 was considered statistically significant. Statistical analyses were carried out using JMP 11.0J software (SAS Institute Inc., Cary, NC, US).

Results

During the study period, 341 patients presented for surgery. Of these, 35 were excluded (Figure 1) and 306 (mean age, 67.5±13.1 years; median follow-up, 419 days [interquartile range, 211-712]) were analyzed. There were 180 and 126 patients in the conv. CR and multi. CR groups, respectively. Propensity score-matching identified 216 matched pairs for comparison (c statistic =0.721; Table 1, Figure S1). No residual imbalance was observed between matched groups (p>0.10 for all variables). Furthermore, the covariate balance in the matched cohort was considerably improved; the absolute standardized mean difference was <0.1. After propensity score-matching, the follow-up periods in the multi. CR and conv. CR groups were comparable (458.5±296.9 days vs. 448.2±267.0 days).

Table 2 compared the clinical results and incidence of unplanned readmission before and after matching. Length of hospital stay and QOL at hospital discharge were not significantly different between the two groups (Table S1). Overall, 56 patients (18.3%) had unplanned readmission (heart failure, n=31; infection, n=11; arrhythmia, n=8;

stroke, n=3; pneumonia, n=2; nonfatal myocardial infarction, n=1). After propensity score matching, the incidence of unplanned readmission was significantly lower in the multi. CR group than in the conv. CR group ($p=0.0148$). A global test based on Schoenfeld residuals found that all covariates and the full model satisfied the proportional hazard assumption. A univariable proportional hazards analysis for predicting unplanned readmission in the entire cohort showed no significant difference between the CR groups ($p=0.174$). However, multivariable regression analyses of unmatched patients showed that compared to conv. CR, multi. CR independently predicted a decreased rate of unplanned readmission (HR, 0.504; 95% CI, 0.282-0.901; $p=0.021$; Table 3). Similarly, multi. CR independently predicted a reduced rate of unplanned readmission in our comparison of the propensity score-matched cohorts (HR, 0.520; 95% CI, 0.277-0.949; $p=0.024$; Table 4). The results of the Kaplan-Meier analysis were shown Figure 2 and Figure S2. The cumulative incidence of unplanned readmission for the multi. CR group was significantly lower than that for the conv. CR group in propensity score-matched cohorts (stratified log-rank test, $p=0.0413$; HR, 0.55, 95%CI, 0.31-0.96; Figure 2).

Discussion

To the best of our knowledge, this is the first study to demonstrate the effectiveness of comprehensive multidisciplinary phase I CR after cardiac surgery. We showed that our comprehensive multi. CR program was independently associated with a significant reduction in postoperative unplanned readmission compared to conv. CR alone after adjusting for many confounding variables. Several investigators have reported the beneficial effects of multi. CR. However, these studies were of phase II CR. We are the

first to demonstrate that phase I multi. CR can effectively reduce the incidence of long-term adverse outcomes. Furthermore, few studies have compared the effects of exercise-based and comprehensive multi. CR. The aim of phase I CR is to minimize the effects of inactivity and maintain or improve muscle strength and mobility [6]. Exercise-based phase I CR effectively improves exercise capacity and prevents cardiac events after cardiac surgery [3, 14, 15]. Notably, our study demonstrated that multidisciplinary educational intervention in addition to exercise-based CR were more effective than exercise-based CR alone even during phase I.

In our cohort, the patients in the multi. CR group had many comorbidities and high surgical risk as shown by their EuroSCOREs. Our selection method, which involved intentionally recruiting high-risk patients into the multi. CR group, may explain the differences between our two study groups. We did not observe a statistically significant reduction in hospital readmission between the conv. CR and multi. CR groups.

Nevertheless, it is interesting to note that, when we used a propensity score matching method to reduce the confounding effect due to differences in demographics between the two groups, multi. CR was independently associated with a significant reduction in unplanned readmission compared to conv. CR. Our results suggested that multi. CR is useful regardless of the patient's background and comorbidities. Previous studies of phase II CR found that patients with an adverse risk factor profile or a poor understanding of risk factors are not likely to attend CR sessions and have low adherence to CR [16]. Furthermore, the CR participation rate is lower in proportion to the delay in CR enrollment after discharge [17, 18]. Phase I CR is conducted in the hospital; Thus, the participation rate tends to be higher than that of phase II CR.

Therefore, we believe that phase I is an opportune time to begin changing lifestyle

habits and improving physical activity in patients at high risk for readmission.

In this study, the incidence of unplanned readmission was 18.3%, which is similar to that of previous studies carried out after cardiac surgery [4, 19]. More than half of these readmissions were for worsening heart failure, often accompanied by volume overload. Previous studies found that most readmissions for heart failure exacerbation are attributable, at least in part, to poor self-care, including non-compliance with medication and diet recommendations and failure to act upon escalating symptoms [20]. Moreover, most heart failure readmissions are caused by factors that patients can be taught to recognize and avoid [21]. We used a wide variety of multidisciplinary strategies focused on patient education and self-management to reduce hospitalizations. Furthermore, it is highly likely that the effects of our multidisciplinary phase I CR educational program persisted long after discharge. A review of interventions promoting self-care in patients with heart failure revealed that specific components, including group-based programs and frequent contact and supervision by researchers or health team members, appear to promote positive outcomes [22]. Thus, phase I is a critical window of opportunity for initiating disease self-management education, especially for high-risk patients.

The second most common cause of readmission was infection, including superficial and deep sternotomy infections. Surgical site infections require meticulous wound care and antibiotics and are complications that could be prevented through quality improvement initiatives. Arrhythmia was also a common cause of readmission.

Postoperative atrial fibrillation (AF) occurred frequently, and the risk of AF persisted after discharge. Reportedly, a nurse-led education program that teaches patients how to self-palpate their pulse may be a useful method for screening asymptomatic AF.

However, the effect of self-pulse palpation on long-term cardiac events remains poorly

understood [23]. Future studies of postoperative AF management after discharge are warranted. Although we could not analyze what component of multi. CR was effective, we think multidisciplinary CR education program for patients and their caregivers effectively promoted heart failure symptom monitoring and daily infection surveillance. In particular, we believe that it is most important to educate patients about self-monitoring and self-management of heart failure symptom by nutritionist or professional nurses.

Recent guidelines for cardiac rehabilitation include a Class I recommendation for patients to receive specific education about heart failure self-care [6]. Nevertheless, phase II CR is underutilized worldwide. Phase II CR was only provided to 30% of patients in Europe, 25% in the United States, and 9% in Japan [24, 25]. Institutional practices, health care system practices, and health insurance system practices could explain this low rate of participation. The under-utilization of phase II CR suggests there is a problem with the continuity of care from inpatient to outpatient rehabilitation. As the rate of phase II CR participation is very low (including formulated exercise prescriptions and patient education programs), phase I CR is of substantial importance to patient management. A recent study demonstrated that patient compliance with phase II CR might be increased with home-based CR or telemonitoring [26]. We believe that CR including patient education and disease management and close cooperation with general practitioner from the early postoperative period may be effective for preventing hospital readmission after cardiac surgery. Future studies investigating the transition from phase I to phase II CR are needed to improve the continuity of rehabilitative care after cardiac surgery.

Our study had several limitations. Firstly, the study was a retrospective analysis, and

no contemporary control group was available for comparison. Therefore, a type 2 statistical error cannot be excluded because of low statistical power to detect a difference between the two groups. However, patients included in our study population could represent a more representative sample of “real world” patients undergoing cardiac surgery than those included in many of the randomized controlled trials that have previously evaluated CR effectiveness. Secondly, our recruitment method may have introduced selection bias, although several different statistical methods were used to adjust for baseline characteristics and confirm the results. There was a possibility that systematic error, bias or hidden confounders, and not true biological effects could occur. Thirdly, we did not adjust for all confounding factors such as patient personality, motivation, or attitude toward CR. Our results may not be generalizable to patients who choose not to enroll in CR programs. Next, we did not have data about cost of CR or the outcomes of long-term QOL. Therefore, we cannot mention whether multidisciplinary CR programs are cost-effective. Furthermore, we could not continue phase II CR after discharge or investigate the effects of general practitioner after discharge and the frequency of intervention in detail. Finally, the lack of complete socioeconomic data, including education and income levels, also limited the study.

In conclusion, this study of the effects of phase I CR after cardiac surgery revealed that, compared to conv. CR alone, comprehensive multi. CR reduced the incidence of unplanned readmission after adjusting for confounding factors. Multi. CR focused on patient education and disease self-management in the early postoperative stage can reduce subsequent hospitalizations. Future studies investigating the clinical effects of phase I CR on long-term outcomes and the transitions between CR phases are needed to improve the care of patients undergoing cardiac surgery.

377

378

Acknowledgments

379

We thank Ms. Hikaru Nishida, Ms. Kazumi Takahama, Ms Keiko Uga, Ms. Junko

380

Uchida, and Prof. Ken-ichi Hirata for their expert assistance in the primary care setting

381

and our institutional colleagues for contributing to the care of our patients.

382

383

Figure Legends

384

Figure. 1 Study flow chart of this study

385

Conv. CR, conventional exercise-based cardiac rehabilitation; Multi. CR,

386

multidisciplinary cardiac rehabilitation

387

388

Figure. 2 The Kaplan–Meier survival curves for cumulative unplanned readmission

389

after cardiac surgery between the conv. CR group (red line) and multi. CR groups (blue

390

line) in a propensity score-matching cohort. The incidence of unplanned readmission in

391

the multi. CR group was significantly lower than that in the conv. CR group (stratified

392

log-rank test, $p=0.0413$). We also revealed confidence limits as the shaded area.

393

Conv. CR, conventional exercise-based cardiac rehabilitation; Multi. CR,

394

multidisciplinary cardiac rehabilitation

395

396

Figure S1. Comparison of propensity score distributions before and after matching

397

between the conventional and multidisciplinary cardiac rehabilitation groups

398

399

Figure S2. The Kaplan–Meier survival curves for cumulative unplanned readmission

400

after cardiac surgery between the conv. CR group (red line) and multi. CR groups (blue

line) in the entire cohort. The incidence of unplanned readmission was not significantly different between the multi. CR group and the conv. CR group (log-rank test, $p=0.0845$). We also revealed confidence limits as a shaded area. Conv. CR, conventional exercise-based cardiac rehabilitation; Multi. CR, multidisciplinary cardiac rehabilitation

Video Legends

Dr Ogawa, the first author and principal statistician of this study, explains the background, major findings, and key relevance of the present study.

References

- [1] Celkan MA, Ustunsoy H, Daglar B, Kazaz H, Kocoglu H. Readmission and mortality in patients undergoing off-pump coronary artery bypass surgery with fast-track recovery protocol. *Heart Vessels* 2005; 20: 251-5.
- [2] Yanatori M, Tomita S, Miura Y, Ueno Y. Feasibility of the fast-track recovery program after cardiac surgery in Japan. *Gen Thorac Cardiovasc Surg* 2007; 55: 445-9.
- [3] Takahashi T, Kumamaru M, Jenkins S, Saitoh M, Morisawa T, Matsuda H. In-patient step count predicts re-hospitalization after cardiac surgery. *J Cardiol* 2015; 66: 286-91.
- [4] Iribarne A, Chang H, Alexander JH, Gillinov AM, Moquete E, Puskas JD et al. Readmissions after cardiac surgery: experience of the National Institutes of Health/Canadian Institutes of Health research cardiothoracic surgical trials network. *Ann Thorac Surg* 2014; 98: 1274-80.
- [5] Hannan EL, Zhong Y, Lahey SJ, Culliford AT, Gold JP, Smith CR et al. 30-day readmissions after coronary artery bypass graft surgery in New York State. *JACC Cardiovasc Interv* 2011; 4: 569-76.
- [6] JCS Joint Working Group. Guidelines for rehabilitation in patients with cardiovascular disease (JCS 2012). *Circ J* 2014; 78: 2022-93.
- [7] Kadda O, Kotanidou A, Manginas A, Stavridis G, Nanas S, Panagiotakos DB. Lifestyle intervention and one-year prognosis of patients following open heart surgery: a randomised clinical trial. *J Clin Nurs* 2015; 24: 1611-21.
- [8] Williams JB, DeLong ER, Peterson ED, Dokholyan RS, Ou FS, Ferguson TB Jr. Secondary prevention after coronary artery bypass graft surgery: findings of a

435 national randomized controlled trial and sustained society-led incorporation into
436 practice. *Circulation* 2011; 123: 39-45.

437 [9] Nashef SA, Roques F, Sharples LD, Nilsson J, Smith C, Goldstone AR et al.
438 EuroSCORE II. *Eur J Cardiothorac Surg* 2012; 41: 734-44; discussion 744-5.

439 [10] Fukuhara S, Bito S, Green J, Hsiao A, Kurokawa K. Translation, adaptation, and
440 validation of the SF-36 Health Survey for use in Japan. *J Clin Epidemiol* 1998;
441 51: 1037-44.

442 [11] Saji M, Higuchi R, Tobaru T, Iguchi N, Takanashi S, Takayama M, Isobe M.
443 Impact of Frailty Markers for Unplanned Hospital Readmission Following
444 Transcatheter Aortic Valve Implantation. *Circ J* 2018; 82: 2191-8.

445 [12] Rubin DB. Estimating causal effects from large data sets using propensity scores.
446 *Ann Intern Med* 1997; 127(8 Pt 2): 757-63.

447 [13] Austin PC, Fine JP. Practical recommendations for reporting Fine-Gray model
448 analyses for competing risk data. *Stat Med* 2017; 36: 4391-4400.

449 [14] Ogawa M, Izawa KP, Satomi-Kobayashi S, Tsuboi Y, Komaki K, Gotake Y et al.
450 Impact of delirium on postoperative frailty and long term cardiovascular events
451 after cardiac surgery. *PLoS One* 2017; 12: e0190359.

452 [15] Ramos Dos Santos PM, Aquaroni Ricci N, Aparecida Bordignon Suster E, de
453 Moraes Paisani D, Dias Chiavegato L. Effects of early mobilisation in patients
454 after cardiac surgery: a systematic review. *Physiotherapy* 2017; 103: 1-12.

455 [16] Daly J, Sindone AP, Thompson DR, Hancock K, Chang E, Davidson P. Barriers
456 to participation in and adherence to cardiac rehabilitation programs: a critical
457 literature review. *Prog Cardiovasc Nurs* 2002; 17: 8-17.

458 [17] Russell KL, Holloway TM, Brum M, Caruso V, Chessex C, Grace SL. Cardiac

459 rehabilitation wait times: effect on enrollment. *J Cardiopulm Rehabil Prev* 2011;
 460 31: 373-7.

461 [18] Pack QR, Mansour M, Barboza JS, Hibner BA, Mahan MG, Ehrman JK et al.
 462 An early appointment to outpatient cardiac rehabilitation at hospital discharge
 463 improves attendance at orientation: a randomized, single-blind, controlled trial.
 464 *Circulation* 2013; 127: 349-55.

465 [19] Maniar HS, Bell JM, Moon MR, Meyers BF, Marsala J, Lawton JS et al.
 466 Prospective evaluation of patients readmitted after cardiac surgery: analysis of
 467 outcomes and identification of risk factors. *J Thorac Cardiovasc Surg* 2014;
 468 147:1013-8.

469 [20] Moser DK, Dickson V, Jaarsma T, Lee C, Stromberg A, Riegel B. Role of self-
 470 care in the patient with heart failure. *Curr Cardiol Rep* 2012; 14: 265-75.

471 [21] Tsuyuki RT, McKelvie RS, Arnold JM, Avezum A Jr., Barretto AC, Carvalho AC
 472 et al. Acute precipitants of congestive heart failure exacerbations. *Arch Intern*
 473 *Med* 2001; 161: 2337-42.

474 [22] Barnason S, Zimmerman L, Young L. An integrative review of interventions
 475 promoting self-care of patients with heart failure. *J Clin Nurs* 2012; 21(3-4): 448-
 476 75.

477 [23] Jaakkola J, Vasankari T, Virtanen R, Juhani Airaksinen KE. Reliability of pulse
 478 palpation in the detection of atrial fibrillation in an elderly population. *Scand J*
 479 *Prim Health Care* 2017; 35: 293-8.

480 [24] Suaya JA, Shepard DS, Normand SL, Ades PA, Prottas J, Stason WB. Use of
 481 cardiac rehabilitation by Medicare beneficiaries after myocardial infarction or
 482 coronary bypass surgery. *Circulation* 2007; 116: 1653-62.

- 483 [25] Goto Y, Saito M, Iwasaka T, Daida H, Kohzuki M, Ueshima K et al. Poor
484 implementation of cardiac rehabilitation despite broad dissemination of coronary
485 interventions for acute myocardial infarction in Japan: a nationwide survey. *Circ*
486 *J* 2007; 71: 173-9.
- 487 [26] Piotrowicz E, Baranowski R, Bilinska M, Stepnowska M, Piotrowska M, Wójcik
488 A et al. A new model of home-based telemonitored cardiac rehabilitation in
489 patients with heart failure: effectiveness, quality of life, and adherence. *Eur J*
490 *Heart Fail* 2010; 12: 164-71.

Readmission






Figure. 1




Figure 2





Figure. S1



No. at risk

	0	3	6	9	12	15	18
Conv. CR group	180	157	136	111	96	83	72
Multi. CR group	126	116	98	86	74	63	52

Figure S2

1 Table 1. Baseline clinical characteristics of conventional and multidisciplinary cardiac rehabilitation groups

2

3

Variables	All study patients				Propensity-matched population			
			Standardized				Standardized	
	Conv. CR	Multi. CR	Mean	p-value	Conv. CR	Multi. CR	Mean	p-value
			Differences				Differences	
Number	180	126			108	108		
Age, years	67.0±14.5	67.6±12.9	0.043	0.692	67.4±13.4	67.3±13.2	0.007	0.941
Sex, female (%)	75 (41.7)	49 (38.9)	0.057	0.626	38 (35.2)	43 (39.8)	0.010	0.596
BMI, kg/m ²	22.9±3.7	23.4±4.3	0.124	0.270	23.0±3.8	23.1±4.3	0.024	0.821
Albumin, g/dl	4.0±0.5	4.0±0.5	0.040	0.837	3.9±0.6	4.0±0.5	0.018	0.879

BNP, pg/ml	248.5±382.1	231.7±294.2	0.049	0.677	217.9±347.6	231.0±298.6	0.043	0.778
Hemoglobin, g/dl	12.4±1.9	12.7±1.7	0.166	0.142	12.7±1.9	12.7±1.6	0.001	0.905
LVEF, %	58.5±13.2	60.3±10.7	0.150	0.194	59.5±13.3	59.8±11.0	0.025	0.850
Hypertension, n (%)	82 (45.6)	76 (60.3)	0.297	0.011	62 (57.4)	61 (56.5)	0.018	0.922
Dyslipidemia, n (%)	47 (26.1)	53 (42.1)	0.322	0.003	36 (33.3)	42 (38.9)	0.097	0.391
Diabetes, n (%)	37 (20.6)	41 (32.5)	0.272	0.018	31 (28.7)	30 (27.8)	0.020	0.923
Chronic kidney disease, n (%)	93 (51.7)	57 (45.2)	0.130	0.268	51 (47.2)	52 (48.2)	0.020	0.892
Atrial fibrillation, n (%)	40 (22.2)	39 (31.0)	0.200	0.086	32 (29.6)	29 (26.9)	0.060	0.766

Smoking, n (%)	23(12.8)	16(12.7)	0.003	0.984	13 (12.0)	14 (13.0)	0.030	0.864
Type of surgery, n (%)								
Valve	130 (72.2)	87 (69.1)	0.068	0.880	76 (70.4)	75 (69.4)	0.022	0.955
CABG	16 (8.9)	12 (9.5)			9 (8.3)	11 (10.2)		
Concomitant	34 (18.9)	27 (21.4)			23 (21.3)	22 (20.4)		
NYHA class n (%)								
I	32 (17.8)	28 (22.2)	0.020	0.279	23 (21.3)	22 (20.4)	0.020	0.779
II	91 (50.6)	65 (51.6)			57 (52.8)	56 (51.9)		
III	53 (29.4)	33 (26.2)			28 (25.9)	30 (27.8)		
Duration of surgery,	347.5±110.6	335.6±91.0	0.118	0.322	343.8±102.8	333.2±87.0	0.092	0.481

min

EuroSCORE II	5.9±3.8	6.8±3.7	0.240	0.032	6.6±4.0	6.7±3.1	0.028	0.807
--------------	---------	---------	-------	-------	---------	---------	-------	-------

Medications, n (%)

β-blocker	112 (62.2)	97 (77.0)	0.326	0.006	74 (77.3)	74 (76.3)	0.024	0.784
-----------	------------	-----------	-------	-------	-----------	-----------	-------	-------

ACE-I	38 (21.1)	37 (29.4)	0.192	0.950	22 (22.7)	22 (22.7)	< 0.001	1.000
-------	-----------	-----------	-------	-------	-----------	-----------	---------	-------

ARB	35 (19.4)	45 (35.7)	0.371	0.001	28 (28.9)	26 (26.8)	0.047	0.451
-----	-----------	-----------	-------	-------	-----------	-----------	-------	-------

Statin	50 (27.8)	46 (36.5)	0.187	0.105	30 (30.9)	31 (32.0)	0.024	0.770
--------	-----------	-----------	-------	-------	-----------	-----------	-------	-------

Diuretics	85 (47.2)	66 (52.4)	0.104	0.374	49 (50.5)	47 (48.5)	0.040	0.774
-----------	-----------	-----------	-------	-------	-----------	-----------	-------	-------

4 Conv. CR, conventional exercise-based cardiac rehabilitation; Multi. CR, multidisciplinary cardiac rehabilitation; BMI, body mass index;

5 BNP, brain natriuretic peptide; LVEF, left ventricular ejection fraction; CABG, coronary artery bypass grafting; NYHA, New York Heart

6 Association; ACE-I, angiotensin converting enzyme inhibitor; ARB, angiotensin II receptor blocker

7 Data are expressed as means \pm standard deviation or numbers (percentage)

8

9

10 Table 2. Comparison of clinical outcomes of conventional and multidisciplinary cardiac rehabilitation groups
11

Variables	All study patients				Propensity-matched population			
	Conv. CR	Multi. CR	Standardized		Conv. CR	Multi. CR	Standardized	
			Mean	p-value			Mean	p-
			Differences				Differences	value
Length of hospital stay, days	22.0±21.0	20.0±10.2	0.135	0.158	21.8±24.3	20.1±10.8	0.090	0.240
Follow up, days	453.4±304.7	405.2±269.8	0.167	0.959	458.5±296.9	448.2±267.0	0.364	0.791
Unplanned readmission, n (%)	39 (21.7)	17(13.5)	0.217	0.065	28 (25.9)	13 (12.0)	0.360	0.0148

Death, n (%)	1 (0.6)	1 (0.8)	0.024	0.799	0 (0.0)	1 (1.0)	0.045	0.222
--------------	---------	---------	-------	-------	---------	---------	-------	-------

12 Conv. CR, conventional exercise-based cardiac rehabilitation; Multi. CR, multidisciplinary cardiac rehabilitation

Table 3. Fine-Gray proportional hazard regression for readmission-free survival after multidisciplinary cardiac rehabilitation for all patients (n=306)

Model	Hazard ratio	95% CI	p-value	Proportional hazard assumption p value
Unadjusted	0.671	0.377-1.193	0.174	0.155
Adjusted for age and sex	0.650	0.365-1.157	0.143	0.232
Multivariable adjusted*	0.504	0.282-0.901	0.0207	0.180

Reference: conventional cardiac rehabilitation group

CI, confidence interval

*Adjusted for age, hypertension, diabetes, dyslipidemia

Table 4. Fine-Gray proportional hazard regression for readmission-free survival after multidisciplinary cardiac rehabilitation for propensity score-matched patients (n=216)

Model	Hazard ratio	95% CI	p-value	Proportional hazard assumption p value
Unadjusted	0.552	0.283-0.987	0.040	0.695
Adjusted for propensity	0.520	0.277-0.949	0.0235	0.722

Reference: conventional cardiac rehabilitation group

CI, confidence interval

1 Table S1. Comparison of health-related quality of life at hospital discharge after surgery in conventional and multidisciplinary cardiac
2 rehabilitation groups

Variables	All study patients				Propensity-matched population			
	Conv. CR	Multi. CR	Standardized		Conv. CR	Multi. CR	Standardized	
			Mean	p-value			Mean	p-value
Differences	Differences							
PCS	38.7±12.6	39.0±13.2	2.325	0.910	38.7±12.8	40.1±13.1	10.810	0.646
MCS	51.9±11.3	53.9±9.2	19.410	0.412	51.8±11.1	53.6±9.6	17.346	0.480

3 Conv. CR, conventional exercise-based cardiac rehabilitation; Multi. CR, multidisciplinary cardiac rehabilitation; PCS, Physical
4 component summary; MCS, Mental component summary