

PDF issue: 2025-05-14

FGF7 supports hematopoietic stem and progenitor cells and niche-dependent myeloblastoma cells via autocrine action on bone marrow stromal cells in vitro

石野, 瑠璃

<mark>(Degree)</mark> 博士(保健学)

(Date of Degree) 2014-03-25

(Date of Publication) 2015-03-01

(Resource Type) doctoral thesis

(Report Number) 甲第6175号

(URL) https://hdl.handle.net/20.500.14094/D1006175

※ 当コンテンツは神戸大学の学術成果です。無断複製・不正使用等を禁じます。著作権法で認められている範囲内で、適切にご利用ください。





# FGF7 supports hematopoietic stem and progenitor cells and niche-dependent myeloblastoma cells via autocrine action on bone marrow stromal cells in vitro

FGF1 and FGF2 support hematopoietic stem and progenitor cells (HSPCs) under stress conditions. In this study, we show that fibroblast growth factor (FGF7) may be a novel niche factor for HSPC support and leukemic growth. FGF7 expression was attenuated in mouse embryonic fibroblasts (MEFs) deficient for the MED1 subunit of the Mediator transcriptional coregulator complex. When normal mouse bone marrow (BM) cells were cocultured with *Med1*<sup>+/+</sup> MEFs or BM stromal cells in the presence of anti-FGF7 antibody, the growth of BM cells and the number of long-time culture-initiating cells (LTC-ICs) decreased significantly. Anti-FGF7 antibody also attenuated the proliferation and cobblestone formation of MB1 stromal cell-dependent myeloblastoma cells. The addition of recombinant FGF7 to the coculture of BM cells and *Med1*<sup>-/-</sup> MEFs increased BM cells and LTC-ICs. FGF7 and its cognate receptor, FGFR2IIIb, were undetectable in BM cells, but MEFs and BM stromal cells expressed both. FGF7 activated downstream targets of FGFR2IIIb in *Med1*<sup>+/+</sup> and *Med1*<sup>-/-</sup> MEFs and BM stromal cells. Taken together, we propose that FGF7 supports HSPCs and leukemia-initiating cells indirectly via FGFR2IIIb expressed on stromal cells.

Keywords: FGF7, hematopoietic niche, bone marrow stromal cells, MED1, Mediator

## Introduction

Postnatal mammalian hematopoietic stem and progenitor cells (HSPCs) are strictly regulated by specific microenvironmental niches in the bone marrow (BM). BM mesenchymal stem cells (MSCs) and their derived or related cells constitute important cellular components of the hematopoietic niche [reviewed in 1-3]. The endosteal region of the BM was first identified as the resting HSPC reservoir, where osteoblasts of mesenchymal origin primarily play a niche function. Meanwhile, recent studies have highlighted the BM sinusoidal vasculature and perivascular region, which are composed of MSCs or related stromal cells (Nestin<sup>+</sup> MSCs, CXC chemokine ligand 12 (CXCL12)-abundant reticular cells, pericytes, and endothelial cells), as a major niche for HSPCs [4-6, reviewed in 1-3]. Molecular components of the HSPC niche include angiopoietin-1 (Ang-1), CXCL12, Wnt signaling, and N-cadherin, which in concert control HSPC behavior [reviewed in 1-3]. We have also reported osteopontin (OPN), produced by stromal cells, as an important niche factor [7].

Fibroblast (FGFs) growth factors comprise а large family of heparin-binding growth factors that regulate cell migration, growth, and differentiation/fate. A family of four receptor tyrosine kinases, designated FGF

receptor (FGFR) 1 through FGFR4, acts as the specific receptors for FGFs to initiate intracellular signaling [reviewed in 8,9]. FGFR1 and FGF2 appear to be dispensable for homeostatic hematopoiesis [10,11]. However, in response to stimulatory stresses to HSPCs, FGFR1 expressed on HSPCs reportedly promotes HSPC expansion through FGF1 (and possibly FGF4) produced by megakaryocytes [10], while FGF2 produced by megakaryocytes indirectly promotes HSPC expansion via the proliferation of stromal cells, which in turn change the levels of niche signals [11]. In this manner, some FGF signaling appears to be a key regulator of HSPC proliferation and recovery in stressed environments.

FGF7 (also known as keratinocyte growth factor). produced by mesenchymal cells, binds to and activates the specific receptor tyrosine kinase FGFR2IIIb, which in turn rapidly activates Ras-mitogen-activated protein kinase (MAPK) and phosphoinositide 3-kinase-Akt signaling pathways. FGFR2IIIb, expressed mainly on epithelial cells, functions in epithelial cell wound healing, growth, and hair development [reviewed in 8,9]. Upon hepatic injury, Thy1<sup>+</sup> mesenchymal cells produce FGF7 in close proximity to liver progenitor cells, which subsequently proliferate contribute liver and to

regeneration. Thus, FGF7 appears to be a functional niche signal for the stimulation of liver progenitor cells [12]. However, the role of FGF7 in HSPC maintenance is not known.

The Mediator transcriptional coregulator complex, а subcomplex of RNA polymerase II holoenzyme, acts as the end-point integrator of a variety of activators and conveysintracellular signals to the general transcription machinery. Among circa 30 subunits, MED1 is crucial for various biological events through its specific interaction with distinct activators, such as nuclear receptors [reviewed in 13]. In this study, we show that FGF7 expression was profoundly attenuated in Med1<sup>-/-</sup> mouse embryonic fibroblasts (MEFs), that FGF7 produced by MEFs and BM stromal cells is a crucial niche factor for HSPC support and leukemic growth in vitro, and that FGF7 acts indirectly by stimulating stromal cells in an autocrine manner.

#### Materials and methods

### <u>Cell culture</u>

Stable lines of  $Med1^{+/+}p53^{-/-}$  and  $Med1^{-/-}p53^{-/-}$  MEFs established from embryonic day 10.0 (E10.0) embryos derived from a single crossing of  $Med1^{+/-}p53^{-/-}$  mice on a C57BL6 background were described previously [7]. Two lines of these MEFs were analyzed in all experiments. The MEFs

were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) at 37°C. The OP-9 BM stromal cells, distributed by RIKEN BRC through the National Bio-Resource of the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT), MC-3T3-E1 cells, and MS-5 cells [14] were maintained in  $\alpha$ -modified Eagle's medium ( $\alpha$ MEM) supplemented with 20% FBS, 10% FBS, and 20% horse serum, respectively. MB-1 stromal cell-dependent myeloblastic leukemia cells were maintained by coculture with mitomycin C-treated OP-9 cells in  $\alpha$ MEM supplemented with 10% FBS, 1× non-essential amino acids, and 100 µM  $\beta$ -mercaptoethanol [15,16].

<u>BM culture and colony-forming cell assay</u> MEFs and MS-5 and OP-9 cells  $(2 \times 10^5)$ , treated with 100 µg/mL of mitomycin C to arrest the cell cycle, were plated on 0.1% gelatin-coated 12-well plates. The next day,  $1 \times 10^6$  BM cells, harvested from the femurs of congenic wild-type mice, were added to each well and cultured in MyeloCult M5300 (Stem Cell Technologies, Canada) and 10 U/mL heparin in the absence or presence of various amounts of recombinant human (rh) FGF7 (R&D Systems), 0.2 µg/mL anti-mouse (m) FGF7 rabbit polyclonal IgG (H-73: sc-7882; Santa Cruz), or normal rabbit IgG (Sigma) at 33°C.

For long-term culture, half of the medium was replaced with fresh medium each week. After an 8-week culture period, trypsinized cells containing HSPCs (adherent and nonadherent) were collected and cultured in complete methylcellulose medium (MethoCult M3434; Stem Cell Technologies) for all types of colonies at 37°C for 14 days, and the colonies were counted.

#### DNA synthesis

Incorporation of bromodeoxyuridine (BrdU) into cells cultured in 24-well plates, after purging for 6 h, was measured by using a cell proliferation enzyme-linked immunosorbent assay (ELISA) for BrdU (chemiluminescence) (Roche).

## Quantitation of mRNA

For quantitative PCR (qPCR), total RNA  $(0.5 \mu g)$  was used to prepare cDNA with the ReverTra Ace qPCR RT Master Mix with a gDNA Remover kit (Toyobo). The expression of various mouse genes was identified by qPCR (StepOnePlus Real-Time PCR Life system; Technologies). Mouse glyceraldehyde-3-phosphate

dehydrogenase (GAPDH) was used as the reference marker. The sequences of the primers used and the PCR conditions for amplification are available upon request.

## Western blot analysis

For western blot analysis, total cell lysates were separated by SDS-PAGE, blotted onto a nitrocellulose membrane, and probed with polyclonal antibodies against mFGF7 (above), β-actin (BioLegend), FGFR substrate 2 (FRS2) (H-91: sc-8318; Santa Cruz), Phospho-FRS2- $\alpha$ (Tyr196) (Cell Signaling), mitogen-activated protein (MAP) kinase ERK1/ERK2 (Millipore), and phospho-MAP kinase ERK1/ERK2 (Tyr202/204) (Cell Signaling) and with monoclonal antibody against mFGFR2(IIIb) (133730; R&D Systems). Chemiluminescence was detected by an ImageQuant LAS 4000mini (GE Healthcare).

## Statistical analysis

All assays were performed in triplicate or quadruplicate in at least three separate experiments. The significance of the differences between independent means was assessed by Student's t-test. We considered a P value of <0.05 statistically s i g n i f i c a n t .

### Results

### FGF7 expression is attenuated in

## <u>Med1<sup>-/-</sup> MEFs</u>

MEFs, primitive mesenchymal cells resembling MSCs, support HSPCs and thus mimic the hematopoietic niche [7].

We employed MEFs as an in vitro model to analyze the role of MED1 in the BM niche and previously reported that the ability of  $Med1^{-/-}$  MEFs to promote BM cell proliferation and support HSPCs is reduced relative to that of  $Med1^{+/+}$  MEFs [7]. Therefore, MED1 in MEFs has a crucial role in activating the transcription of genes encoding molecules with potency for HSPC support. Microarray analysis comparing  $Med1^{+/+}p53^{-/-}$  and  $Med1^{-/-}p53^{-/-}$  MEFs revealed reduced expression of Fgf7 and Opn in  $Med1^{+/+}p53^{-/-}$  MEFs (GEO accession number GSE22471) [7].

RT-qPCR (Fig. 1A) and western blot (Fig. 1B) analyses confirmed the reduced level of FGF7 in  $Med1^{-/-}$  MEFs. However, Fgf7 expression was not restored after MED1 was reintroduced into  $Med1^{-/-}$  MEFs (Rev- $Med1^{-/-}$  MEFs) (Fig. 1A), indicating that the reduction in *Fgf7* expression was an indirect effect of *Med1* ablation.

## *FGF7 is expressed in BM mesenchymal cells but not in BM hematopoietic cells*

RT-qPCR (Fig. 1A) and western blot (Fig. 1B) analyses showed that FGF7 was expressed in BM mesenchymal cells, including stromal cells (MS-5 and OP-9) and osteoblasts (MC3T3-E1), as in  $Med1^{+/+}$  MEFs, but *Fgf7* mRNA was undetectable in hematopoietic cells (Fig. 1A). This implies that the quantity of

FGF7 secreted by stromal cells reflected FGF7 levels in the coculture niche model.

# <u>MEF FGF7 mediates mitogenic signal(s)</u> to BM cells and the support of long-term culture-initiating cells

We asked whether FGF7 is responsible for the phenotypes of  $Med1^{-/-}$  MEFs. To answer this question, we first analyzed the effect of FGF7 on the mitogenicity of cocultured BM cells. When BM cells on  $Med1^{+/+}$  MEFs were cocultured for 2 weeks in the presence of anti-FGF7 IgG or control rabbit whole IgG, the number of BM cells in cultures with anti-FGF7 IgG was lower than in the control cultures (Fig. 1C, supplementary Fig. 1A). This result, which could be due to a specific blocking effect of the antibody, prompted us to culture BM cells on  $Medl^{-/-}$  MEFs in the presence or absence of rhFGF7. As expected, the number of BM cells increased in FGF7 an dose-dependent manner (Fig. 1D. supplementary Fig. 1B).

To assess if changes in the cell number reflect altered mitogenicity, DNA synthesis was measured. After 1 week of coculture, the incorporation of BrdU into BM cells on  $Med1^{+/+}$  MEFs was lower in the presence of anti-FGF7 antibody than in the control, and the incorporation into BM cells on  $Med1^{-/-}$  MEFs was higher in the presence of rhFGF7 (Fig. 1E). Together, these data strongly suggest that FGF7 produced by MEFs had mitogenic activity on the cocultured BM cells.

We then conducted a long-term BM culture to assess whether FGF7 affected HSPC support. We first cultured BM cells on  $Med1^{+/+}$  MEFs in the presence of anti-FGF7 or control IgG for 8 weeks and counted the resulting long-term culture-initiating cells (LTC-ICs) by colony formation assay. The number of LTC-ICs was reduced in the presence of anti-FGF7 antibody, probably through its blocking effect on FGF7 secreted by MEFs (Fig. 1F). BM coculture on  $Med1^{-/-}$  MEFs in the absence or presence of rhFGF7 enhanced the number of LTC-ICs in an FGF7 dose-dependent manner (Fig. 1F). These data suggested a role for FGF7 in HSPC support in a MEF-based long-term BM culture.



Fig. 1. MEF-produced FGF7 promotes BM cell growth and supports LTC-ICs on MEFs.

(A and B) Expression of FGF7. Quantitative PCR (A) and western blot analysis (B). FGF7 was expressed in  $Med1^{+/+}$  MEFs, BM stromal cells (MS-5 and OP-9), and osteoblasts (MC3T3), and its expression was reduced in  $Med1^{-/-}$  MEFs (A, B). *Fgf*7 mRNA expression was not restored in Rev-*Med1*<sup>-/-</sup> MEFs and was not detected in BM hematopoietic cells (A). (C) The number of BM cells, when cocultured on  $Med1^{+/+}$  MEFs, decreased in the presence of anti-FGF7 antibody (Ab).  $\alpha$ , anti. (D) The number of BM cells, when cocultured on  $Med1^{-/-}$  MEFs, increased in the presence of rhFGF7. (E) BM cell DNA synthesis on  $Med1^{+/+}$  MEFs, measured by BrdU incorporation, decreased in the presence of anti-FGF7 Ab. BM cell DNA synthesis on  $Med1^{+/+}$  MEFs increased in the presence of rhFGF7. (F)  $Med1^{+/+}$  MEFs were reduced in the presence of anti-FGF7 Ab. The number of LTC-ICs on  $Med1^{-/-}$  MEFs increased in the presence of rhFGF7. The values are the mean  $\pm$  SD (A, C–E) or mean  $\pm$  SE (F) of a representative experiment performed in triplicate (A) or quadruplicate (C–F) (\*, P < 0.05; \*\*, P < 0.01).

## <u>BM stromal cell FGF7-mediated growth</u> of BM cells and LTC-IC support

Two independently established mouse BM stromal cell lines, MS-5 and OP-9, represent a BM niche and provide long-term HSPC support in vitro [14,17]. performed Therefore. we similar experiments using these cells. As expected, when BM cells were cocultured with MS-5 or OP-9 cells for 2 weeks, the addition of anti-FGF7 antibody attenuated the growth of the cocultured BM cells (Fig. 2A and supplementary Fig. 2A). Similarly, when BM cells were cocultured for 8 weeks, the number of LTC-ICs was reduced by the addition of anti-FGF7 antibody (Fig. 2Band supplementary Fig. 2B). These data further support the MEF-based conclusion that FGF7 has a role in BM cell growth and HSPC support in vitro.

# <u>BM stromal cell FGF7-mediated growth</u> of niche-dependent myeloblastic leukemia cells

MB-1 niche-dependent myeloblastoma cells, originally derived from a patient with myeloid blast crisis chronic myeloid leukemia, require BM stromal cells such as OP-9 and MS-5 cells for survival and optimal growth. During coculture, typically with OP-9 and MS-5 BM cells. **MB-1** cells form stromal cobblestone areas, which most probably reconstitute a stochastic model of leukemic stem or initiating cells in vitro [15,16]. We next asked if FGF7 produced by MS-5 or OP-9 BM stromal cells has a role in cocultured MB-1 cell growth. When anti-FGF7 antibody was added to MB-1 cells cocultured with MS-5 or OP-9 cells, the proliferation of MB-1 cells was significantly attenuated, although the cells were alive, as judged by trypan blue exclusion (Fig. 2C and D). Anti-FGF7 antibody also attenuated DNA synthesis in MB-1 cells cocultured with MS-5 or OP-9 cells (Fig. 2E, supplementary Fig. 3A). The number of cobblestone areas was also reduced significantly when anti-FGF7 antibody was added to the coculture, indicating reduced stem cell-like features of MB-1 cells (Fig. 2F, supplementary Fig. 3B). These results clearly indicate that BM stromal cell FGF7 is important for the optimal growth and support of stromal cell-dependent myeloblastic leukemia cells; it may constitute a leukemic stem cell niche.



Fig. 2. FGF7 depletion attenuates the growth of BM cells and MB-1 niche-dependent myeloblastoma cells and reduces the number of LTC-ICs cocultured on BM stromal cells.

(A, B) BM cells (A) and LTC-ICs (B) on MS-5 BM stromal cells were counted in the presence of anti-FGF7 Ab. The numbers of BM cells and LTC-ICs on stromal cells decreased in the presence of anti-FGF7 Ab. (C, D) The number of MB-1 cells on MS-5 (C) or OP-9 (D) BM stromal cells decreased in the presence of anti-FGF7 Ab. (E) BrdU incorporation into MB-1 cells cocultured with MS-5 cells was impaired in the presence of anti-FGF7 Ab. The values of BrdU incorporation are plotted as the fold increase versus the value with control IgG. (F) The number of cobblestone areas per visual field formed by MB-1 cells cocultured with MS-5 cells was counted. The number decreased in the presence of anti-FGF7 Ab. The values are the mean  $\pm$  SD (A, C, D) or SE (B, E, F) of a representative experiment performed in quadruplicate (\*, P < 0.05; \*\*, P < 0.01).

## FGF7 receptor FGFR2IIIb is expressed on BM mesenchymal cells but not on BM hematopoietic cells

We next investigated the mechanism of FGF7 in BM cell growth and HSPC support. First, we performed QT-PCR analysis to measure the expression of FGF receptors on BM hematopoietic cells and found that some FGFRs were variously expressed. However, FGFR2IIIb, the sole and specific receptor for FGF7, was completely undetectable (Fig. 3A).

Therefore, FGF7 may act on stromal cells in an autocrine manner. Various mesenchymal cells were checked for the expression of each FGFR. BM mesenchymal stromal cells (MS-5 and OP-9), osteoblastic cells (MC3T3-E1), and MEFs (Med1<sup>+/+</sup> and Med1<sup>-/-</sup>) expressed all of the FGFRs at various levels (Fig. 3A). The expression of FGFR2IIIb was also examined at the protein level by western blot and was comparable in all of these cells (Fig. 3B). The data suggest that FGF7 might act on BM stromal cells in an autocrine (or paracrine) manner to elicit niche function indirectly by these cells.

# FGF7 initiates FGFR-mediated intracellular signaling in MEFs and BM stromal cells

Because FGF7 appeared to act on stromal cells in the niche model system,

we asked if treating these cells with FGF7 elicited intracellular signaling. To this end, we assessed the phosphorylation of FRS2 (the immediate target of FGFR tyrosine kinase), the phosphorylation of MAP kinase (the intermediate hub of various intracellular signals), and the mRNA levels of immediate early response genes [8,9].

In both  $Med1^{+/+}$  and  $Med1^{-/-}$  MEFs, the phosphorylation of FRS2 and MAP kinases was upregulated 10 min after the addition of 10 or 50 ng/mL FGF7 (Fig. 4A), indicating that FGF7 effectively elicited FGFR2IIIb tyrosine kinase activity with subsequent activation of the signaling downstream cascade of We next analyzed FGFR2IIIb. the response of immediate early gene expression. In both  $Med1^{+/+}$  and  $Med1^{-/-}$ MEFs, c-Fos, c-Jun, and c-Myc were robustly transcribed after the addition of 10 ng/mL (supplementary Fig. 4) or 50 ng/mL (Fig. 4B) of FGF7, with a peak within 1 hr, followed by attenuation and a second wave of transcription (Fig. 4B, supplementary Fig. 4).

Experiments using MS-5 BM stromal cells yielded similar results, with prompt activation of FRS2 and MAP kinases and subsequent transcription of c-*Fos* and c-*Myc* (supplementary Fig. 5A and B). The phosphorylation of FRS2 and MAP kinases was also evident after the addition of FGF7 to OP-9 cells

(supplementary Fig. 5A). Taken together, these results clearly show that FGF7 effectively elicits FGFR2IIIb activation and subsequent intracellular signaling in MEFs and BM stromal cells and that MED1 is dispensable in these systems.



Fig. 3. Fgfr2IIIb is expressed in MEFs and BM stromal cells, but not in BM hematopoietic cells.

(A) Quantitative PCR. *Fgfr1*, *2IIIb*, *2c*, *3*, and *4* mRNAs were expressed in BM stromal cells (MS-5 and OP-9), osteoblasts (MC3T3-E1), and MEFs. BM hematopoietic cells expressed *Fgfr1*, *2c*, *3*, and *4* mRNAs but not *Fgfr2b* mRNA. The 23 values (mean  $\pm$  SD of a representative experiment performed in triplicate) are plotted as the fold increase versus the value in *Med1*<sup>+/+</sup> MEFs. (B) Western blot. FGF7 receptor FGFR2IIIb was expressed comparably in BM stromal cells, osteoblasts, and MEFs.



Fig. 4. FGF7 initiates FGFR2IIIb-mediated intracellular signaling in both *Med1<sup>+/+</sup>* and *Med1<sup>-/-</sup>* MEFs.
(A) The phosphorylation of FRS2 and MAP kinases was induced 10 min after rhFGF7 addition in MEFs. (B) The expression of immediate early genes, c-*Fos*, c-*Jun*, and c-*Myc*, was induced after rhFGF7 addition in MEFs.

## Discussion

We determined the role of FGF7 in a hematopoietic niche model. Our findings suggest that FGF7 and its cognate receptor FGFR2IIIb are specifically expressed in mesenchymal stromal cells, that FGF7 supports HSPC maintenance, and that this activity may be elicited indirectly by FGF7-activated stromal cells in an autocrine or paracrine manner. BM stromal cell FGF7 also constitutes a niche for myeloblastic leukemia cells.

FGF systems have been highlighted in recent years as important molecular components of microenvironmental signals both in hematopoietic (FGF1 and FGF2) and liver (FGF7) systems in stress conditions. In the liver niche, FGF7 produced by fibroblastic stroma-like cells acts on liver progenitor cells, thus constituting a genuine functional niche signal [12]. The FGF7-FGFR2IIIb system in the hematopoietic niche model and liver niche contrasts with the FGF1- and FGF2-elicited HSPC expansion in that, whereas FGF1 and FGF2 are supplied primarily by megakaryocytes, FGF7 is produced by stromal cells both in the hematopoietic niche (this study) and liver niche [12]. In addition, FGF7 acts on FGFR2IIIb on mesenchymal stromal cells in an autocrine or paracrine manner, whereas FGF1 and FGF2 act on cells distinct from their origin.

The mechanism of niche function by

FGF7 appears to be essentially indirect: it acts not on HSPCs (nor, most likely, on leukemic stem cells) but on stromal cells themselves, which subsequently act on normal and malignant HSPCs by an as yet unknown mechanism(s). A precedent is the indirect action of FGF7 on thymic mesenchymal cells to mediate thymocyte proliferation and T cell maturation [18]. The thymic mesenchymal cells express FGFR2IIIb, and these cells reportedly produce bone morphogenic protein 2 (BMP2), BMP4, Wnt5b, and Wnt10b, probably through an indirect mechanism downstream of transcription factors such as p53 and NF-kB [18]. It will be interesting to see if these molecules are altered in BM mesenchymal cells as well and if they contribute to HSPC support leukemic growth. a n d Although FGF7 is downregulated in *Med1*<sup>-/-</sup> MEFs, the inability to restore its expression by reintroducing FGF7 in the cells indicates that the effect of MED1 in this process is again indirect; it might reflect subtle differences in the differentiation status of Med1+/+ MEFs and Med1--- MEFs, reflected in their different manner of myogenic differentiation when MyoD is added [19]. In *Med1*<sup>-/-</sup> MEFs, we previously showed that niche molecules, angiopoietin-1, including Jagged-1, N-cadherin, Wnt, and BMP4, remain unchanged However, [7]. the transcription of genes encoding the

chemokines CXCL1, CXCL5, CXCL10, and CXCL15 is significantly reduced in Med1--- MEFs (GEO accession number GSE22471) [7]. These chemokines, whose functions in the BM microenvironment are essentially unknown, might contribute to the niche function uniquely for HSPC support and leukemic cell growth. The action of these chemokines on HSPCs in the presence or absence of FGF7 must be carefully examined in the future.

The BM microenvironment is complex in terms of cell types, molecules, oxygen concentration, and gravity, etc., which HSPC contribute to maintenance. proliferation, and differentiation. Therefore, the effect of niche molecules, especially those in families, such as FGFs, not be detected because may of redundancies within a living animal. Thus, a simplified niche model system, such as the one adopted in this study, is an ideal strategy for assaying molecules of interest. However, having established FGF7 as a potentially important (both normal and malignant) niche factor in vitro, the next step is to determine whether it has the same role in BM mesenchymal stromal cells in vivo. A series of mouse knockout models, with single or double ablation of genes including Fgf7 or Fgfr2IIIb, as well as transgenic models, will help answer this question in the future.

## Acknowledgements

I thank my collaborators, Kaori Minami, Satowa Tanaka, Mami Nagai, Keiji Matsui, Natsumi Hasegawa, Robert G. Roeder, Shigetaka Asano and Mitsuhiro Ito. We thank K. Ito for MS-5 cells, H. Kaji for MC3T3-E1 cells, S. Yamagishi and Y. Ikebata for technical assistance, and Y. Takai and members in our laboratories and Takai's laboratory for helpful discussion. This study was supported by grants from the MEXT, the Global Center for Excellence Program Global Center of Excellence for Education and Research on Signal Transduction Medicine in the Coming Generation from MEXT, and the Takeda Science Foundation (to M.I.).

### References

- P.S. Frenette, S. Pinho, D. Lucas, C. Scheiermann, Mesenchymal stem cell: keystone of the hematopoietic stem cell niche and a stepping-stone for regenerative medicine, Annu. Rev. Immunol. 31 (2013) 285-316.
- [2] J.N. Smith, L.M. Calvi, Current concepts in bone marrow microenvironmental regulation of hematopoietic stem and progenitor cells, Stem Cells 31 (2013) 1044-1050.
- [3] A. Nakamura-Ishizu, T. Suda,

Hematopoietic stem cell niche: an interplay among a repertoire of multiple functional niches, Biochim. Biophys. Acta. 1830 (2013) 2404-2409.

- [4] A. Greenbaum, Y.M. Hsu, R.B. Day, L.G. Schuettpelz, M.J. Christopher, J.N. Borgerding, T. Nagasawa, D.C. Link, CXCL12 in early mesenchymal progenitors is required for haematopoietic stem-cell maintenance, Nature 495 (2013) 227-230.
- [5] L. Ding, S.J. Morrison, Haematopoietic stem cells and early lymphoid progenitors occupy distinct bone marrow niches, Nature 495 (2013) 231-235.
- [6] M. Corselli, C.J. Chin, C. Parekh,
  A. Sahaghian, W. Wang, S. Ge, D.
  Evseenko, X. Wang, E. Montelatici,
  L. Lazzari, G.M. Crooks, B. Péault,
  Perivascular support of human
  hematopoietic stem/progenitor
  cells, Blood 121 (2013)
  2891-2901.
- A. Sumitomo, R. Ishino, [7] N. Urahama, K. Inoue, K. Yonezawa, N. Hasegawa, O. Horie, H. Matsuoka, T. Kondo, R.G. Roeder, M. Ito. The transcriptional mediator subunit MED1/TRAP220 in stromal cells is involved in hematopoietic stem/progenitor cell support through osteopontin

expression, Mol. Cell. Biol. 30 (2010) 4818-4827.

- [8] A. Beenken, M. Mohammadi, The FGF family: biology, pathophysiology and therapy, Nat. Rev. Drug. Discov. 8 (2009) 235-253.
- [9] V. Knights, S.J. Cook, De-regulated FGF receptors as therapeutic targets in cancer, Pharmacol. Ther. 125 (2010) 105-117.
- [10] M. Zhao, J.T. Ross, T. Itkin, J.M. Perry, A. Venkatraman, J.S. Haug, M.J. Hembree, C.X. Deng, T. Lapidot, X.C. He, L. Li, FGF signaling facilitates postinjury recovery of mouse hematopoietic system, Blood 120 (2012) 1831-1842.
- [11] T. Itkin, A. Ludin, B. Gradus, S. Gur-Cohen, A. Kalinkovich, A. Schajnovitz, Y. Ovadya, O. Kollet, J. Canaani, E. Shezen, D.J. Coffin, G.N. Enikolopov, T. Berg, W. Piacibello, E. Hornstein, T. Lapidot, FGF-2 expands murine hematopoietic stem and progenitor cells via proliferation of stromal c-Kit activation. cells. and CXCL12 down-regulation, Blood 120 (2012) 1843-1855.
- [12] H.M. Takase, T. Itoh, S. Ino, T. Wang, T. Koji, S. Akira, Y. Takikawa, A. Miyajima, FGF7 is a

functional niche signal required for stimulation of adult liver progenitor cells that support liver regeneration, Genes Dev. 27 (2013) 169-181.

- [13] S. Malik, R.G. Roeder, The metazoan Mediator co-activator complex as an integrative hub for transcriptional regulation, Nat. Rev. Genet. 11 (2010) 761-772.
- [14] K. Itoh, H. Tezuka, H. Sakoda, M. Konno, K. Nagata, T. Uchiyama, H. Uchino, K.J. Mori, Reproducible establishment of hemopoietic supportive stromal cell lines from murine bone marrow, Exp. Hematol. 17 (1989) 145-153.
- [15] K. Funayama, Y. Saito-Kurimoto,
  Y. Ebihara, M. Shimane, H.
  Nomura, K. Tsuji, S. Asano,
  Adhesion-mediated self-renewal abilities of Ph+ blastoma cells,
  Biochem. Biophys. Res. Commun. 396 (2010) 193-198.
- [16] K. Funayama, M. Shimane, H. Nomura, S. Asano, An evidence for adhesion-mediated acquisition of acute myeloid leukemic stem cell-like immaturities, Biochem. Biophys. Res. Commun. 392 (2010) 271-276.
- [17] P. Feugier, N. Li, D. Y. Jo, J. H. Shieh, K.L. MacKenzie, J.F. Lesesve, V. Latger-Cannard, D. Bensoussan, R.G. Crystal, S. Rafii,

J.F. Stoltz, M.A. Moore, Osteopetrotic mouse stroma with thrombopoietin, c-kit ligand, and flk-2 ligand supports long-term mobilized CD34+ hematopoiesis in vitro, Stem Cells Dev. 14 (2005) 505-516.

- [18] S.W. Rossi, L.T. Jeker, T. Ueno, S. Kuse, M.P. Keller, S. Zuklys, A.V. Gudkov, Y. Takahama, W. Krenger, B.R. Blazer, G.A. Holländer, Keratinocyte growth factor (KGF) enhances postnatal T-cell development via enhancements in proliferation and function of thymic epithelial cells, Blood 109 (2007) 3803-3811.
- [19] K. Ge, M. Guermah, C.-X. Yuan, M. Ito, A.E. Wallberg, B.M. Spiegelman, R.G. Roeder, Transcription coactivator TRAP220 is required for PPARγ2-stimulated adipogenesis, Nature 417 (2002) 563-567.

## **Supplementary Figures**



Supplementary Fig. 1. FGF7 mediates the growth of BM cells on MEFs.

The number of BM cells decreased after the addition of anti-FGF7 antibody in cocultures with  $Med1^{+/+}$  MEFs (upper panel) and increased after the addition of recombinant human (rh) FGF7 in cocultures with  $Med1^{-/-}$  MEFs (lower panel). Scale bar, 100 µm.



**Supplementary Fig. 2. FGF7 depletion attenuates the growth of BM cells and the number of LTC-ICs** cocultured on OP-9 BM stromal cells. (A, B) BM cells (A) and LTC-ICs (B) on OP-9 BM stromal cells were counted in the presence of anti-FGF7 Ab. The numbers of BM cells and LTC-ICs on stromal cells decreased in the presence of anti-FGF7 Ab.



Supplementary Fig. 3. DNA synthesis and cobblestone formation of MB-1 cells cocultured with OP-9 BM stromal cells.

(A) BrdU incorporation into MB-1 cells cocultured with OP-9 cells was impaired in the presence of anti-FGF7 Ab. The values of BrdU incorporation are plotted as the fold increase versus the value with control IgG. (B) The number of cobblestone areas per visual field formed by MB-1 cells cocultured with OP-9 cells was counted. The number decreased in the presence of anti-FGF7 Ab. The values are the mean  $\pm$ SD of a representative experiment performed in quadruplicate (\*, *P* < 0.05; \*\*, *P* < 0.01).



**Supplementary Fig. 4.** FGF7 initiates FGFR2IIIb-mediated intracellular signaling in MEFs. The expression of immediate early genes, c-*Fos*, c-*Jun* and c-*Myc*, was induced after the addition of 10 ng/mL rhFGF7 in both *Med1*<sup>+/+</sup> and *Med1*<sup>-/-</sup> MEFs.



Supplementary Fig. 5. FGF7 initiates FGFR2IIIb-mediated intracellular signaling in BM stromal cells. (A) The phosphorylation of FRS2 and MAP kinases was induced 10 min after rhFGF7 addition in OP-9 and MS-5 BM stromal cells. (B) The expression of immediate early genes, c-*Fos* and c-*Myc*, was induced after rhFGF7 addition in MS-5 BM stromal cells.