



Relationship between activities of daily living and readmission within 90 days in hospitalized elderly patients with heart failure

Kitamura, Masahiro

(Degree)

博士（保健学）

(Date of Degree)

2018-03-25

(Date of Publication)

2019-03-01

(Resource Type)

doctoral thesis

(Report Number)

甲第7173号

(URL)

<https://hdl.handle.net/20.500.14094/D1007173>

※ 当コンテンツは神戸大学の学術成果です。無断複製・不正使用等を禁じます。著作権法で認められている範囲内で、適切にご利用ください。



博 士 論 文

Relationship between activities of daily living and readmission
within 90 days in hospitalized elderly patients with heart failure

(入院高齢心不全患者における ADL と 90 日以内再入院の関連性)

平成 3 0 年 1 月 1 7 日

神戸大学大学院保健学研究科保健学専攻

北村 匡大

(Masahiro Kitamura)

1. Introduction

Heart failure (HF) affects about 1% of individuals in their 50s and 10% of those in their 80s, and its incidence is increasing rapidly with age worldwide [1]. Among individuals with HF, problems such as an increase in the readmission rate and medical expenses, and many readmissions in the short-term occur [2, 3]. In epidemiological studies of Japanese elderly patients with HF, complications, an increase in the length of hospital stay and medical expenses, high readmission rates, and increased medical expenses have been reported [4]. Among patients who are readmitted, there are increased cases of disease onset other than heart disease [5]. The readmission rate is high within 6 months [3, 6]. Age, severity, the length of hospital stay, comorbidities, and disease management are risk factors for short-term readmission [7-10].

One of the purposes of rehabilitation in patients with HF is the recovery of activities of daily living (ADL) [11]. ADL and functional limitations in patients with HF are associated with readmission [12, 13]. Few studies have shown the relationship between readmission and ADL in patients with HF, and the cutoff values of ADL to predict readmission are unknown.

Therefore, we assumed that in patients with HF, the group with poor ADL would have a higher readmission rate than the group with good ADL. The purpose of the present study was to investigate the relationship between ADL and readmission within 90 days in elderly patients with HF.

2. Materials and Methods

2. 1. Study Design and Participants

Five hundred eight-nine consecutive patients with HF who underwent rehabilitation at one acute care hospital from May 2012 to May 2016 were included in this retrospective cohort study. Of these patients, those aged ≥ 65 years and those who could walk with assistance before hospitalization and during the initial hospitalization were included. Patients who underwent pacemaker operation during hospitalization, those who were transferred to other departments, those who were not discharged home, those who died during hospitalization, and those who were difficult to follow for 90 days were excluded from this study. The reason for exclusion of pacemaker patients is because the rehabilitation protocol is different.

The Kokura Rehabilitation College Institutional Review Committee on Human Research approved this study (approval no. 29-03), and informed consent was obtained from each patient.

2. 2. Rehabilitation of hospitalization

Patients of this study, has received the rehabilitation in accordance with the Japanese guidelines [11]. About the start of rehabilitation, after confirming with the doctor that there are no symptoms at light activity, we have urged rehabilitation such as the sitting position, standing, walking and ADL. If the patient was able to walk, we have urged the aerobic

exercise for the purpose of increasing the endurance necessary to home life.

2. 3. Investigation

Patients' characteristics and clinical parameters, including age, gender, body mass index (BMI), left ventricular ejection fraction (LVEF), brain natriuretic peptide (BNP) concentration, New York Heart Association (NYHA) class at discharge, estimated glomerular filtration rate (eGFR), creatinine level at discharge, hemoglobin level at discharge, albumin level at discharge, acute management, comorbidity, Charlson comorbidity index, medications, time of initiation of walking exercise, length of hospital stay, motor Functional Independence Measure (FIM) score at discharge, and cognitive FIM score at discharge, were evaluated by reviewing medical records retrospectively. We divided patients into two groups, the non-readmission or readmission group within 90 days, based on a previous study [9, 10]. We also evaluated the FIM as a measurement of ADL [14].

2. 4. Assessment of ADL

The FIM was developed to suit rehabilitative aspects of patients with disabilities, and it consisted of two domains: motor and cognitive [14]. The motor domain (motor FIM) consists of 13 items: eating; grooming; bathing; dressing upper body; dressing lower body; toileting; bladder management; bowel management; transfer to bed, chair, or wheelchair; transfer to toilet; transfer to tub or shower; walking/wheelchair; and stairs. The cognitive domain (cognitive FIM) consists of 5 items: comprehension, expression, social interaction, problem

solving, and memory. A scoring scale from 1 to 7 points was used (1 point for total assistance, 2 points for maximal assistance, 3 points for moderate assistance, 4 points for minimal contact assistance, 5 points for supervision, 6 points for modified independence, and 7 points for complete independence). The minimum total FIM score was 18 points, and the maximum total FIM score was 126 points; whereas, the minimum scores for the motor FIM and cognitive FIM were 13 points and 5 points, and maximum scores for the motor FIM and cognitive FIM were 91 points and 35 points, respectively. This measurement was obtained by two physical therapists from the time of discharge.

2. 5. Assessment of follow-up

Patients enrolled in this study were followed up with for 90 days. The first follow-up clinic visit was scheduled within the first 2 weeks after discharge. The following readmission information was obtained from medical records by two physical therapists: the date of readmission, number of days from discharge to readmission, and reasons for readmission. The definition of readmission was admission for all causes within 90 days after discharge, except hospitalization for examination.

2. 6. Statistical Analysis

Patients' characteristics and clinical parameters were reported using percentages for categorical variables and the mean \pm standard deviation for continuous variables. The unpaired t-test and chi-square test were used to compare patients' characteristics and clinical

parameters between the two groups. A Cox proportional hazard model for readmission within 90 days was used to ascertain whether ADL at discharge was an independent predictor of readmission within 90 days (hazard ratio and 95% confidence interval). The objective variables used in this model were readmission (the end point), data 0 (non-readmission), and data 1 (readmission). The explanatory variables used in this model were variables that showed statistical significance at $p < 0.05$ in univariate analysis. The detailed items between two groups on factors which were significant by these analyzes were examined. To determine the cut-off value of the most influential factor obtained by these analyzes, a receiver operating characteristic (ROC) curve was constructed by plotting the sensitivity against the false positive rate. Patients were classified into two groups according to these cut-off values, a Kaplan-Meier curve was constructed, and a log-rank test was used. A p -value < 0.05 indicated statistical significance. Statistical analyses were performed using IBM SPSS 23.0 J statistical software (IBM SPSS Japan, Inc., Tokyo, Japan).

3. Results

3. 1. Flow of included patients

A flow chart of patients included in this study is shown in Figure 1. Of 589 consecutive patients with HF who underwent rehabilitation, 323 met the inclusion criteria, but 210 patients were excluded later because of pacemaker operation during hospitalization (14),

transfer to other departments (8), non-home discharge (78), died during hospitalization (17) or were difficult to follow for 90 days (93 patients). Therefore, 113 patients were ultimately included and divided into the non-readmission group (n = 90) or readmission group (n = 23).

3. 2. Patients' characteristics

A comparison of the patients' clinical characteristics between the non-readmission group and readmission group is shown in Table 1. Compared to the non-readmission group, the readmission group was significantly older and had a lower BMI, poorer NYHA class, lower hemoglobin level at discharge, and lower motor FIM score ($p < 0.05$).

3. 3. Factor of readmission

Results of the Cox proportional hazard models, as provided in Table 2, demonstrate the associations between each parameter and readmission within 90 days. In the univariate Cox proportional hazard model with age, BMI, NYHA class at discharge, hemoglobin level at discharge, and motor FIM score at discharge as covariates, all these variables were independent predictors of readmission. In the multivariate cox proportional hazard model with age, BMI, NYHA class at discharge, hemoglobin level at discharge, and motor FIM score at discharge as covariates, BMI (hazard ratio: 0.87; 95% confidence interval: 0.76-0.99) and motor FIM score at discharge (hazard ratio: 0.94; 95% confidence interval: 0.89-0.99) were independent predictors of readmission (Table 3). Comparison between groups of motor FIM items is shown in Table 4.

3. 4. Cut-off value of the motor FIM score for predicting readmission

The cut-off value of the motor FIM score at discharge that predicted the occurrence of readmission in the ROC curve was 75 points (area under the curve: 0.78, $p < 0.001$, sensitivity: 0.767, false positive rate: 0.348) (Figure 2).

3. 5. Readmission rates based on the motor FIM score

In the Kaplan-Meier analysis, we divided patients into two groups based on the cutoff values of the motor FIM score. The group with a motor FIM score ≥ 75 points had significantly higher readmission avoidance rates than the group with a motor FIM score < 75 points (log-rank test, $p < 0.001$) (Figure 3).

4. Discussion

To our knowledge, this is the first study to report the differences in motor ADL in elderly hospitalized patients with HF that are associated with readmission within 90 days.

4. 1. Characteristics of the readmission group of elderly patients with HF

The elderly patients with HF in the readmission group were significantly older and a poorer NYHA class, lower hemoglobin level, and lower motor FIM score than those in the non-readmission group. These findings were largely in agreement with the characteristics of patients with HF who were readmitted in previous studies. In a past study, an older age and low BMI in patients with HF were risk factors for short-term readmission [15]. Regarding the BMI of patients with HF, a low BMI is known to reduce readmission [16]. Additionally, the

poor NYHA class is associated with readmission within 90 days [17]. Anemia in patients with HF is a predictor of readmission within 90 days [9, 18], and their low hemoglobin levels are likely to result in readmission because of heart load [19,20]. Additionally, the low ADL in patients with HF is associated with readmission within 30 days [21]. However, the readmission rate (20.4%) within 90 days in this study is lower than that reported in these aforementioned previous studies. The subjects of this study included walkable and initial hospitalization and excluded non-home discharge. From these criteria, there were many HF patients in good condition and the readmission rate may be low. Therefore, in our study, although subjects' readmission rate was low, the characteristics of patients with HF are almost consistent with those of previous studies; thus, these patient characteristics are considered partially generalizable.

4. 2. Relationship between readmission and motor ADL

In patients with HF who were readmitted because of poor ADL recovery during hospitalization and declining ADL after discharge, new events may occur from increased heart load. Low ADL at discharge in patients with internal disorders is associated with a high readmission rate, and change in the rate of ADL during hospitalization is related to readmission. Intervention to prevent ADL decline during those hospitalizations is important [22]. Patients with HF after discharge are likely to decline in physical function, with the possibility of readmission due to events such as disease, falls, etc. [23,24]. In addition, in

patients with a disability who underwent rehabilitation during hospitalization, a low ADL at discharge was associated with a high rate of readmission within 90 days, and HF was a risk factor of complications [25]. Therefore, in patients with hospitalization, physicians need to further research of ADL during hospitalization and the status of ADL after discharge. In readmission group, motor FIM low items were self-care, transfer and locomotion. Walking is known as readmission factor in HF patients [26]. Self-care is reported as a prognostic factor in elderly hospitalized patients [27]. Improvements in items such as self-care, transfer and locomotion shown in this study may prevent readmission. It is important to investigate the relationship between physical function and ADL in order to prevent readmission in the future.

4. 3. Clinical implication

Motor ADL was an independent factor of readmission within 90 days in elderly patients with HF. Improvement of ADL at discharge may reduce readmission. The cut-off value of the motor FIM score may be an indicator for readmission. These findings suggest the importance of intervention to improve ADL during hospitalization and after discharge.

4. 4. Limitations

This retrospective cohort study was conducted at one facility with a small sample.

According to inclusion and exclusion criteria, the final subjects was approximately 20%. The motor FIM may have a ceiling effect [28]. Also, not consider gender difference in this study.

Additionally, we did not investigate physical function [12, 29], and we were unable to follow-

up with some patients, and to examine clinical characteristics and ADL for non-home discharge patients and home discharge cases. Further, we did not assess outpatient rehabilitation after discharge.

5. Conclusion

The motor ADL score in elderly patients with HF was an independent factor of readmission, and its cut-off value was 74.5 points.

Acknowledgment

This work was supported by JSPS KAKENHI grant number JP17K01500.

Compliance with ethical standards

Participants received oral explanations of the research procedures from the researchers.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this article.

Ethical approval

This study was approved by the Kokura Rehabilitation College Institutional Committee.

Informed consent

Informed consent was obtained from each patient.

References

1. W. B. Kannel, A. J. Belanger, "Epidemiology of heart failure," *Am Heart J*, vol. 121, no. 4 Pt 1, pp. 951–957, 1991.
2. T. E. Owan, D. O. Hodge, R. M. Herges, et al., "Trends in prevalence and outcome of heart failure with preserved ejection fraction," *N Engl J Med*, vol. 355, no. 3, pp. 251–259, 2006.
3. S. F. Jencks, M. V. Williams, E. A. Coleman, "Rehospitalizations among patients in the Medicare fee-for-service program," *N Engl J Med*, vol. 360, no. 14, pp. 1418–1428, 2009.
4. H. Tsutsui, M. Tsuchihashi-Makaya, S. Kinugawa, et al., "Clinical characteristics and outcome of hospitalized patients with heart failure in Japan," *Circ J* vol. 70, no. 12, pp. 1617–1623, 2006.
5. K. Dharmarajan, K. M. Strait, T. Lagu, et al., "Acute decompensated heart failure is routinely treated as a cardiopulmonary syndrome," *PLoS One*, vol. 8, no. 10, pp. e78222, 2013.
6. A. S. Desai, L. W. Stevenson, "Rehospitalization for heart failure: predict or prevent?"

- Circulation, vol. 126, no. 4, pp. 501-506, 2012.
7. K. Dharmarajan, A. F. Hsieh, Z. Lin, et al., “Diagnoses and timing of 30-day readmissions after hospitalization for heart failure, acute myocardial infarction, or pneumonia,” JAMA, vol. 309, no. 4, pp. 355–363, 2013.
 8. J. T. Black, P. S. Romano, B. Sadeghi, et al., “A remote monitoring and telephone nurse coaching intervention to reduce readmissions among patients with heart failure: study protocol for the Better Effectiveness After Transition - Heart Failure (BEAT-HF) randomized controlled trial,” Trials vol. 15, pp. 124–124, 2014.
 9. M. Saito, K. Negishi, T. H. Marwick, “Meta-analysis of risks for short-term readmission in patients with heart failure,” Am J Cardiol, vol. 117, no. 4, pp. 626–632, 2016.
 10. H. Deeka, H. Skouri, S. Nouredine, “Readmission rates and related factors in heart failure patients: a study in Lebanon,” Collegian, vol. 23, no. 1, pp. 61–68, 2016.
 11. Japanese Circulation Society Joint Working Group, “Guidelines for rehabilitation in patients with cardiovascular disease 2012,” Circ J, vol. 78, no. 8, pp. 2022-2093, 2014.
 12. S. Yamada, Y. Shimizu, M. Suzuki, et al., “Functional limitations predict the risk of rehospitalization among patients with chronic heart failure,” Circ J, vol. 76, no. 7, pp. 1654–1661, 2012.
 13. S. M. Dunlay, S. M. Manemann, A. M. Chamberlain, et al., “Activities of daily living and outcomes in heart failure,” Circ Heart Fail, vol. 8, no. 2, pp. 261–267, 2015.

14. K. J. Ottenbacher, Y. Hsu, C. V. Granger, et al., "The reliability of the functional independence measure: a quantitative review," *Arch Phys Med Rehabil*, vol. 77, no. 12, pp. 1226–1232, 1996.
15. H. Aizawa, S. Imai, K. Fushimi, "Factors associated with 30-day readmission of patients with heart failure from a Japanese administrative database," *BMC Cardiovasc Disord*, vol. 134, pp. 134, 2015.
16. A. Zapatero, R. Barba, N. Gonzalez, et al., "Influence of obesity and malnutrition on acute heart failure," *Rev Esp Cardiol (Engl Ed)*, vol. 65, no. 5, pp. 421–426, 2012.
17. E. Zaharias, J. Cataldo, L. Mackin, et al., "Simple measures of function and symptoms in hospitalized heart failure patients predict short-term cardiac event-free survival," *Nurs Res Pract*, vol. 2014, pp. 815984, 2014.
18. S. Muzzarelli, G. Leibundgut, M. T. Maeder, et al., "Predictors of early readmission or death in elderly patients with heart failure," *Am Heart J*, vol. 160, no. 2, pp. 308–314, 2010.
19. S. Hamaguchi, M. Tsuchihashi-Makaya, S. Kinugawa, et al., "Anemia is an independent predictor of long-term adverse outcomes in patients hospitalized with heart failure in Japan. A report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD)," *Circ J* vol. 73, no. 10, pp. 1901–1908, 2009.
20. M. A. Varat, R. J. Adolph, N. O. Fowler, "Cardiovascular effects of anemia," *Am Heart J*,

vol. 83, no. 3, pp. 415–426, 1972.

21. S. R. Greysen, I. Stijacic Cenzer, A. D. Auerbach, et al., “Functional impairment and hospital readmission in Medicare seniors,” *JAMA Intern Med*, vol. 175, no. 4, pp. 559–565, 2015.
22. O. Tonkikh, E. Shadmi, N. Flaks-Manov, et al., “Functional status before and during acute hospitalization and readmission risk identification,” *J Hosp Med*, vol. 11, no. 9, pp. 636–641, 2016.
23. H. M. Krumholz, “Post-hospital syndrome--an acquired, transient condition of generalized risk,” *N Engl J Med*, vol. 368, no. 2, pp. 100–102, 2013.
24. C. Rodríguez-Pascual, E. Paredes-Galán, A. I. Ferrero-Martínez, et al., “The frailty syndrome is associated with adverse health outcomes in very old patients with stable heart failure: a prospective study in six Spanish hospitals,” *Int J Cardiol*, vol. 236, no. 1, pp. 296–303, 2017.
25. R. V. Galloway, A. M. Karmarkar, J. E. Graham, et al., “Hospital readmission following discharge from inpatient rehabilitation for older adults with debility,” *Phys Ther*, vol. 96, no. 2, pp. 241–251, 2016.
26. M. T. Vidán, V. Blaya-Novakova, E. Sánchez, et al., “Prevalence and prognostic impact of frailty and its components in non-dependent elderly patients with heart failure.” *Eur J Heart Fail*, Vol. 18, no. 7, 869-875, 2016.

27. D.E. Barnes, K. M. Mehta, W. J. Boscardin, et al., "Prediction of recovery, dependence or death in elders who become disabled during hospitalization." *J Gen Intern Med*, Vol. 28, no. 2, 261-268, 2013.
28. S. Kwon, A. G. Hartzema, P. W. Duncan, et al., "Disability measures in stroke: Relationship among the Barthel index, the functional independence measure, and the modified Rankin scale," *Stroke*, vol. 35, no. 4, 918–923, 2004.
29. G. Pulignano, D. Del Sindaco, A. Di Lenarda, et al., "Usefulness of frailty profile for targeting older heart failure patients in disease management programs: a cost-effectiveness, pilot study," *J Cardiovasc Med (Hagerstown)*, vol. 11, no. 10, pp. 739–747, 2010.

Table 1. Patients' characteristics

	Non- readmission n=90	Readmission n=23	F or χ^2 value	p Value
Age, years	79.6±6.9	83.8±5.9	1.12 ^a	0.008
Gender, male, %	55.6	47.8	0.44	0.51
BMI, kg/m ²	22.8±3.0	21.3±3.5	0.89 ^a	0.04
Clinical parameter				
LVEF, %	47.2±16.6	49.4±13.5	2.13 ^a	0.55
BNP, pg/mL	783.8±826.4	696.2±410.2	2.83 ^a	0.62
NYHA class I/II, %	84.4/15.6	65.2/34.8	4.32	0.04
Creatinine level, mg/dL	1.3±0.9	1.9±1.7	9.43	0.16
eGFR, ml/min/1.73 m ²	50.1±21.8	40.3±24.7	0.92 ^a	0.06
Hemoglobin level, g/dL	11.5±2.0	10.4±2.2	0.13 ^a	0.04
Albumin level at discharge, g/dL	3.5±0.6	3.4±0.4	0.14	0.35
Acute management, %	18.9	8.7	1.36	0.24
Comorbidity, %				

Hypertension	86.7	82.6	0.25	0.62
Diabetes	44.4	30.4	1.48	0.22
Ischemic heart disease	51.1	47.8	0.08	0.78
Valvular disease	27.8	43.5	2.11	0.15
Atrial fibrillation	41.1	56.5	1.76	0.18
Orthopedic disease	37.8	34.8	0.07	0.79
Neurological disease	24.0	5.6	3.04	0.08
Respiratory disease	21.1	21.7	0.004	0.95
CCI	2.2±1.9	2.8±2.3	1.82	0.45
Medication				
Diuretic	93.3	95.7	0.17	0.68
β-blockers	58.9	52.2	0.34	0.56
ACEI/ARB	40.0	47.8	0.46	0.50
Rehabilitation progress				
Initiation of walking exercise, days	4.6±4.9	6.2±7.1	3.15 ^a	0.21
Length of hospital stay, days	17.3±7.4	15.9±7.1	0.23 ^a	0.43
Motor FIM score at admission	39.4±18.0	34.8±15.2	0.85 ^a	0.26
Motor FIM score at discharge	79.8±8.1	70.9±9.5	0.94 ^a	<0.001

Cognitive FIM score at admission	29.6±7.0	26.5±8.2	0.61 ^a	0.07
Cognitive FIM score at discharge	33.0±3.9	31.2±5.2	4.78 ^a	0.08

Values are presented as a mean±standard deviation or %.

ACEI = angiotensin-converting enzyme inhibitor; ARB = angiotensin receptor blocker; BMI = body mass index; BNP = brain natriuretic peptide; CCI = Charlson Comorbidity Index; eGFR = estimated glomerular filtration rate; FIM = Functional Independence Measurement; LVEF = left ventricular ejection fraction; NYHA = New York Heart Association.

^a F value.

1 **Table 2.** Results of univariate analysis

	Cox	95% CI		p
	proportional			value
	hazard ratio			
Age, years	1.09	1.02	1.17	0.01
BMI, kg/m ²	0.86	0.75	0.98	0.03
NYHA class at discharge	2.53	1.07	5.96	0.03
I/II, %				
Hemoglobin level at	0.76	0.61	0.95	0.02
discharge, g/dL				
m-FIM score at discharge	0.92	0.89	0.96	<0.001

2 BMI = body mass index; CI = confidence interval; FIM = Functional Independence

3 Measurement; NYHA = New York Heart Association.

4 **Table 3.** Results of multivariate analysis

	Cox	95% CI		p
	proportional			value
	hazard ratio			
Age, years	1.02	0.94	1.10	0.70
BMI, kg/m ²	0.87	0.76	0.99	0.047
NYHA class at discharge	1.52	0.61	3.77	0.28
I/II, %				
Hemoglobin level at	0.88	0.70	1.11	0.37
discharge, g/dL				
m-FIM score at discharge	0.94	0.89	0.99	0.012

5 BMI = body mass index; CI = confidence interval; FIM = Functional Independence

6 Measurement; NYHA = New York Heart Association.

Table 4. Motor FIM item between groups

	Non- readmission n=90	Readmission n=23	F or χ^2 value	p Value
Eating	6.8±0.6	6.7±0.5	0.46 ^a	0.430
Grooming	6.6±0.8	6.0±1.0	1.39 ^a	0.004
Bathing	6.2±1.1	5.2±1.2	0.04 ^a	0.001
Dressing upper body	6.5±0.8	5.7±1.0	0.36 ^a	<0.001
Dressing lower body	6.5±0.9	5.6±1.1	1.14 ^a	<0.001
Toileting	6.5±0.8	5.8±1.2	3.42 ^a	0.002
Bladder management	6.8±0.6	6.5±0.8	8.74 ^a	0.100
Bowel management	6.8±0.6	6.4±1.1	21.09 ^a	0.060
Transfer of Bed/ chair/ wheelchair	6.7±0.6	6.4±0.7	1.96 ^a	0.044
Transfer of toilet	6.6±0.7	6.1±0.9	0.54 ^a	0.005
Transfer of tub/ shower	5.2±1.7	4.2±1.5	1.10 ^a	0.012
Walking/ wheelchair	6.3±0.9	5.6±1.3	0.60 ^a	0.007
Stairs	2.4±2.1	1.4±1.2	17.55 ^a	0.005

Values are presented as a mean±standard deviation.

^a F value.




Figure1. Patient flow during study




Figure2. Cutoff value of the motor FIM score for predicting readmission




Figure3. Comparison of readmission rates according to the FIM